

## **Assessment of heavy metals concentrations in water, plankton and fish of Lake Manzala, Egypt**

**Mohamed H. Bahnasawy, Abdel Aziz A. Khidr , Nadia A. Dheina**

Zoology Department, Damietta Faculty of Science, Damietta, Egypt

### **ABSTRACT**

The levels of some heavy metals (Cu, Zn, Cd, Pb) were determined in water, plankton and fish (*Liza aurata*) collected from five sites in Lake Manzala. Metals in water and fish exhibited a significant seasonal and regional variations in which all metals attained their maximum values during summer, while the lowest level was reported during winter. The accumulation of different metals in water, plankton and fish tissues followed the order Zn > Cu > Pb > Cd. The mean concentrations of the tested metals in water were: Cu (0.055), Zn (0.311), Cd (0.020) and Pb (0.022) mg/l. Cd level in water was found to be higher than the permissible limit recommended for drinking water. Metals in plankton were much higher than those in water and fish. Gills of the examined fish contained the highest concentrations of all the measured metals, while muscles retained the lowest levels. In spite of the contamination of Lake Manzala by such heavy metals, the levels of these metals in the edible fish muscle did not exceed the recommended permissible limits and thus are considered safe for human consumption.

**Key words:** Lake Manzala, heavy metals, water, plankton, fish.

### **INTRODUCTION**

In aquatic systems, heavy metals have received considerable attention due to their toxicity and accumulation in biota (Mason, 1991). Metals generally enter the aquatic environment through atmospheric deposition, erosion of geological matrix or due to anthropogenic activities caused by industrial effluents, domestic sewage and mining wastes (Tarvainen *et al.* 1997; Stephen *et al.*, 2000). Some of these metals are toxic to living organisms even at quite low concentrations, whereas others are biologically essential as natural constituents of aquatic ecosystem and only become toxic at very high concentrations.

Heavy metals may affect organisms directly by accumulating in the body or indirectly by transferring to the next trophic level of the food chain. Being non-biodegradable like many organic pollutants, they can be concentrated along the food chain, producing their toxic effects at points often far away from the source of pollution (Fernandez *et al.*, 2000). Accumulation of heavy metals in the food web can occur either by accumulation from the surrounding medium such as water or sediment, or by bioaccumulation from the food source (Tulonen *et al.*, 2006). Aquatic organisms have been widely used in biological monitoring and assessment of safe environmental levels of heavy metals.

Lake Manzala is one of the large lakes in northern region of Egypt (about 52611 hectares surface area) and the most productive for fisheries. The lake receives heavy loads of organic and inorganic pollutants via several agricultural drains (Badawy *et al.*, 1995). Due to the toxicity of heavy metals, accurate information about their concentration in aquatic ecosystem is needed (Janssen *et al.*, 2000). Therefore, the objective of this study was to evaluate the pollution level of Lake Manzala via determining the accumulation of Cu, Zn, Cd and Pb in water, plankton and some tissues of *Liza aurata*.

## MATERIALS AND METHODS

Lake Manzala is bounded by the Mediterranean Sea to the north, the Suez canal to the east and Damietta branch of Nile to the west (Fig.1).




Figure (1): Location of sampling sites (\*) in Lake Manzala; El-Manzala (I), El-Diba (II), El-Ratama (III), El-Sirw (IV) and Bahr El-Bakar (V)

The lake has gradually changed from a brackish environment to eutrophic freshwater basin due to the increased amounts of agricultural drainage water and sewage discharge into it via seven major drains (Abdel-Baky *et al.*, 1998). Water, plankton and fish (*Liza aurata*, mugilidae) samples were collected from five different locations in the Lake (Fig.1) during four seasons from winter 2001 to autumn 2002. The locations were chosen to represent different levels of pollution. Water samples were collected monthly from 50 cm depth in two liters

polyethylene bottle acidified with nitric acid and kept for analysis. Plankton (zoo and phytoplankton) samples were collected with a plankton net of 55  $\mu\text{m}$  mesh size, through vertical hauls from the upper 10cm layer. Filtered plankton samples were acidified with HCl and kept for analysis. Parts of gills, skin and dorsal muscle were taken from each fish, weighed, put in small Erlenmeyer flasks, dried in an oven at 105 °C for about 24 hours and digested by conc. Nitric acid and perchloric acid on a hotplate until the solution became clear. Cu, Zn, Cd and Pb concentrations in water were determined by extraction method (APHA,1998) using atomic absorption spectrophotometer. Plankton and fish samples were prepared for heavy metals analysis according to the method described by Kalay *et al.*(1999). Two-way ANOVA was employed to find the significant differences of heavy metals concentration in water, plankton and fish organs with regard to sites and seasons (Bailey, 1982).

## RESULTS AND DISCUSSION

The mean concentrations of Cu, Zn, Cd, Pb in water samples collected from Lake Manzala are shown in Table 1. The mean concentrations of the tested metals in water were found in the following order: Cd (0.020)< Pb (0.022)< Cu(0.055)<Zn (0.311) mg/l. This order of occurrence agrees with the previous studies performed on Lake Manzala (Abdel-Baky *et al.*, 1998; Ibrahim *et al.*, 1999). All the metals attained their maximum values at site V which receives huge quantities of sewage and industrial wastes, beside agricultural drainage water via Bahr Al-Bakar drain. Badaway and Wahaab (1997) reported that water in Bahr Al-Bakar region is not suitable for human use. It was found that this site is rich in organic carbon (Dheina, 2007) and some authors found a correlation between the concentration of heavy metals in water and the abundance of organic matter (Radwan *et al.*1990a; Abdel-Baky *et al.*, 1998). Site II appeared to be the least polluted region of the lake as it contained the lowest levels of the investigated metals. Since it did not receive much of agricultural, industrial and sewage drains. The levels of metals exhibited seasonal fluctuations, where Cu, Cd, Pb showed significant differences between seasons. Their highest levels were found during summer, while their lowest values occurred during winter. These seasonal variations may be due to the fluctuation of the amount of agricultural drainage water, sewage effluents and industrial wastes discharged into the lake (Zyada, 1995). Ali and Abdel-Satar (2005) attributed the increase of metals concentration in water during hot seasons (spring and summer) to the release of heavy metals from sediment to the overlying water under the effect of both high temperature and fermentation process due to decomposition of organic matter. The seasonal variations of metals in water were reported by different authors in different water bodies: El-Safy and Al-Ghannam (1996), Abdel-Baky *et al.*(1998), Ibrahim *et al.* (1999) in Lake Manzala, Hamed(1998) in River Nile. Compared with the previous studies of Lake Manzala, Abdel-Hamid and El-Zareef (1996) found lower values of Cu (0.01-0.02)mg/l, El-Safy and

Al-Ghannam (1996) obtained lower Cd but higher Pb, Abdel-Baky *et al.* (1998) recorded higher values of Cu (0.08), Zn (7.94), Cd (0.11), Pb (0.64) mg/l, Ibrahim *et al.* (1999) reported higher value of pb (0.09) but lower levels of Cu (0.03), Zn (0.23) and Cd (0.005) mg/l. Compared with other lakes, Cu, Zn, Cd, Pb in Lake Manzala are higher than those of Piaseczno Lake (Poland) (0.015, 0.058, 0.001, 0.018 mg/l respectively) (Radwan *et al.*, 1990a), Lapland Lake (Finland) had higher Zn (1.84 mg/l) (Mannio *et al.*, 1995), Dominic Lake had higher Cu (3.93mg/l) (Szymanowska *et al.*, 1999). Higher concentrations of Cd (0.11) and Pb (0.086) mg/l were found in Beysehir Lake, Turkey (Altindag and Yigit, 2005). Uluabat Lake (Turkey) contained higher Cu (0.14), Cd (0.04), Pb (0.03) mg/l (Elmaci *et al.*, 2007).

According to USEPA (1986) Cu, Zn, Pb levels in Lake Manzala were within the permissible limit recommended for drinking and irrigation purposes, while that of Cd was found higher than those recommended.

Studying of heavy metals concentrations in plankton is very important because plankton is often the main diet for many predators and may remarkably contribute to the transfer of heavy metals to higher trophic levels. The results (Table 2) indicate that Cu, Zn, Cd, Pb concentrations in plankton were much higher than those of water. This may be related to the large surface of plankton organisms (phyto + zooplankton) in relation to their mass unit, and their active metabolism leading to rapid adsorption of various pollutants (Ravera, 2001). The latter author added that some algal species protect themselves by trapping and accumulating pollutants (e.g. metals) in their polysaccharides wall. The order of abundance of metals in plankton was Zn > Cu > Pb > Cd. This corresponds to the same order of abundance of these metals in water, which supports the hypothesis that water is an important source of plankton contamination. Elmaci *et al.* (2007) reported that the quantity of heavy metals in plankton depends on their concentration in water and partially on sediment. All the metals in plankton attained their maximum values at site V, where its water had the highest concentrations of these metals. The accumulation of heavy metals in plankton has been reported to depend upon several factors, such as the productivity of water body, the physico-chemical properties of the water, quantitative and qualitative species composition of zoo and phytoplankton, capacity of heavy metals absorbance and season (Radwan *et al.*, 1990b; Kerrison *et al.*, 1998; Elmaci *et al.*, 2007). There were no significant differences in metals in plankton between sites and seasons. Compared with other studies, small plankton and macro zooplankton from American lakes accumulated lower levels of Cu, Zn, Cd, Pb (Chen *et al.*, 2000). Plankton from lakes in southern Finland showed higher level of Cu but lower levels of Cd, Zn, Pb (Tulonen *et al.*, 2006). Elmaci *et al.* (2007) recorded enormously higher concentrations of Cu (6820.0), Zn (20290.0), Cd (1450.0), Pb (580.0) µg/g dry weight.

Heavy metals concentration in muscle, skin and gills of *Liza aurata* are shown in Tables (3-6). There were significant differences between sites, seasons

and fish organs. The highest concentrations of Cu, Zn, Cd, Pb were found in tissues of fish from site V, where its water contained the highest levels of the measured metals. This agrees with Shakweer (1998) who concluded that the level of bioaccumulation of trace metals in various organs of fish reflects the degree of water pollution in aquatic environment in which such fish are living. Ravera (2001) reported that if an environment receives foreign pollutants the organism living in it will take up the pollutants from the water or/and food, and concentrate it in its body. The order of detection of metals in the fish organs was as follow: gills>skin>muscles. Gills accumulated the highest level of Zn (62.018-99.80), Cu (11.88-15.48), Pb (6.9-10.26) Cd (2.93-5.19),  $\mu\text{g/g}$  dry weight. The high content of metals in gill tissues can be attributed to the fact that fish gills play a distinct role in metal uptake from the environment. Due to its respiratory function, gills are in direct contact with the contaminated water and have the thinnest epithelium of all organs (Kotze *et al.*, 1999). This result agrees with many authors who reported that gills have a high tendency to accumulate heavy metals (Unlu *et al.*, 1996; Kotze *et al.*, 1999; Wong *et al.*, 2001; Coetzee *et al.*, 2002; Altindag and Yigit, 2005). Compared with other studies, gills in the present study showed higher concentrations of Cu, Zn, Cd, Pb than those reported in *Mugil cephalus* from northeast Mediterranean Sea (Canli and Atli, 2003). Following the gills, the skin accumulated lesser concentrations of the metals. The skin tissue together with the gill tissues are characterized by a mucus layer on the outer surface. This can indicate them as excretion routes involving the slaughting off mucus from their surface (Varanci and Markey, 1978, Yilmaz 2003). Skin of *Mugil cephalus* from Iskenderun Bay (Turkey) accumulated higher levels of Pb and Zn (Yilmaz, 2005). Muscles retained the lowest concentrations of the measured metals. This finding confirms the observations of many authors who showed that fish muscles have low tendency to accumulate heavy metals to which they are exposed (Blasco *et al.*, 1998, Canli *et al.*, 1998, Ibrahim *et al.*, 1999, Canli and Atli, 2003, Karaded *et al.*, 2004, Yilmaz, 2005). In light of the recommended permissible limits of heavy metals in fish tissue for human consumption according to the National Health Medical Research Council (NHMRC) (cited from Ibrahim *et al.*, 1999b), it can be declared that the muscles of *Liza aurata*, in the present study, are considered safe for human consumption. Metals concentrations in fish organs exhibited seasonal variations in which all the detected metals attained their highest levels during summer, while their lowest values were found during winter. These seasonal variations can be attributed to the increase or decrease of drainage water discharged into the lake (Abdel-Baky *et al.*, 1998). Compared with other studies, *Liza aurata* from the middle eastern Coast of Tunisia accumulated in their muscle higher levels of Cu (4.78), Zn (45.0) but lower level of Cd (0.07)  $\mu\text{g/g}$  dry weight (Hamza-Chaffai *et al.*, 1996). Enormously higher concentrations of Cu (23.16) and Zn (27.26)  $\mu\text{g/g}$  wet weight were found in muscle of *Liza abu* from Tigris River (Turkey) (Unlu *et al.*, 1996). Blasco *et al.* (1998) measured a

remarkably high concentration of Cu and Zn in muscle of *Liza aurata* from Cadiz Bay (Spain). Higher levels of Cu, Zn, Cd, Pb were detected also in muscles of *Liza ramada* from Lake Manzala and from Damietta Nile Estuary (Ibrahim *et al.*, 1999a&b). Higher concentrations of Zn (37.39), Pb (5.32), Cu (4.41) but lower Cd (0.66)  $\mu\text{g/g}$  dry weight were recorded in muscle of *Mugil cephalus* from the north east Mediterranean Sea (Canli and Atli, 2003).

### CONCLUSION

Results of the present study clearly demonstrate that Lake Manzala is highly contaminated with Cu, Zn, Cd and Pb due to the continuous discharge of different pollutants into it. Great efforts and co-operation between different authorities are needed to protect the lake from pollution and reduce environmental risk. This can be achieved by treatment of the agricultural, industrial and sewage discharges. Regular evaluation of pollutants in the lake is also very important.

### REFERENCES

- Abdel-Baky, T. E.; Hagra, A. E.; Hassan, S. H. and Zyadah, M. A (1998). Environmental impact assessment of pollution in Lake Manzalah, 1-Distribution of some heavy metals in water and sediment. J. Egypt. Ger. Soc. Zool., 26 (B): 25-38.
- Abdelhamid, A. M. and El-Zareef, A. A. (1996). Further studies of the pollution status on the southern region of El-Manzalah Lake. Proc. Conf. Food Borne contamination and Egyptian's Health, Mansoura, Nov., 26-27, pp: 141-150.
- Ali, M.H. and Abdel-Satar, A.M. (2005). Studies of some heavy metals in water, sediment, fish and fish diets in some fish farms in El-Fayoum province, Egypt. J. Aquat. Res., 31(2):261-273.
- Altindag, A. and Yigit, S. (2005). Assessment of heavy metal concentrations in the food web of lake Beysehir, Turkey. Chemosphere, 60: 552-556.
- APHA (1998): Standard methods for the examination of water and wastewater. 20<sup>th</sup> Edition. American Public Health Association, Bultimore, USA.
- Badawy, M. I.; Wahaab, R. A. and Abou Waly, H. F. (1995). Petroleum and chlorinated hydrocarbons in water from Lake Manzala and associated canals. Bull. Environ. Contam. Toxicol. 55: 258-263.

- Badawy, M. I. and Wahaab, R. A. (1997). Environmental impact of some chemical pollutants on Lake Manzala. *International Journal of Environmental Health Research*, 7: 161-170.
- Bailey, N. (1982). *Statistical methods in biology*. 2<sup>nd</sup> Ed. (Biological Science Texts), pp: 216.
- Blasco, J.; Rubio, J. A.; Forja, J.; Gomezparra, A. and Establier, R. (1998). Heavy metals in some fishes of the Mugilidae family from salt ponds of Cadiz Bay, Spain. *Ecotoxicology and Environmental Restoration*, 1 (2): 71-77.
- Canli, M.; Ay, Ö. and Kalay, M. (1998). Levels of heavy metals (Cd, Pb, Cu, Cr and Ni) in tissues of *Cyprinus carpio*, *Barbus capito* and *Chondrostoma regium* from the Seyhan River, Turkey. *Turkish, J. Zoology*, 22: 149-157.
- Canli, M. and Atli, G. (2003). The relationships between heavy metal (Cd, Cr, Cu, Fe, Pb, and Zn) levels and the size of six Mediterranean fish species. *Environmental Pollution*, 121: 129-136.
- Chen, C.Y.; Stemberger, R. S.; Klaue, B.; Blum, J. D.; Pickhardt, P. C. and Folt, C. L. (2000). Accumulation of heavy metals in food web components across a gradient of lakes. *Limnol. Oceanogr.*, 45(7): 1525-1536.
- Coetzee, L.; du Preez, H. H. and van Vuren, J. H. J. (2002). Metal concentrations in *Clarias gariepinus* and *Labeo umbratus* from the Olifant and Klein Olifants River, Mpumalanga, South Africa: Zinc, copper, manganese, lead, chromium, nickel, aluminium and iron. *Water SA*, 28 (4): 433-448.
- Dheina, N.A. (2007). Biological and ecological studies on some species of family Mugilidae from different habitats in relation to their parasitic fauna. Ph. D. Thesis. Damietta Faculty of Science, Mansoura University, Egypt.
- Elmaci, A.; Teksoy, A.; OlcayTopac, F.; Ozengin, N.; Kurtoglu, S. and Savas Baskaya, H. (2007). Assessment of heavy metals in lake Uluabat, Turkey. *Afr. J. Biotech.*, 6 (19): 2236-2244.
- El-Safy, M. K. and Al-Ghannam, M. L. (1996). Studies on some heavy metal pollutants in fish of El-Lake Manzala. *Proc. Conf. Food Borne Contamination and Egyptian's Health*, Mansoura, Nov. 26-27, pp.151-180.

- Fernandez, M.; Cuesta, S.; Jimene, O.; Garcia, M. A. Hernandez, L.M.; Marina, M. L. and Gonzalez, M. J.(2000). Organochlorine and heavy metal residues in the water, sediment system of the southeast regional park in Madrid, Spain. *Chemosphere*, 41:801-812.
- Hamed, M. A. (1998). Distribution of trace metals in the River Nile ecosystem Damietta branch between Mansoura city and Damietta province. *J. Egypt. Ger. Soc. Zool.*, 27 (A): 399-415.
- Hamza-Chaffai, A.; Romeo, M. and El-Abed, A. (1996). Heavy metals in different fishes from the Middle Eastern coast of Tunisia. *Bull. Environ. Contam. Toxicol.*, 56: 766-773.
- Ibrahim, A. M.; Bahnasawy, M. H.; Mansy, S. E. and El-Fayomy, R. I. (1999a). Heavy metal accumulation in water, Sediment and some fishes in Lake Manzala, Egypt. *J. Egypt. Ger. Soc. Zool.*, 29 (B): 43-58.
- Ibrahim, A. M.; Bahnasawy, M. H.; Mansy S. E. and EL-Fayomy, R. I. (1999b). Distribution of heavy metals in the Damietta Nile Estuary ecosystem. *Egypt. J. Aquat. Biol & Fish.*, 3(4): 369-397.
- Ibrahim, A. M.; Bahnasawy, M. H.; Mansy, S. E. and EL-Fayomy, R. I. (2000). On some heavy metal levels in water, sediment and marine organisms from the Mediterranean coast of Lake Manzala. *Egypt. J. Aquat. Biol & Fish.*, 4(4): 61-81.
- Janssen, C.R., DeSchamphelaere, K., Heijerick, D., Muysen, B., Lock, K., Bossuyt, B., Vangheluwe, M. and VanSprang, P.(2000). Uncertainties in the environmental risk assessment of metals. *Hum.Ecol.Risk.Assess.*,6: 1003-1018.
- Kalay, M.; Ay, Ö. and canli, M. (1999). Heavy metal concentrations in fish tissues from the Northeast Mediterranean Sea. *Bull. Environ. Contam. Toxicol.*, 63: 673-681.
- Karadede, H.; Oymak, S. A. and Ünlü, E. (2004). Heavy metals in mullet, *Liza abu*, and Catfish, *Silurus triostegus*, from the Atatürk Dam Lake (Euphrates), Turkey. *Environment International*, 30: 183-188.
- Kerrison, P.H., Annoni, D., Zerini, S., Ravera, O. and Moss. B. (1998). Effects of low concentrations of heavy metals on plankton community dynamics in a small, shallow, fertile lake. *J. Plankton Res.*,10:779-812.

- Kotze, P.; Du preez, H. H. and Van Vuren, J. H. (1999). Bioaccumulation of copper and zinc in *Oreochromis mossambicus* and *Clarias gariepinus*, from the Olifants River, Mpumalanga, South Africa. *water S. A.*, 25 (1):99-110.
- Mannio, J., Jarvien, O., Tuominen, R. and Verta, M. (1995). Survey of trace elements in lake waters of Finnish Lapland using the ICP-MS technique. *Sci. Total Environ.*, 160-161: 433-439.
- Mason, C. (1991). *Biology of freshwater pollution*. Harlow: Longman Scientific and Technical.
- Radwan, S.; Kowalik, W. and Kornijow, R. (1990a). Accumulation of heavy metals in a Lake ecosystem. *Sci. Total Environ.*, 96: 121-129.
- Radwan, S.; Kowalik, W. and Kowalczyk, C. (1990b). Occurrence of heavy metals in water, phytoplankton and zooplankton of mesotrophic Lake in Eastern Poland. *Sci. Total Environ.*, 96: 115-120.
- Ravera, O. (2001). Monitoring of the aquatic environment by species accumulator of pollutants: a review. *J. Inmol.*, 60: 63-78.
- Shakweer, L. M. (1998). Concentration levels of some trace metals in *Oreochromis niloticus* at highly and less polluted areas of Mariut Lake. *J. Egypt. Ger. Soc. Zool.*, 25(A): 125-144.
- Stephen, C.; Jewett, A. and Sathy Naidu, A. (2000). Assessment of heavy metals in red king crabs following offshore placer gold mining. *Mar. Pollut. Bull.*, 40(6): 478-490.
- Szymanowska, A.; Samecka-Symerman, A. and Kampers, A. J. (1999). Heavy metals in three lakes in west Poland. *Ecotoxicol. Environ. Safety*, 43: 21-29.
- Tarvainen, T.; Lahermo, P. and Mannio, J. (1997). Source of trace metals in streams and headwater lakes in Finland. *Water, Air, Soil Pollut.*, 94: 1-32.
- Tulonen, T.; Pihlstrom, M.; Arvola, L. and Rask, M. (2006). Concentrations of heavy metals in food web components of small, boreal lakes. *Boreal Environ. Res.*, 11:185-194
- U. S. E. P. A. (United States Environmental Protection Agency) (1986). *Quality criteria for water*. 1986. EPA 440 15-86-001. May 1986.

- Ünlü, E.; Akba, O.; Sevim, S. and Gümüş, B. (1996). Heavy metal levels in mullet, *Liza abu* (Heckel, 1843) (Mugilidae) from the Tigris River, Turkey. *Fresenius Envir. Bull.*, 5: 107-112.
- Varanasi, U. and Markey, Y. D. (1978). Uptake and release of lead and cadmium in skin and mucus of coho salmon (*Oncorhynchus kisutch*). *Comp. Biochem. Physiol.* 60C: 187-191.
- Wong, C.K.; Wong, P. P. K. and Chu, L. M. (2001). Heavy metal concentrations in marine fishes collected from fish culture sites in Hong Kong. *Arch. Environ. Contam. Toxicol.* 40:60-69.
- Yilmaz, A.B. (2003). Levels of heavy metals (Fe, Cu, Ni, Cr, Pb and Zn) in tissue of *Mugil cephalus* and *Trachurus mediterraneus* from Iskenderun Bay, Turkey. *Environ. Res.*, 92: 277-281.
- Yilmaz, A.B.(2005). Comparison of heavy metal levels of grey mullet (*Mugil cephalus*) and sea bream (*Sparus aurata*) caught in Iskenderun Bay (Turkey). *Turk. J. Vet. Anim. Sci.*, 29:257-262.

Table 1: Seasonal variations of heavy metals concentrations (mg/l) in water of Lake Manzala.

| Elements | Site  | Seasons             |                     |                     |                     |                     | ANOVA         |    |         |       |
|----------|-------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------|----|---------|-------|
|          |       | Winter              | Spring              | Summer              | Autumn              | Total               | Factor        | df | F value | Sig.  |
| Cu       | I     | 0.038<br>±<br>0.002 | 0.051<br>±<br>0.042 | 0.061<br>±<br>0.02  | 0.040<br>±<br>0     | 0.048<br>±<br>0.031 |               |    |         |       |
|          | II    | 0.009<br>±<br>0     | 0.028<br>±<br>0.004 | 0.031<br>±<br>0.013 | 0.016<br>±<br>0.029 | 0.021<br>±<br>0.009 | Site          | 4  | 16.096  | 0     |
|          | III   | 0.025<br>±<br>0.003 | 0.040<br>±<br>0.004 | 0.055<br>±<br>0.009 | 0.038<br>±<br>0.007 | 0.040<br>±<br>0.004 | Season        | 3  | 14.288  | 0     |
|          | IV    | 0.032<br>±<br>0.002 | 0.065<br>±<br>0.002 | 0.083<br>±<br>0.004 | 0.049<br>±<br>0.004 | 0.057<br>±<br>0.004 | Site x Season | 12 | 13.479  | 0     |
|          | V     | 0.053<br>±<br>0.009 | 0.111<br>±<br>0.002 | 0.192<br>±<br>0.002 | 0.088<br>±<br>0.02  | 0.111<br>±<br>0.007 |               |    |         |       |
|          | Total | 0.031<br>±<br>0.003 | 0.059<br>±<br>0.011 | 0.084<br>±<br>0.010 | 0.046<br>±<br>0.012 | 0.055<br>±<br>0.011 |               |    |         |       |
| Zn       | I     | 0.177<br>±<br>0.221 | 0.370<br>±<br>0.136 | 0.472<br>±<br>0.179 | 0.246<br>±<br>0.112 | 0.316<br>±<br>0.154 |               |    |         |       |
|          | II    | 0.139<br>±<br>0     | 0.281<br>±<br>0.058 | 0.301<br>±<br>0.065 | 0.198<br>±<br>0.009 | 0.230<br>±<br>0.031 | Site          | 4  | 2.421   | 0.064 |
|          | III   | 0.181<br>±<br>0.013 | 0.310<br>±<br>0.161 | 0.372<br>±<br>0.013 | 0.226<br>±<br>0.114 | 0.272<br>±<br>0.058 | Season        | 3  | 3.156   | 0.035 |
|          | IV    | 0.198<br>±<br>0.047 | 0.382<br>±<br>0.036 | 0.493<br>±<br>0.147 | 0.288<br>±<br>0.042 | 0.340<br>±<br>0.257 | Site x Season | 12 | 2.726   | 0.009 |
|          | V     | 0.232<br>±<br>0.226 | 0.470<br>±<br>0.087 | 0.529<br>±<br>0.183 | 0.352<br>±<br>0.031 | 0.396<br>±<br>0.143 |               |    |         |       |
|          | Total | 0.185<br>±<br>0.101 | 0.363<br>±<br>0.096 | 0.433<br>±<br>0.117 | 0.262<br>±<br>0.062 | 0.311<br>±<br>0.129 |               |    |         |       |
| Cd       | I     | 0.018<br>±<br>0.002 | 0.021<br>±<br>0.018 | 0.025<br>±<br>0.002 | 0.014<br>±<br>0.025 | 0.020<br>±<br>0.004 |               |    |         |       |
|          | II    | N.D.                | 0.015<br>±<br>0.031 | 0.019<br>±<br>0.007 | 0.011<br>±<br>0.02  | 0.011<br>±<br>0.007 | Site          | 4  | 12.854  | 0     |
|          | III   | 0.009<br>±<br>0.011 | 0.018<br>±<br>0.002 | 0.022<br>±<br>0.007 | 0.014<br>±<br>0.011 | 0.016<br>±<br>0.011 | Season        | 3  | 4.607   | 0.007 |
|          | IV    | 0.016<br>±<br>0.007 | 0.026<br>±<br>0.002 | 0.031<br>±<br>0.007 | 0.021<br>±<br>0.009 | 0.024<br>±<br>0.004 | Site x Season | 12 | 5.614   | 0     |
|          | V     | 0.021<br>±<br>0.011 | 0.031<br>±<br>0.009 | 0.038<br>±<br>0.002 | 0.027<br>±<br>0.009 | 0.029<br>±<br>0.009 |               |    |         |       |
|          | Total | 0.016<br>±<br>0.008 | 0.022<br>±<br>0.012 | 0.027<br>±<br>0.005 | 0.017<br>±<br>0.015 | 0.020<br>±<br>0.007 |               |    |         |       |
| Pb       | I     | 0.006<br>±<br>0.002 | 0.026<br>±<br>0.011 | 0.034<br>±<br>0.002 | 0.011<br>±<br>0.002 | 0.019<br>±<br>0.007 |               |    |         |       |
|          | II    | N.D.                | 0.011<br>±<br>0.007 | 0.017<br>±<br>0.002 | 0.006<br>±<br>0.002 | 0.009<br>±<br>0.002 | Site          | 4  | 11.707  | 0     |
|          | III   | N.D.                | 0.017<br>±<br>0.016 | 0.029<br>±<br>0.007 | 0.008<br>±<br>0.002 | 0.014<br>±<br>0.007 | Season        | 3  | 4.601   | 0.007 |
|          | IV    | 0.008<br>±<br>0.004 | 0.032<br>±<br>0.007 | 0.054<br>±<br>0.011 | 0.020<br>±<br>0.007 | 0.029<br>±<br>0.007 | Site x Season | 12 | 10.943  | 0     |
|          | V     | 0.012<br>±<br>0.007 | 0.046<br>±<br>0.002 | 0.074<br>±<br>0.018 | 0.029<br>±<br>0.002 | 0.040<br>±<br>0.004 |               |    |         |       |
|          | Total | 0.009<br>±<br>0.004 | 0.026<br>±<br>0.009 | 0.042<br>±<br>0.008 | 0.015<br>±<br>0.003 | 0.022<br>±<br>0.005 |               |    |         |       |

N.D. : not detected

Table 2: Seasonal variations of heavy metals concentrations ( $\mu\text{g/g}$  dry weight) in plankton from Lake Manzala.

| Elements | Site    | Seasons |         |         |         |               | ANOVA  |       |         |      |
|----------|---------|---------|---------|---------|---------|---------------|--------|-------|---------|------|
|          |         | Winter  | Spring  | Summer  | Autumn  | Total         | Factor | df    | F value | Sig. |
| Cu       | I       | 88.430  | 111.430 | 118.860 | 96.340  | 103.760       | Site   | 4     | 2.315   | 0.74 |
|          |         | ±       | ±       | ±       | ±       | ±             |        |       |         |      |
|          | 13.830  | 25.160  | 23.050  | 15.370  | 18.180  |               |        |       |         |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
|          | 48.570  | 69.440  | 75.480  | 59.910  | 63.350  |               |        |       |         |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
|          | 21.230  | 8.780   | 31.840  | 20.670  | 25.360  |               |        |       |         |      |
| II       | 71.890  | 84.860  | 93.830  | 78.000  | 82.140  | Season        | 3      | 1.142 | 0.344   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 8.100    | 23.050  | 23.500  | 15.370  | 20.050  |         |               |        |       |         |      |
| III      | 90.570  | 115.750 | 135.720 | 104.000 | 111.510 | Site x Season | 12     | 2.166 | 0.034   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 23.060   | 20.490  | 30.460  | 30.740  | 17.200  |         |               |        |       |         |      |
| IV       | 108.760 | 136.620 | 154.430 | 126.000 | 131.450 | Site          | 4      | 1.557 | 0.204   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 34.890   | 39.110  | 41.510  | 32.120  | 49.640  |         |               |        |       |         |      |
| V        | 81.644  | 103.620 | 115.664 | 92.850  | 98.442  | Season        | 3      | 2.386 | 0.083   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 20.222   | 23.318  | 30.072  | 22.854  | 26.086  |         |               |        |       |         |      |
| Total    | 406.420 | 537.720 | 549.090 | 462.180 | 488.850 | Site x Season | 12     | 2.048 | 0.045   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 140.220  | 71.710  | 97.020  | 89.950  | 168.740 |         |               |        |       |         |      |
| I        | 248.540 | 358.040 | 380.150 | 267.190 | 313.480 | Site          | 4      | 1.557 | 0.204   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 126.800  | 64.740  | 92.860  | 185.200 | 115.400 |         |               |        |       |         |      |
| II       | 251.640 | 428.960 | 460.650 | 390.410 | 382.910 | Season        | 3      | 2.386 | 0.083   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 104.860  | 177.860 | 125.260 | 52.360  | 124.150 |         |               |        |       |         |      |
| III      | 430.670 | 610.280 | 695.730 | 520.140 | 564.210 | Site x Season | 12     | 2.048 | 0.045   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 74.050   | 174.540 | 216.230 | 110.890 | 77.180  |         |               |        |       |         |      |
| IV       | 490.850 | 620.280 | 737.650 | 560.460 | 602.310 | Site          | 4      | 1.557 | 0.204   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 66.650   | 144.180 | 91.970  | 48.940  | 91.080  |         |               |        |       |         |      |
| V        | 365.624 | 511.056 | 564.654 | 440.076 | 470.352 | Season        | 3      | 2.386 | 0.083   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 102.516  | 126.606 | 124.668 | 97.468  | 115.310 |         |               |        |       |         |      |
| Total    | 20.170  | 27.000  | 32.000  | 25.670  | 26.210  | Site x Season | 12     | 2.947 | 0.005   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 1.620    | 6.450   | 8.940   | 8.600   | 9.400   |         |               |        |       |         |      |
| I        | 14.290  | 21.280  | 25.340  | 17.760  | 19.670  | Site          | 4      | 2.065 | 0.104   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 7.750    | 2.450   | 7.750   | 3.270   | 9.970   |         |               |        |       |         |      |
| II       | 18.610  | 26.000  | 30.430  | 22.670  | 24.430  | Season        | 3      | 2.350 | 0.087   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 5.090    | 6.450   | 6.580   | 9.660   | 9.540   |         |               |        |       |         |      |
| III      | 24.820  | 33.220  | 36.670  | 29.470  | 31.040  | Site x Season | 12     | 2.947 | 0.005   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 12.910   | 2.860   | 8.600   | 8.940   | 11.220  |         |               |        |       |         |      |
| IV       | 27.380  | 39.970  | 46.000  | 31.270  | 36.150  | Site          | 4      | 2.065 | 0.104   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 7.920    | 6.000   | 12.910  | 8.600   | 9.800   |         |               |        |       |         |      |
| V        | 21.054  | 29.494  | 34.088  | 25.368  | 27.500  | Season        | 3      | 2.350 | 0.087   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 7.058    | 4.842   | 8.956   | 7.814   | 9.986   |         |               |        |       |         |      |
| Total    | 56.750  | 79.130  | 91.130  | 69.530  | 74.130  | Site x Season | 12     | 0.628 | 0.806   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 13.450   | 33.620  | 33.620  | 22.410  | 16.630  |         |               |        |       |         |      |
| I        | 44.750  | 61.210  | 72.050  | 55.310  | 58.330  | Site          | 4      | 3.644 | 0.013   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 13.450   | 28.870  | 15.930  | 18.280  | 47.010  |         |               |        |       |         |      |
| II       | 51.300  | 74.780  | 83.720  | 67.450  | 69.310  | Season        | 3      | 0.961 | 0.421   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 2.040    | 27.500  | 22.460  | 24.820  | 24.510  |         |               |        |       |         |      |
| III      | 77.130  | 109.300 | 118.450 | 91.170  | 99.010  | Site x Season | 12     | 0.628 | 0.806   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 33.620   | 34.350  | 51.640  | 44.830  | 27.270  |         |               |        |       |         |      |
| IV       | 95.690  | 132.830 | 149.130 | 118.530 | 124.040 | Site          | 4      | 3.644 | 0.013   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 25.180   | 43.050  | 33.620  | 53.930  | 31.460  |         |               |        |       |         |      |
| V        | 65.124  | 91.450  | 102.896 | 80.398  | 84.964  | Season        | 3      | 0.961 | 0.421   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |
| 17.548   | 33.478  | 31.454  | 32.854  | 29.376  |         |               |        |       |         |      |
| Total    | 17.548  | 33.478  | 31.454  | 32.854  | 29.376  | Site x Season | 12     | 0.628 | 0.806   |      |
|          | ±       | ±       | ±       | ±       | ±       |               |        |       |         |      |

Table.3: Seasonal variations of copper concentration ((µg/g dry weight) in different organs of Liza aurata from Lake Manzala.

| Organ   | Site | Seasons              |                      |                      |                      |                      | ANOVA             |    |          |       |
|---------|------|----------------------|----------------------|----------------------|----------------------|----------------------|-------------------|----|----------|-------|
|         |      | Winter               | Spring               | Summer               | Autumn               | Total                | Factor            | df | F value  | Sig.  |
| Gills   | I    | 11.520<br>±<br>0.610 | 12.340<br>±<br>0.770 | 13.850<br>±<br>1.250 | 13.620<br>±<br>0.280 | 12.830<br>±<br>1.230 | Site              | 4  | 80.467   | 0     |
|         | II   | 10.510<br>±<br>1.010 | 11.660<br>±<br>1.180 | 12.770<br>±<br>0.800 | 12.580<br>±<br>0.540 | 11.880<br>±<br>1.240 | Season            | 3  | 48.159   | 0     |
|         | III  | 11.820<br>±<br>0.580 | 12.820<br>±<br>1.030 | 13.250<br>±<br>0.700 | 12.660<br>±<br>0.840 | 12.640<br>±<br>0.910 | Organ             | 2  | 3565.515 | 0     |
|         | IV   | 11.830<br>±<br>0.810 | 14.370<br>±<br>1.810 | 14.420<br>±<br>2.890 | 13.220<br>±<br>0.670 | 13.460<br>±<br>1.970 | Site x<br>Season  | 12 | 1.325    | 0.204 |
|         | V    | 14.220<br>±<br>0.830 | 16.510<br>±<br>1.290 | 16.520<br>±<br>2.320 | 14.680<br>±<br>1.440 | 15.480<br>±<br>1.790 | Site x<br>Organ   | 8  | 7.986    | 0     |
| Skin    | I    | 7.780<br>±<br>0.080  | 7.870<br>±<br>0.150  | 8.940<br>±<br>0.310  | 8.870<br>±<br>0.190  | 8.370<br>±<br>0.590  | Season x<br>Organ | 6  | 4.024    | 0.001 |
|         | II   | 6.730<br>±<br>0.080  | 6.820<br>±<br>0.200  | 7.930<br>±<br>0.420  | 7.340<br>±<br>0.120  | 7.200<br>±<br>0.540  |                   |    |          |       |
|         | III  | 6.740<br>±<br>0.570  | 7.860<br>±<br>0.230  | 8.920<br>±<br>0.530  | 7.840<br>±<br>0.540  | 7.840<br>±<br>0.910  | Site              |    |          |       |
|         | IV   | 7.830<br>±<br>0.150  | 8.920<br>±<br>0.200  | 8.960<br>±<br>0.290  | 8.860<br>±<br>0.070  | 8.640<br>±<br>0.510  | x                 |    |          |       |
|         | V    | 8.960<br>±<br>0.210  | 9.210<br>±<br>0.280  | 9.710<br>±<br>0.490  | 9.590<br>±<br>0.450  | 9.370<br>±<br>0.460  | Season            |    |          |       |
| Muscles | I    | 3.430<br>±<br>0.250  | 3.960<br>±<br>0.240  | 4.700<br>±<br>0.230  | 3.880<br>±<br>0.220  | 3.990<br>±<br>0.510  | x                 |    |          |       |
|         | II   | 3.610<br>±<br>0.480  | 3.810<br>±<br>0.110  | 4.030<br>±<br>0.260  | 3.550<br>±<br>0.140  | 3.750<br>±<br>0.330  | Organ             | 24 | 1.555    | 0.052 |
|         | III  | 3.460<br>±<br>0.210  | 3.990<br>±<br>0.340  | 4.150<br>±<br>0.190  | 4.110<br>±<br>0.210  | 3.930<br>±<br>0.360  |                   |    |          |       |
|         | IV   | 3.990<br>±<br>0.300  | 4.570<br>±<br>0.120  | 4.740<br>±<br>0.410  | 4.520<br>±<br>0.180  | 4.460<br>±<br>0.380  |                   |    |          |       |
|         | V    | 4.000<br>±<br>0.270  | 4.930<br>±<br>0.320  | 5.490<br>±<br>0.150  | 5.430<br>±<br>0.280  | 4.960<br>±<br>0.660  |                   |    |          |       |

Table 4: Seasonal variations of Zinc concentration ( $\mu\text{g/g}$  dry weight) in different organs of *Liza aurata* from Lake Manzala.

| Organ   | Site | Seasons |         |         |        |        | ANOVA          |    |          |      |
|---------|------|---------|---------|---------|--------|--------|----------------|----|----------|------|
|         |      | Winter  | Spring  | Summer  | Autumn | Total  | Factor         | df | F value  | Sig. |
| Gills   | I    | 52.720  | 75.360  | 86.250  | 68.460 | 70.700 | Site           | 4  | 748.475  | 0    |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | II   | 47.410  | 66.740  | 74.660  | 59.260 | 62.020 | Season         | 3  | 1091.732 | 0    |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | III  | 52.640  | 74.450  | 81.730  | 67.290 | 69.030 | Organ          | 2  | 9358.979 | 0    |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | IV   | 64.350  | 91.580  | 103.470 | 79.630 | 84.760 | Site x Season  | 12 | 21.840   | 0    |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | V    | 67.210  | 101.360 | 136.170 | 94.460 | 99.800 | Site x Organ   | 8  | 53.910   | 0    |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
| Skin    | I    | 35.310  | 54.350  | 60.280  | 46.980 | 49.230 | Season x Organ | 6  | 132.850  | 0    |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | II   | 30.630  | 42.700  | 51.340  | 39.370 | 41.010 |                |    |          |      |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | III  | 33.720  | 54.660  | 65.630  | 41.460 | 48.870 | Site           |    |          |      |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | IV   | 41.420  | 62.560  | 78.240  | 52.260 | 58.620 | x              |    |          |      |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | V    | 47.520  | 72.460  | 85.380  | 61.360 | 66.680 | Season         |    |          |      |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
| Muscles | I    | 15.500  | 19.660  | 24.060  | 18.340 | 19.390 | x              |    |          |      |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | II   | 12.460  | 18.070  | 17.460  | 14.270 | 15.570 | Organ          | 24 | 9.000    | 0    |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | III  | 12.980  | 19.680  | 21.290  | 16.760 | 17.680 |                |    |          |      |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | IV   | 19.410  | 27.760  | 29.780  | 22.440 | 24.850 |                |    |          |      |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         | V    | 25.380  | 30.240  | 35.450  | 27.640 | 29.680 |                |    |          |      |
|         |      | ±       | ±       | ±       | ±      | ±      |                |    |          |      |
|         |      | 3.190   | 1.850   | 3.340   | 2.300  | 4.600  |                |    |          |      |

Table 5: Seasonal variations of cadmium concentration ( $\mu\text{g/g}$  dry weight) in different organs of *Liza aurata* from Lake Manzala.

| Organ   | Site | Seasons             |                     |                     |                     |                     | ANOVA             |    |          |       |
|---------|------|---------------------|---------------------|---------------------|---------------------|---------------------|-------------------|----|----------|-------|
|         |      | Winter              | Spring              | Summer              | Autumn              | Total               | Factor            | df | F value  | Sig.  |
| Gills   | I    | 3.350<br>±<br>0.590 | 3.620<br>±<br>0.820 | 4.190<br>±<br>0.470 | 2.920<br>±<br>0.700 | 3.520<br>±<br>0.770 | Site              | 4  | 67.547   | 0     |
|         | II   | 2.630<br>±<br>0.520 | 3.360<br>±<br>0.550 | 3.400<br>±<br>0.310 | 2.330<br>±<br>1.170 | 2.930<br>±<br>0.810 | Season            | 3  | 34.024   | 0     |
|         | III  | 3.170<br>±<br>0.370 | 3.970<br>±<br>0.450 | 4.170<br>±<br>0.400 | 3.290<br>±<br>0.260 | 3.650<br>±<br>0.560 | Organ             | 2  | 1152.758 | 0     |
|         | IV   | 4.180<br>±<br>0.720 | 5.090<br>±<br>0.830 | 5.420<br>±<br>0.840 | 4.380<br>±<br>0.330 | 4.770<br>±<br>0.830 | Site x<br>Season  | 12 | 0.691    | 0.759 |
|         | V    | 4.570<br>±<br>0.740 | 5.280<br>±<br>0.620 | 6.060<br>±<br>1.160 | 4.850<br>±<br>1.230 | 5.190<br>±<br>1.060 | Site x<br>Organ   | 8  | 19.836   | 0     |
| Skin    | I    | 1.620<br>±<br>0.120 | 2.140<br>±<br>0.090 | 2.210<br>±<br>0.110 | 1.770<br>±<br>0.160 | 1.940<br>±<br>0.280 | Season x<br>Organ | 6  | 7.364    | 0     |
|         | II   | 1.590<br>±<br>0.070 | 1.730<br>±<br>0.090 | 1.720<br>±<br>0.090 | 1.640<br>±<br>0.200 | 1.670<br>±<br>0.130 |                   |    |          |       |
|         | III  | 1.520<br>±<br>0.070 | 2.170<br>±<br>0.090 | 1.870<br>±<br>0.150 | 1.680<br>±<br>0.140 | 1.810<br>±<br>0.270 | Site              |    |          |       |
|         | IV   | 1.670<br>±<br>0.090 | 2.280<br>±<br>0.320 | 2.340<br>±<br>0.090 | 1.970<br>±<br>0.130 | 2.070<br>±<br>0.320 | x                 |    |          |       |
|         | V    | 2.070<br>±<br>0.410 | 2.560<br>±<br>0.270 | 2.760<br>±<br>0.130 | 2.520<br>±<br>0.270 | 2.480<br>±<br>0.370 | Season            |    |          |       |
| Muscles | I    | 0.970<br>±<br>0.370 | 1.060<br>±<br>0.210 | 1.390<br>±<br>0.150 | 1.110<br>±<br>0.070 | 1.130<br>±<br>0.270 | x                 |    |          |       |
|         | II   | 0.810<br>±<br>0.120 | 0.970<br>±<br>0.060 | 1.140<br>±<br>0.080 | 1.070<br>±<br>0.100 | 1.000<br>±<br>0.150 | Organ             | 24 | 0.411    | 0.994 |
|         | III  | 0.960<br>±<br>0.120 | 1.130<br>±<br>0.060 | 1.230<br>±<br>0.070 | 1.110<br>±<br>0.070 | 1.110<br>±<br>0.130 |                   |    |          |       |
|         | IV   | 1.190<br>±<br>0.150 | 1.260<br>±<br>0.060 | 1.420<br>±<br>0.090 | 1.170<br>±<br>0.100 | 1.260<br>±<br>0.140 |                   |    |          |       |
|         | V    | 1.270<br>±<br>0.090 | 1.340<br>±<br>0.060 | 1.610<br>±<br>0.160 | 1.230<br>±<br>0.090 | 1.370<br>±<br>0.180 |                   |    |          |       |

Table 6: Seasonal variations of lead concentration ( $\mu\text{g/g}$  dry weight) in different organs of *Liza aurata* from Lake Manzala.

| Organ   | Site | Seasons             |                      |                      |                      |                      | ANOVA             |    |          |       |
|---------|------|---------------------|----------------------|----------------------|----------------------|----------------------|-------------------|----|----------|-------|
|         |      | Winter              | Spring               | Summer               | Autumn               | Total                | Factor            | df | F value  | Sig.  |
| Gills   | I    | 7.140<br>±<br>0.760 | 8.720<br>±<br>0.420  | 9.320<br>±<br>0.680  | 8.480<br>±<br>0.590  | 8.420<br>±<br>1.000  | Site              | 4  | 104.379  | 0     |
|         | II   | 6.440<br>±<br>0.670 | 7.020<br>±<br>0.490  | 7.660<br>±<br>0.560  | 6.480<br>±<br>0.570  | 6.900<br>±<br>0.730  | Season            | 3  | 40.010   | 0     |
|         | III  | 6.940<br>±<br>0.470 | 7.880<br>±<br>0.540  | 8.320<br>±<br>0.740  | 7.360<br>±<br>0.630  | 7.630<br>±<br>0.770  | Organ             | 2  | 5872.010 | 0     |
|         | IV   | 8.120<br>±<br>0.700 | 9.460<br>±<br>0.650  | 9.840<br>±<br>0.670  | 9.080<br>±<br>0.740  | 9.130<br>±<br>0.910  | Site x<br>Season  | 12 | 0.504    | 0.911 |
|         | V    | 9.460<br>±<br>0.680 | 10.280<br>±<br>0.350 | 10.940<br>±<br>0.670 | 10.380<br>±<br>0.580 | 10.270<br>±<br>0.760 | Site x<br>Organ   | 8  | 34.132   | 0     |
| Skin    | I    | 2.440<br>±<br>0.310 | 2.780<br>±<br>0.380  | 2.950<br>±<br>0.310  | 2.620<br>±<br>0.620  | 2.700<br>±<br>0.440  | Season x<br>Organ | 6  | 9.482    | 0     |
|         | II   | 2.340<br>±<br>0.240 | 2.490<br>±<br>0.530  | 2.630<br>±<br>0.370  | 2.410<br>±<br>0.380  | 2.470<br>±<br>0.370  |                   |    |          |       |
|         | III  | 2.420<br>±<br>0.340 | 2.610<br>±<br>0.360  | 2.860<br>±<br>0.320  | 2.530<br>±<br>0.350  | 2.610<br>±<br>0.350  | Site              |    |          |       |
|         | IV   | 2.620<br>±<br>0.360 | 2.860<br>±<br>0.350  | 3.070<br>±<br>0.310  | 2.720<br>±<br>0.430  | 2.820<br>±<br>0.380  | x                 |    |          |       |
|         | V    | 2.750<br>±<br>0.310 | 3.070<br>±<br>0.430  | 3.270<br>±<br>0.380  | 2.810<br>±<br>0.370  | 2.980<br>±<br>0.410  | Season            |    |          |       |
| Muscles | I    | 1.720<br>±<br>0.370 | 1.900<br>±<br>0.390  | 2.110<br>±<br>0.300  | 1.920<br>±<br>0.220  | 1.910<br>±<br>0.330  | x                 |    |          |       |
|         | II   | 1.410<br>±<br>0.200 | 1.620<br>±<br>0.520  | 1.750<br>±<br>0.180  | 1.540<br>±<br>0.190  | 1.580<br>±<br>0.310  | Organ             | 24 | 0.464    | 0.986 |
|         | III  | 1.540<br>±<br>0.300 | 1.820<br>±<br>0.400  | 1.940<br>±<br>0.350  | 1.750<br>±<br>0.280  | 1.760<br>±<br>0.340  |                   |    |          |       |
|         | IV   | 1.970<br>±<br>0.480 | 2.160<br>±<br>0.360  | 2.370<br>±<br>0.370  | 2.130<br>±<br>0.320  | 2.160<br>±<br>0.380  |                   |    |          |       |
|         | V    | 2.190<br>±<br>0.460 | 2.470<br>±<br>0.230  | 2.660<br>±<br>0.360  | 2.420<br>±<br>0.430  | 2.440<br>±<br>0.390  |                   |    |          |       |