



Effect of non-ionizing radiation on structure and medical content for plant

RH. Ebrahim^a, Hamed M El-Shora^c, Abu Bakr El-Bediwi^b, A. Abdelrazek^b



CrossMark

^aHigher Future Institute of Engineering and Technology, Telha, Egypt

^bPhysics Department, Faculty of Science, Mansoura University, 35516 Mansoura, Egypt

^cBotany Department, Faculty of Science, Mansoura University, 35516 Mansoura, Egypt

Abstract

Plants are living chemical factories for enormous array of the secondary metabolites. Due to toxic and carcinogenic effects of the synthetic antioxidants, a great deal of attention has been focused in recent decades on natural antioxidants such as polyphenols derived from various natural sources. The aim of research is to study the effect of UVA on structure and medical content of yellow mustard. The results show non-enzymatic such as glutathione and phenolic contents in yellow mustard improved after exposure to UVA. Total proline, tocopherol total flavonoids in yellow mustard varied after exposure to UVA. Also internal structure as hydroxyl group of yellow mustard changed after exposure to UVA.

Key words: phenolic, glutathione, DPPH scavenging activity, proline, yellow mustard, UVA

2. Introduction

Solar radiation is a complex mixture of ultraviolet, visible light and infrared wavelengths. UVA radiation (315– 400 nm) is a component of solar radiation. Plants are living chemical factories for the biosynthesis of a huge array of the secondary metabolites. Plants are used medicinally in different countries and are a source of many powerful drugs. The World Health Organization stated that about 80% of the world's population depends mainly on traditional medicine that mainly includes the use of plant extracts. UV light is an important abiotic elicitor, and had use in phytochemical production in a variety of plant cultures in the past [1]. Exposure to UV light stress causes stimulation of defense mechanisms in plants, thus, producing commercially important secondary compounds [2]. Some reports discussed the effect of low levels of UV radiation on plant growth [3, 4]. There are many beneficial uses of radiation that offer few risks when properly employed. The exposure to radiations can have stimulatory effects on specific morphological parameters. The mustard plant belongs to the Cruciferae (Brassicaceae) family, used in medicine is external as a liniment and relieving pain from bruises or a stiff neck and relieving colic and respiratory problems. Growth behavior, secondary metabolites and vitamins of *Nigella Sativa*

and *garden cress* changed after exposure by UVC [5, 6]. UVs have adequate energy to break the chemical bonds causing photochemical reactions and inducing changes in plant metabolic enzyme, subsequently trigger the production of secondary metabolites [7- 9]. Effect of UV is varied with duration and irradiation intensity. The objective of this work is to assess the effect of UVA radiation on structure, non-enzymatic and enzymatic antioxidants for yellow mustard.

3. Experimental methods

3.1 Structure measurements

Internal structure and molecular structure of yellow mustard are studied by Shimadzu X-ray diffractometer, (Dx-30, Japan), scanning electron microscope (JEOL JSM-6510LV, Japan) and Nicolet™ iS™ 10 FT-IR Spectrometer from USA.

3.2 GSH determination: GSH is determined using UV/V spectrophotometer, Jenway, England.

3.3 DPPH determination: Concentration ranging from 0.4g/100g to 2g/100g are prepared with methanol from each sample (100 µl) extract and DPPH radical (100 µl, 2Mm) dissolved in methanol. The mixture is stirred and left to stand for 15 min in dark. Then the absorbance is measured at 517 nm against a blank.

Percentage scavenging effect is calculated as $[(A_0 -$

*Corresponding author e-mail: baker_elbediwi@yahoo.com; (Abu Bakr El-Bediwi).

EJCHEM use only: Received date here; revised date here; accepted date here

DOI: 10.21608/EJCHEM.2022.117683.5305

©2022 National Information and Documentation Center (NIDOC)




Figure 2: SEM of normal yellow mustard and after exposed to UVA

4.2 Molecular structure

Figure 3 shows IR spectrum of yellow mustard, a plot of wave number (X- axis) vs. present transmittance (Y- axis). IR analysis of yellow mustard listed in Table 2 show transmittance the intensity increased

after exposure to UVA at 5 cm for 1 and 4 hours but decreased at 20 cm for 1 and 4 hours. A significant change occurred in the main peak position, O-H, after exposure to UVA for 1 hour at 5 cm, but a little variation is occurred during other exposure times at 5 and 20 cm distance. That is because the absorption of UVA in cell, break or modify position or degrade some molecular bond or switching off of the transcription-translation machinery during radiation exposure [12].




Figure 3: IR spectrum of yellow mustard after exposure to UVA

Table 2: IR spectrum analysis of yellow mustard after exposure to UVA

Control (untreated sample)					
Position	Intensity %		Band		
1746	45.6		C- O		
2925	31.04		C-H		
3422	69.4		O-H		

1 hour at 5 cm			4 hour at 5 cm		
Positio n	Intensit y %	Ban d	Positio n	Intensit y %	Ban d
1746	50.4	C- O	1746	54.12	C- O
2925	34.36	C-H	2925	38.11	C-H
3450	78.4	O-H	3423	74.26	O-H

1 hour at 20 cm			4 hour at 20 cm		
Positio n	Intensit y %	Ban d	Positio n	Intensit y %	Ban d
1746	8.32	C- O	1746	32.9	C- O
2925	2.75	C-H	2925	18.21	C-H
3421	19.22	O-H	3421	57.37	O-H

4.3 Glutathione content

The glutathione which one of the most important non-enzymatic antioxidants. Glutathione content for yellow mustard is increased after exposure to UVA at 5 and 20 cm for different period of times as seen in Table 3 and Figure 4. The results show it is increased by 17.44%, 19.3%, 7.8% and 4.58% and, by 14.68%, 19.3%, 0.45% and 5.5% at 5 and 20 cm distance far from UVA source for different period times. That is because is positively affected by radiation as a hermetic type of response under applied of radiation [13, 14].




Figure 4: glutathione in yellow mustard after exposed to UVA

Table 3: glutathione content of yellow mustard after exposed to UVA

Exposure time (hour)	GSH (mg/ 100g)	
Zero (Control)	72.66	
	5 cm	20 cm
1	85.33	83.33
2	86.66	86.66
3	78.33	72.99
4	75.99	76.66

4.4 Phenolic content

Phenolic compounds are plant secondary metabolites that hold an aromatic ring bearing at least one hydroxyl groups. Total phenolic content for yellow

mustard increased by 32.57%, 43.5%, 19.5% and 48.35% and by 18.17%, 8.31%, 59.71% to 1.82.5% after exposure to UVA at distances 5 and 20 cm for 1, 2, 3 and 4 hours as shown in Table 4 and Figure 5. That is because UV radiation increases the accumulation of phenolic compounds along with antioxidant properties. Also a change in the intensity and position of O-H band caused increased in phenolic. The other results also shown phenolic compounds is increased or induced after UV radiation exposure during different period times [15- 19].




Figure 5: total phenolic content in yellow mustard exposed to UVA

Table 4: total phenolic content of yellow mustard exposed to UVA

Exposure time (hour)	Total phenolic (mg/ 100g)	
Zero (Control)	409.48	
	5 cm	20 cm
1	542.86	483.89
2	587.57	443.50
3	489.33	654.21
4	607.44	416.92

4.5 Flavonoids content

Flavonoid is a term that is a bit ambiguous literally it means flavone-like compound. Table 5 and Figure 6 show total flavonoids in yellow mustard decrease after exposure for 1, 3 and 4 hours and 1 and 3 hours at 5 and 20 cm from UVA source but it increased after exposure for 2 hours at 5 cm distance and 2 and 4 hours at 20 cm. That is because flavonoids is plant secondary metabolites that hold an aromatic ring bearing at least one hydroxyl groups which changed as shown in IR analysis.




Figure 6: total flavonoids content in yellow mustard exposed to UVA

Table 5: total flavonoids content of yellow mustard exposed to UVA

Exposure time (hour)	Total flavonoids (mg/ 100g)	
Zero (Control)	297.6	
	5 cm	20 cm
1	148.45	141.95
2	395.71	446.75
3	198.54	236.8
4	183.07	422.64

4.6 Total proline content

Proline plays important roles in protein synthesis and structure, metabolism and nutrition. Total proline content in yellow mustard decreased after exposure to UVA for different period times and distances as shown in Table 6 and Figure 7. The chemical composition of yellow mustard such as protein changed after exposure to UVA. That is because proline is a substrate for protein synthesis. Also proline accumulation is response to biotic and abiotic stresses caused to the effect of UVA.




Figure 7: total proline content in yellow mustard exposed to UVA

Table 6: total proline content of yellow mustard exposed to UVA at 5 and 20 cm for different period times

Exposure time (hour)	Total proline (mg/ 100g)	
Zero (Control)	925.84	
	5 cm	20 cm
1	889.96	899.24
2	883.56	862.40
3	917.54	911.96
4	914.41	888.98

4.7 DPPH scavenging activity

UVR carries higher energy than visible light and its effects on tissues include gene mutations, DNA damage, oxidative stress immunosuppression, and inflammatory responses. Therefore, some compounds increase and others may decrease due to abiotic stress as pathways for secondary metabolite production are interrelated. DPPH scavenging activity (%) for yellow mustard decreased after exposure to UVA for 1, 2, 3 and 4 hours at 5 and 20 cm as shown in Table 7 and Figure 8.




Figure 8: DPPH scavenging activity of yellow mustard exposed to UVA

Table 7: DPPH scavenging activity (%) of yellow mustard exposed to UVA

Exposure time (hour)	DPPH scavenging activity (%) 5 cm		
	0.4%	1%	2%
Zero (Control)	53.90	64.89	85.82
1	30.56	58.33	67.01
2	26.67	39.56	62.85
3	27.8	39.03	47.43
4	26.04	40.60	60.76

Exposure time (hour)	DPPH scavenging activity (%) 20 cm		
	0.4%	1%	2%
Zero (Control)	53.90	64.89	85.82
1	23.14	31.25	57.29
2	29.51	51.39	68.75
3	29.17	35.07	64.24
4	24.31	34.03	63.54

4.8 Tocopherol content

Table 8 and Figure 9 show, tocopherol content in yellow mustard decreased by 10.1%, 6.01%, 47.5% and 42.6% after exposure to UVA at 5 cm for 1, 2, 3 and 4 hours. Also it decreased gradually after exposure for 1, 2, 3 and 4 at 20 cm.




Figure 9: tocopherol content in yellow mustard exposed to UVA

Table 8: tocopherol of yellow mustard exposed to UVA

Exposure time (hour)	Tocopherol (mg/ kg)	
	Zero (Control)	2029.19
		5 cm 20 cm
1		1824.13 1965.43
2		1907.79 1726.63
3		1065.43 1335.00
4		1165.43 1234.16

Conclusion

Variation in radiation resulted in, significant differences in biomass accumulation, protein and total non-structural carbohydrates contents in plants. Exposure to UVA improve/or disprove some antioxidants for yellow mustard. Also UVA caused a change in yellow mustard internal structure.

References

- Xuan T.D, Khanh T.D, Khang D.T, Quan, N.T, Elzaawely A.A., Changes in chemical composition, total phenolics and antioxidant activity of *Alpinia* (*Alpinia zerumbet*) leaves exposed to UV. *Int. Lett. Nat. Sci.* 55, 25-34(2016).
- Yin X., Singer S.D., Qiao H., Liu Y., Jiao C., Wang H., Li Z., Fei Z., Wang Y., Fan C., Insights into the mechanisms underlying ultraviolet-C induced resveratrol metabolism in grapevine (*V. amurensis* Rupr.) cv. "Tonghua-3". *Front. Plant Sci.* 7, 503(2016).
- Hideg E., Jansen M. A., Strid A., UV-B exposure, ROS, and stress: inseparable companions or loosely linked associates? *Trends Plant Sci.* 18(2), 107-115(2013).
- Bornman J. F., Barnes P. W., Robinson S. A., Ballaré C. L., Flint, S. D., Caldwell, M. M., Solar ultraviolet radiation and ozone depletion-driven climate change: effects on terrestrial ecosystems. *Photochem. Photobiol. Photochem. Photobiol. Sci.* 14(1), 88-107 (2015).
- El-Bediwi A.B, Hasanin S, Abdelrazek A, El-Shora H.M., Influence of UVC on growth behavior, internal structure, enzymes and free radical of *Nigella Sativa* plant *R. Rev. BioSci.*, 13(2), 1-13(2018).
- El-Bediwi A.B, Hasanin S, Abdelrazek A, El-Shora H.M., Effect of ultraviolet on morphological and secondary metabolites content of *garden cress* *IJSRSET*, 4(1), 187- 194 (2018).
- Zhang W.J, and Björn L.O., *Fitoterapia*, 80 (4), 2009.
- Hectors K, Van Oevelen S, Geuns J, Guisez Y, Jansen M.A.K and Prinsen E., *Physiol. Plant.* 152(2), 219-230(2014).
- Ghasemi S, Kumleh H.H and Kordrostami M., *Protoplasma*. 256(1), 279-290 (2019).
- Kovács E and Keresztes A., Effect of gamma and UV-B/C radiation on plant cells. *Micron*, 33(2), 199- 210(2002).
- Ashraf M, Cheema A.A, Rashid M, and Qamar Z., Effect of g-rays on M1 generation in basmati rice. *Pak. J. Bot.* 35, 791–795 (2003).
- Sen Raychaudhuri S and Deng X.W., The role of superoxide dismutase in combating oxidative stress in higher plants. *Bot. Rev.* 66(1), 89-98 (2000).
- Štajner D, Popovic B, and Taški K., Effects of g-irradiation on antioxidant activity in soybean seeds. *Cent. Eur. J. Biol.* 4(3), 381-386(2009).
- Chakravarty B and Sen S., Enhancement of regeneration potential and variability by g-irradiation in cultured cells of *Scilla indica* *Biol. Plant.* 44(2), 189-193(2001).
- Surjadinata, B. B., Jacobo-Velázquez D.A. and Cisneros-Zevallos L., UVA, UVB and UVC light enhances the biosynthesis of phenolic antioxidants in fresh-cut carrot through a synergistic effect with wounding. *Molecules*. 22(4), 1-13(2017).
- Teoh, L. S., Lasekan. O, Adzahan N.M., and Hashim. N., The effect of ultraviolet treatment on enzymatic activity and total phenolic content of minimally processed potato slices. *J. Food Sci. Technol.* 53(7), 3035-3042(2016).
- Ullah, M.A., Tungmunthum D, Garros L, Drouet S, Hano C and Abbasi B. H., Effect of ultraviolet-C radiation and melatonin stress on biosynthesis of antioxidant and antidiabetic metabolites produced in in vitro callus cultures of *Lepidium sativum* L. *Int. J. Mol. Sci.* 20(7), 1787 (2019).
- Zagoskina NV, Alivia AK, Gladysenko TO, Lapshin PV, Egorova EA, Bukhov NG. *Russ J Plant Physiol.* 52,731-9(2005).
- Zagoskina NV, Dubravina GA, Alyavina AK, Goncharuk EA. *Russ J Plant Physiol.* 50, 270(2003).