Micronutrient Intake Profile of Egyptian Women in Reproductive Ages

Salwa M. Saleh MD¹; Hanaa H El-Sayed Ph.D.²; Hoda M El Gezery Ph.D.¹ and Asmaa EM Mostafa Ph.D.¹

1. Department of Field Studies and Survey- National Nutrition Institute, Cairo, Egypt

2. Department of Nutritional chemistry and metabolism- National Nutrition Institute, Cairo, Egypt

Corresponding author: Asmaa E. Mostafa, Tel: +20 101 512 3186. E-mail: dr_asmaa707@yahoo.com

ABSTRACT

omen in low-and middle-income nations face major public health issues related to micronutrient deficiencies. From every region of Egypt, samples of women of fertile age who were neither pregnant nor lactating were chosen at random in several governorates. 4694 women between the ages of 18 and less than 50 years old participated in this study from 2015-2020. Using data from the participant's social relationships, daily consumption, and anthropometric measures, the micronutrient nutritional status was assessed. The results showed that most of the sample participants were between the ages of 25 and 35. In upper Egypt districts women made up 75.1% of university graduates. The frontier regions had the highest numbers of illiteracies, whereas Lower Egypt had the lowest. The highest percentages of professionals working were found in urban areas (37.1%) and upper Egypt (32.4%). Female unemployment was the highest in frontier areas (58.2%). The sample was overweight (29.1%) and obese (49.0%). Females consumed more minerals than they needed, such as sodium and copper. While intake of potassium, calcium, and magnesium fell below 50% of the reference daily intake (RDI). Furthermore, the utilization of phosphorus, iron, and zinc fell within the permissible ranges of 77-81%, 77-84%, and 92-105%, respectively. Upper Egypt regions were found to have the lowest vitamin A intake (51%) while the frontier region had the highest rate (122%). All water-soluble vitamins, with the exception of vitamin B1, were above 75% of RDI. This study indicated present poor dietary practices among women and their need for additional nutrition education.

Keywords: Egyptian governorates- women- Micronutrient intake- National Egypt intake 2020

INTRODUCTION

Micronutrient deficiencies (MNDs) are significant public health issues that affect а significant portion of global citizens (Manjeru and Others They prevalent 2017). are worldwide, with the largest danger being too expectant mothers and their young children under the age of five (Bailev and else searchers **2015).** Around the world, there are over 800 million malnourished persons and 1.5 to 2 billion individuals who have one or more chronic MNDs. Notable mineral and vitamin deficiencies include those of calcium, iodine, iron, zinc, selenium, and vitamin A and folate (Beal et al., 2017; FAO 2020; Kumssa and Another's 2015; WHO and UNICEF 2017). Deficits in any one of these micronutrients can cause major issues like greater sensitivity to diseases, congenital disabilities, blindness. stunting, mental impairment, subpar academic and professional performance, and even mortality (Petry et al., 2016; Black, 2003; Zimmermann 2009; Roth and Others 2018: van Hagen et al., 2017). Young children, pregnant and lactating adolescent girls, women, and women of fertile age are more

susceptible to the effects of micronutrient insufficiency because of their high requirements. It has been demonstrated that improving diets, fortifying staple foods and condiments. and biofortification. and supplementing populations with increased needs to raise their intake of micronutrients can reduce mother and infant illness, cognitive impairment, and death (Bhutta et al., 2008; Bhutta et al., 2013; Keats et al., 2021).

However, it is difficult to estimate the prevalence and population of people who are micronutrient deficient globally. This is partly because most micronutrient deficiencies go untreated since they have vague symptoms, which is why they are frequently referred to as hidden hunger. Additionally, populationbased surveys rarely contain biomarkers for micronutrient leaving significant status. a knowledge gap regarding the burden of micronutrient deficiency both nationally and globally (Brown et al., 2021). Based on the anticipated number of anemic individuals. the nutrition community has reported for more than three decades that 2 billion people worldwide suffer from micronutrient deficiencies (WHO, **1991**). To determine financing and programming priorities and to support advocacy activities to advance these interests, reliable and well-documented global and estimates of regional persons from micronutrient suffering deficiencies are necessarv (Haddad et al., 2015). There are still few population-representative markers for micronutrient status, and studies that are not representative are not very useful for calculating burden. For preschool-aged children (aged 6-59 months) and **non-pregnant** women of reproductive age, there are few and frequently out-of-date statistics available, but very few population-representative surveys include data on school-aged adolescent children. males. pregnant women, men, and older persons (WHO,2020).

This study seeks to offer a distinctive perspective on the food and micronutrient intake of a sizable cohort of Egyptian women who are of reproductive age. The study goals were to (1) describe these women's micronutrient intakes, (2) evaluate the sufficiency of those intakes in comparison to references, and (3) pinpoint

significant dietary sources of micronutrients.

Study design: Cross section study was employed in this study.

Sample size: This study was carried out on 4694 women, whose ages ranged from 18 to 50 years old. This study was undertaken as part of a study of the national survey done by the National Nutrition Institute (NNI) Egypt for updating in food consumption patterns among different population groups during the period from 2015 to 2020. Egypt was divided into four geographical areas as follows: Urban, Lower, Upper, and Frontier Egypt. The following governorates were selected randomly from each geographical area Urban (Cairo, Alexandria, Suez, and Port Said), while Lower Egypt (Ismailia, Minofyia, Kafr El-Shiekh, Damietta, Oalvubia, Sharkia, Dakahlia, Beheira, and Gharbia), Upper Egypt (Giza, Minya, Sohag, Assuit, Qena, Fayoum, Beni surf, Aswan and Luxor) and Frontier (North Sinai, Matrouh, Red Sea, New Valley, and South Sinai). The sample was selected from the nominated governorates randomly as follows: -The districts were selected randomly. Each studied governorate except Cairo and Alexandria was divided into urban and rural sampling units. All residence areas had different socioeconomic classes.

Exclusion criteria: Samples with specific nutritional problems and specific dietary recommendations (vegetarian, pregnant, lactating,

Diabetes, chronic renal diseases, chronic liver diseases).

Study tools: The tools of this study consisted of structured inter-viewing questionnaires. This consists of three parts: The first is to elicit the socioeconomic characteristics of women including information about their age, educational levels, and occupation (Park and Park, 1979). The second is to collect daily food intake by using the 24-hour recall method and food frequency. The third recorded anthropometric measurements of studied women.

Ethical considerations: the permission of the ministry of health and population research ethics committee.

METHODOLOGY:

Dietary intake of nutrients: The sample weighting procedure and the 24-hour recall to record the daily intake. On the day before the interview day, the field team acquired oral reports about the meals and drinks the target participants ingested at each dining event. It started with the first meal consumed after waking up and continued with each successive meal until sleep. Utilizing a dietitian's kit of utensils with known weights

and supported by weighing comparable portions of what was consumed, amounts of foods and beverages were calculated. the nutritive value of the intake was determent using food composition tables for Egypt (**NNI**, **2006**) **and** compared with the per capita RDI used by **FDA** (**2020**) according to age and physical state.

- Dietary habits: "Food Frequency Questionnaire": This method was used to collect qualitative data about the habitual food and beverage consumption habits of the family members per week (daily, less than 3 times per week, or equal to or more than 3 times per week) and monthly.
- Anthropometric measures: According to WHO (2006) guidelines, anthropometric data, include length, body weight, and body mass index (BMI).
- Statistical analysis: The SPSS version 20 program was used to analyze the data. Data were expressed in numbers and percentages for qualitative nonparametric variables the median

percentiles were used while in quantitative parametric data, the mean \pm standard deviation (SD) was used. The Chisquare test also was used for qualitative variables to determine the *P*-values and confidence interval (CI), T-ANOVA. and test. the Pearson correlation test was also used. Significant statistical correlations between different variables were detected by using the Chisquare test. The statistical significance correlation reaches if the *P*-value is less than 0.05according to Levesque (2007).

Dietary data analysis: Data • from 24-hour recall was analyzed using the food analyzing program (using Egyptian food composition 2006 as the database) in the Statistics Unit at the National Nutrition Institute. The result for each subject was concentrated on the amount of micronutrient intake consumed on this day. Data obtained from this program was then first compared to the RDI check for to the adequacy of intake and then the intake of each group was

compared to the others using an independent sample t-test or one-way ANOVA test.

RESULTS and DISCUSSION

Characteristics of the Participants:

In Figure (1) the participants in the survey samples examined the 4 main demographic which included 4694 areas. Egyptian women of reproductive age. In Lower Egypt, there are more samples (1714). The majority of the sample's women had the ages 25-35 years (figure 2). Among the four demographic groups, there was a considerable variation in the education of women in figure (3). Women had the highest percentage of university graduates showed in upper Egypt areas (75.1%), while the lowest percentage was found in frontier and lower Egypt areas (3.4% and 8.3%) respectively. Despite this, lower Egypt had the lowest percentage of illiteracy and the frontier regions had the highest. The majority of the sample were attending secondary education. Relating to women's employment, the 4 demographic areas differed significantly from one to another. Urban regions and upper Egypt had the largest proportions of professionals working, with 37.8%

and 37.6%, respectively. However, just 5.9% of professional occupations were found in frontier areas, which is the lowest rate. Frontier areas had the highest ratio of female unemployment (58.2%), whereas lower Egypt had the lowest percentage (28.4%), their results showed in (**figure 4**).

The perchance distribution of women's BMI correlated with geographic dispersion, was shown in Table (1). According to BMI calculations, the majority of the sample representatives had obese (49%). and (0.77%)were underweight. On the report of the data, there is no relationship between urban and rural areas with women's BMIs an $(X^2 \text{ value equal})$ (0.215). Rural areas had a higher percentage of underweight to normal weight than urban areas did. An indication of poor nutritional habits among women of that age and the need for more nutrition women's education programs is the fact that around 49% of Egyptian women and 29.1% are overweight.

From the food frequency questionnaire, **Figure 5** demonstrated the mean food groups intake of households (per capita) in urban, lower, and upper Egypt and frontier regions. Cereals and tubers revealed the highest daily item intake (380.5g and 94.6g), as a good source of calories. Extra calories are deposited as fat throughout the body. Adipose tissue with excessive amounts of fat cells can harm health, and an energy imbalance between calories ingested and calories burned is what leads obesity and to overweight (WHO, 2014).

The participants' dietary intake of vitamins and minerals

The importance of micronutrient consumption bv women of reproductive age nonand lactated pregnant was highlighted by this study. The data seen in the table (2) indicated that, for the majority of participants from different areas, mean sodium intake was more than three times higher than the requirements, exceeding the recommended daily intake of 2.3 g /day (FDA, 2020). It is commonly recognized that consuming more salt causes high blood pressure, which in turn causes heart disease, stroke, and kidney damage (Cappuccio, 2013). In the Nordic cohorts of Europe, an independent relationship between sodium intake and the risk of coronary heart disease has been described.

Higher salt intake of 100 mmol was associated with 51% higher death from coronary heart disease, 45% higher mortality from cardiovascular disease, and 26% higher mortality from all causes. (Tuomilehto et al., 2001). Since too much sodium is linked to elevated blood pressure, stroke, and body water retention, sodium consumption should be restricted to a maximum of requirements. The use of cheese and meat products like sausages may be to blame for the study participants' higher sodium intake. These goods often contain more than 1 g of salt per 100 g of the item, which is a large quantity of salt (and thus sodium) (Farquhar et al., 2015). Conversely, the average daily intake of sodium is still 4.0 g or more in many regions (Hyseni, et al., 2017; Vasara, et al., 2017). Population-based prospective studies reveal that the risk of cardiovascular disease rises by 18% for every daily increase of 1000 mg in salt excretion (Ma et al., 2022). Therefore, the WHO (2013)recommends 30% а reduction in salt intake in the Plan Global Action for the Prevention and Control of Noncommunicable Diseases NCDs. Planning and evaluating the

success of national salt reduction strategies requires an evaluation of the population's salt consumption (WHO, 2020). In the current population-based investi-gation, salt intake was significantly higher than advised. The critical need for coordinated action to address salt consumption is reiterated by this in Egypt.

For the study samples, the computed potassium consumption was less than half of the daily dose of 4700 mg advised by the FDA (2020). However, plant-based food items like fresh and dried fruits, legumes, green leafy vegetables, etc. are easy to consume for potassium, studies from the United States (Pratt, et al., 2014) and Poland (Bzikowska-Jura et al., **2018**) have found that in lactating women's potassium intake is lower than that advised by the national nutrition guidelines. In order to decrease the risk of CVDs in adults. such hypertension, as coronary heart disease, and stroke, the WHO also suggests increasing potassium consumption from meals (WHO, 2012, 2014; D'Elia, et al., 2011). However, the risk of CVD can be predicted more accurately by the sodium-topotassium ratio in 24-hour urine collections rather than by dietary

consumption alone (Stone et al, 2016). A combination of low sodium and high potassium intake may be a more accurate indicator of the risk of cardiovascular disease than sodium and potassium intake separately, therefore measuring the ratio is crucial. The ideal Na/K ratio when adhering to recommended intakes is roughly one, according to the WHO (2012).

Table (2) showed that the main Egyptian demographic groups varied significantly in their calcium intake. Lower Egypt was found to have the highest calcium intake, while frontier regions had the lowest calcium intake. The amount of calcium consumed in this trial was only a third of the RDI of "1300 mg." According to questionnaire, the frequency women consumed less than half a cup of milk daily. Adults' bone health may be negatively impacted by low calcium consumption. To create national programs to maximize consumption, it is vital to acknowledge the existence of low calcium intake (Balk et al, **2017).** According to the available data. greater dietary calcium consumption is linked to lower body weight, BMI, and obesity. The activation of lipolysis and

suppression of lipogenesis may be the mechanism through which dietary calcium reduces body fat (**DeJongh et al., 2006**). These notes are shown in the study's sample. It has been demonstrated that dietary salt intake increases urine calcium loss and consequently increases calcium requirements in those with highsodium diets.

In all of the selected regions, women's phosphorus (P) usage ranged from 77 to 81% of RDI; this ratio was satisfactory. Additionally, thev had significantly lower mean calcium and phosphorus levels, both of which are crucial for maintaining strong bones (Alshammari et al., **2017).** By changing cytokine levels, increased dietary phosphorus consumption may have an impact on inflammatory illness. Phosphates are linked to a number of diseases, including premature aging and vascular calcification; this association may be due to an augmented inflammatory response (Goodson et al., 2019). Despite the fact that phosphate insufficiency is probably uncommon, there is some weak evidence that it can be caused by dietary deficiencies. In fact, average dietary intakes of P and magnesium (Mg) are typically far

higher than their biological needs, and if not, enhanced absorption is the result (Waterlow, 2006).

According to FDA (2020). the average amount of Mg consumed was 40% of the RDI (420 mg). Mg is a critical dietary component for people since it is engaged in important biological Neuromuscular processes. conduction abnormalities, clinical and hypocalcemia, illnesses affecting the metabolism of the muscles and bones, such as osteoporosis and loss of muscle all linked mass. are to hypomagnesemia. Α national survey of 4497 American adults found that higher levels of systemic inflammation were linked to reduced magnesium intake (Kim et al., 2010). A meta-analysis of 32,918 adults found that elevated serum C-reactive protein (CRP) were linked to levels poor magnesium intake (Dibaba et al., 2014). Numerous adult crosssectional research found links between magnesium consumption and increased muscle size and strength. (Welch et al., 2016; Hayhoe et al., 2019). Magnesium and calcium compete for binding calcium-sensing receptors. on which lowers serum parathyroid hormone (PTH) levels (Mazur et al., 2007). Chronically elevated serum PTH is linked to muscle loss by negatively affecting muscular

energy metabolism, which reduces the generation of amino acids and protein synthesis (Garber 1983). The insulin signaling pathways may also play a role in how magnesium consumption controls inflammation and muscle mass. It is crucial for appropriate insulin secretion and activity, as well as for the best performance of numerous enzymes involved in glucose and energy metabolism, to keep serum Mg²⁺ concentrations within the reference range. Insulin resistance is correlated with low serum magnesium levels (Chutia and Lynrah, 2015). Mg⁺² can control pancreatic beta cells' release of insulin (Kostov 2019).

Table 2 also indicated that the urban area had the lowest iron intake but that Upper Egypt had the highest. With a variance of 77-84%, women's consumption of iron is lower than their RDI. Women are prone to issues including fatigue as well as a variety of micronutrient deficiencies as a result of the poor diet brought on by meals high in calories but low in nutrients (Daniels et al., 2005). Low dietary consumption in developing nations is related to both the diet's poor iron bioavailability as well as its low iron content. Chronic inflamemation. inflammatory bowel disease (IBD), congestive heart failure. chronic renal disease. obesity, cancer, and rheumatoid arthritis all interfere with iron absorption (Gilreath. et al., 2014: Lopez, et al., 2016). It is wellaccepted that the incidence of iron insufficiency is widespread in poor nations due to low intakes of animal products and diets rich in phytates. which prevent iron absorption (de Benoist, et al., **2007**). The primary contributor to micronutrient deficiency is a diet heavy in staple foods and deficient in animal sources (Bouis, 2003). A deficiency is brought on by excessive loss or insufficient iron absorption. Depending on the individual's iron level, the type of iron, and other dietary parameters, absorption is strictly controlled in the intestines (Beard 2007).

Lower Egypt and urban regions had the lowest zinc intake (92%) of RDI, whereas upper and frontier Egypt reported the highest zinc capacity (100-150% of RDI). Many essential metalloenzymes in the body, such as carbonic anhydrase and alkaline phosphatase, have zinc as an essential component of their catalytic site. The structural differentiation of numerous organs, including the skin, highly depends on zinc finger proteins, such as those found in

retinoic acid and vitamin D receptors. Additionally, zinc functions as an ionic signal that controls gene expression. It is abundant in foods high in protein, particularly those high in animal protein. Phytates and iron diminish their bioavailability (**Maverakis et al., 2007**).

According to FDA (2020), lower Egypt had the highest reported copper intake at 153% of the RDI (0.9 mg/day), followed by urban and higher Egypt at 131%. Nuts, organs meat, fish, legumes, dried fruit, particularly cashews, and vegetables are the main dietary sources of copper. A significant source of copper aside from diet is water, albeit the metallic content of varies greatly. The water concentration of copper in water is determined by elements such as natural mineral content, pH, and plumbing system (National Research Council, US 2000). High copper concentrations have been reported in numerous nations, including Europe. When the concentrations have been measured multiple times, there is significant variation between the recorded amounts even within the sources. The maximum same permissible level of copper in water for human consumption in

the European Union (EU) is 2 mg/L (EFSA, 2015). The primary function of copper is the formation oxidases. which of transport electrons to reduce molecular oxygen. As a result, copper is essential for energy metabolism at the cellular level (Kubiak, et al., **2010**). Additionally, the presence of copper promotes mesenchymal stem cells' development into the osteogenic pathway. (Ding, et al., 2014).

 Table 3 demonstrated that
there were notable variations in vitamin A intake among Egypt's main demographic groups. The districts in upper Egypt had the lowest vitamin A intake. In the frontier region, the RDI percentage reached its greatest point at 122%. Immune system health, cellular communication, growth, and development, also male and female reproduction are all impacted by vitamin A (Blaner, 2020). Supporting cell growth and differentiation, vitamin A is essential for the healthy development and upkeep of the heart, lungs, eyes, and other organs (Ross, **2014**). Because it supports the differentiation appropriate and operation of conjunctival the membranes and cornea, as well as being a necessary component of rhodopsin, the light-sensitive protein in the retina that reacts to light entering the eye, vitamin A is also vital for vision (Carazo et al., 2021).

Infection risk and intensity are increased by vitamin A deficiency (VAD) (Stevens, et al., 2015). One of the leading causes of preventable childhood blindness is vitamin A deficiency (Bailey et al., 2015). Additionally, a chronic vitamin A deficit has been linked to incorrect lung development, respiratory conditions, a higher risk of anemia, and even mortality (Wiseman et al., 2017: Timoneda et al., 2018). The body retains excess vitamin A because it is fat soluble, particularly in the liver, where levels can build up. After taking one or more very high doses of vitamin A (usually more than 100 times the RDA), a person will have acute vitamin A toxicity, also known as hypervitaminosis A, within days to weeks (NIDDKD, 2020). Severe headaches, blurred vision, nausea, dizziness. aching muscles. and difficulties with coordination are frequently the resulting signs and symptoms. When conditions are extreme, an increase in cerebral spinal fluid pressure can cause drowsiness, eventually a coma, and even death 2020). (NIDDKD. Chronic hypervitaminosis A (consumption of excessive dosages of vitamin A on a regular basis) can result in abnormal liver test results as well as dry skin, sore muscles and joints, weariness, and depression. (NIDDKD, 2020).

Table 3 showed thatdifferent regions had vitamin Cintake that was between 61% and

92% of the RDI (90 mg/day). Vitamin C plays a role in protein metabolism as well as being necessary for the manufacture of collagen, L-carnitine, and certain neurotransmitters (Li and Schellhorn, 2007). Collagen is a crucial component of connective tissue, which is crucial for the healing of wounds. Vitamin C is also а vital physiological antioxidant (Frei et al., 1989), is crucial for immune system health (Jacob and Sotoudeh, 2002), and enhances nonheme iron absorption (Gershoff, 1993). Data in table 3 also showed that vitamin B1 is consumed more than RDI in Upper Egypt and equal RDI in lower Egypt, while the frontier area was 96% and the urban was 95%. Humans store thiamin primarily in the liver, but in very small amounts. The vitamin has a short half-life, so people require a continuous supply of it from their diet (Bemeur and Butterworth 2014).

Between the main demographic groups in Egypt, appreciable there were no differences in B2 consumption, and it was within the acceptable range of >75% from RDI (1.2mg/day). Blood homocysteine levels are kept at normal levels with the aid of riboflavin B2 (**Rivlin, 2010**). only little levels of riboflavin in the kidneys, liver, and heart. If too much is taken, it either isn't absorbed or the little that is absorbed is eliminated in urine (**IMFNB, 1998**). The quantity of free riboflavin that the large intestine can absorb depends on the diet and is produced by bacteria there. After consuming foods with a vegetable base rather than a meat base, more riboflavin is created (**Said and Ross 2014**).

The prevalence of obesity and rising vitamin intake may be positively correlated (**Zhou and Zhou, 2014**). Because B vitamins can enhance fat synthesis (**McHenry and Gavin, 1941**), research has shown that higher B vitamin intake (B1, B2, and niacin) is closely connected with the prevalence of obesity and diabetes (**Zhou et al., 2010**).

CONCLUSION

The education levels of the study sample were different with 39.2%, females had the highest secondary education rate, and roughly a quarter of females were litterateurs. Urban areas had the highest proportion of residents with a university degree. Frontier regions had a greater rate of

illiteracy. In terms of employment status, (68%) of the ladies were unemployed. Most adult females consumed insufficient amounts of potassium, calcium, magnesium, vitamin A. and other minerals. All adult females received enough vitamin C to meet their needs. The intake of sodium, copper, and vitamin B1 exceeded the needs. While consumption of potassium, calcium, and magnesium fell below 50% of the recommended daily amounts. Furthermore, the utilization of phosphorus, iron, and zinc fell within the permissible ranges of 77-81%, 77-84%, and 92-105%, respectively.

RECOMMENDATION

Many of these nutritional deficits can be avoided by learning about nutrition and eating a healthy, balanced diet that includes a variety of food types that are sufficient for each nutrient. When it comes to factors that affect food intake and dietary practices, such as eating habits, food purchasing, food preparation, food safety, and environmental factors, nutrition education has the power to improve dietary behavior and nutritional status. For nations afflicted by globalization, urbanization, and a risky dietary shift toward

inexpensive processed foods high in sugar, fat, and salt, proper nutrition education is essential.

REFERENCES

Alshammari E; Suneetha E; Adnan M; Khan S and Alazzeh A (2017):

> Growth Profile and Its Association with Nutrient Intake and Dietary Patterns among Children and Adolescents in Hail Region of Saudi Arabia. *BioMed Research International,* vol. 2017, 1-9

Bailey RL; West KP and Black RE (2015):

The epidemiology of global micronutrient deficiencies. *Annals of Nutrition and Metabolism* 66 (Suppl. 2):22–33

Balk EM; Adam GP; Langberg VN; Earley A; Clark P; Ebeling P R; Mithal A; Rizzoli R; Zerbini CAF; Pierroz DD; and Dawson-Hughes B (2017):

Global dietary calcium intake among adults: a systematic review. *Osteoporos* Int 28, 3315–3324

Beal TE; Massiot JE; Arsenault M; Smith R and Hijmans RJ (2017):

Global trends in dietary micronutrient supplies and estimated prevalence of inadequate intakes. *PLoS One* 12 (4): e0175554

Beard J (2007):

Manual Reference for Laboratory Considerations -Iron Status Indicators for Population Assessments. WHO: Geneva. Switzerland: Indicators of the iron status of populations: Free erythroncyte protoporphyrin and zinc protoporphyrin; serum and plasma iron, total iron binding capacity and transferrin saturation; and serum transferrin receptor.

Bemeur C and Butterworth RF (2014):

Thiamin. In: Ross AC, Caballero B, Cousins RJ, Tucker KL, Ziegler TR, eds. Modern Nutrition in Health and Disease. 11th ed. Baltimore, MD: Lippincott Williams & Wilkins:317-24. Bhutta ZA; Ahmed T; Black RE; Cousens S; Dewey K; Giugliani E; Haider BA; Kirkwood B; Morris SS; Sachdev HP and Shekar M (2008):

> Maternal and Child Undernutrition Study Group. What works? Interventions for maternal and child undernutrition and survival. *Lancet*. 2;371(9610):417-40.

Bhutta ZA; Das JK; Rizvi A; Gaffey MF; Walker N; Horton S; Webb P; Lartey A and Black RE (2013):

Lancet Nutrition Interventions Review Group, the Maternal and Child Nutrition Study Group. Evidence-based interventions for improvement of and maternal child nutrition: what can be done and at what cost? Lancet. 3;382(9890):452-477.

Black R (2003)

Micronutrient deficiency an underlying cause of morbidity and mortality. *Bull World Health Organ*; 81:79–79.

Blaner WS (2020):

Vitamin A and Provitamin A Carotenoids. In: Marriott BP, Birt DF, Stallings VA, Yates AA, eds. Present Knowledge in Nutrition. 11th ed. Cambridge, Massachusetts: *Wiley-Blackwell;*73-91.

Bouis HE (2003):

Micronutrient fortification of plants through plant breeding: can it improve nutrition in man at low cost? *Proc Nutr Soc.*; 62: 403–11.

Brown KH, Moore SE, Hess SY,

McDonald CM; Jones KS; Meadows SR; Manger MS; Coates J; Alayon S and Osendarp SJM (2021):

> Increasing the availability and utilization of reliable data on population micronutrient (MN) status globally: the MN Data Generation Initiative. *Am J Clin. Nutr.;* 114: 862–70.

Bzikowska-Jura A; Czerwonogrodzka-Senczyna A; Ol edzka G; Szostak-Wegierek D; Weker H and Wesołowska A (2018):

Maternal Nutrition and Body Composition During Breastfeeding: Association with Human Milk Composition. *Nutrients*, 10, 1379

Carazo A; Macakova K; Matousova K; Krcmova LK; Protti M and Mladenka P (2021):

Vitamin A update: Forms, sources, kinetics, detection, function, deficiency, therapeutic use, and toxicity. *Nutrients;* 13: 1703.

Chutia H and Lynrah KG (2015):

Association of Serum Magnesium Deficiency with Insulin Resistance in Type 2 Diabetes Mellitus. *J. Lab. Physicians* 2015, 7, 75–78.

Daniels SR; Arnett DK; Eckel RH; Gidding SS; Hayman LL; Kumanyika S; Robinson TN; Scott BJ; St Jeor S and Williams CL (2005):

> Overweight in children and adolescents: pathophysiology, consequences, prevention, and treatment. *Circulation*. 19; 111 (15): 1999-2012.

DeBenoist B; Darnton-Hill I; Davidsson L; Fontaine O and Hotz C. (2007):

Conclusions of the Joint WHO/UNICEF/IAEA/IZi NCG Interagency Meeting on Zinc Status Indicators. *Food and Nutrition Bulletin.*; 28 (3_ supp 13): S480-S484

DeJongh DE; Binkley TL and Specker BL (2006):

Fat mass gain is lower in calcium-supplemented than in un-supplemented preschool children with low dietary calcium intakes. *The American Journal of Clinical Nutrition*, 84, 5,1123–1127

D'Elia L; Barba G; Cappuccio FP and Strazzullo P (2011):

Potassium Intake, Stroke, and cardiovascular disease a Meta-Analysis of Prospective Studies. J. Am. Coll. Cardiol. 2011, 57, 1210–1219.

Dibaba DT; Xun P and He K (2014):

Dietary Magnesium Intake Is Inversely Associated with Serum C-reactive Protein Levels: Metaanalysis and Systematic Review. Eur. J. Clin. Nutr., 68, 510–516

Ding H; Gao YS; Wang Y; Hu C; Sun Y and Zhang C (2014):

Dimethyloxalylglycine increases the bone healing capacity of adipose-derived stem cells by promoting osteogenic differentiation and angio-genic potential. *Stem Cells Dev.*, 23, 990– 1000.

European Food Safety Authority (EFSA) (2015):

Scientific Opinion on Dietary Reference Values for copper. *EFSA Journal*, 13:4253. DOI: 10.2903.

FAO. 2020:

Food and Agriculture Organization of the United Nations. Rome, Italy: FAO STAT database. Accessed May 15, 2020.

Farquhar WB; Edwards DG; Jurkovitz CT and Weintraub WS (2015):

Dietary sodium and health: more than just blood pressure. *J Am Coll* *Cardiol.* 17; 65 (10):1042-50.

Food and drug administration (FDA) (2020):

Reference Guide: Daily Value Changes for Nutrients.

Frei B; England L and Ames BN (1989):

Ascorbate is an outstanding antioxidant in human blood plasma. *Proc Natl Acad Sci U S A*; 86: 6377-81

Garber AJ (1983):

Effects of Parathyroid Hormone on Skeletal Muscle Protein and Amino Acid Metabolism in the Rat. J. *Clin. Investig.*, 71, 1806–1821.

Gershoff SN (1993):

Vitamin C (ascorbic acid): new roles, new requirements? *Nutr. Rev.*; 51:313-26

Gilreath JA and Stenehjem DD (2014):

Rodgers, G.M. Diagnosis and treatment of cancerrelated anemia. *Am. J. Hematol*, 89, 203–212.

Goodson JM; Shi P and Razzaque MS (2019):

Dietary	Phosphorus
Enhances	Inflammatory
Response:	A Study of
Human	Gingivitis. J.
Steroid E	Biochem. Mol.
<i>Biol.</i> 188, 1	66–171.

Haddad L: Achadi E: Bendech MA; Ahuja A; Bhatia K; Bhutta Z; Blössner M; Borghi **E**: Colecraft E; de Onis M; Eriksen K: Fanzo J: Flores-Avala R: Fracassi P; Kimani-Murage E; Nago Koukoubou E; Krasevec J; Newby H; Nugent R; Oenema S; Martin-Prével Y: Randel J: Requeio J: Shyam T: **Udomkesmalee E and Reddy KS** (2014):

> The Global Nutrition Report 2014: actions and accountability to accelerate the world's progress on nutrition. J Nutr.; 145 (4): 663-71.

Hayhoe RPG; Lentjes MAH; Mulligan AA; Luben RN; Khaw KT and Welch AA (2019):

Cross-sectional associations of dietary and circulating magnesium with skeletal muscle mass in the EPIC-Norfolk

cohort. *Clin. Nutr.*, 38, 317–323.

Hyseni L; Elliot-Green A; Lloyd-Williams F; Kypridemos C; O'Flaherty M; McGill R; Orton L; Bromley H; Cappuccio FP and Capewell S (2017):

Systematic Review of Dietary Salt Reduction Policies: Evidence for an Effectiveness Hierarchy? PLoS ONE 12.

Institute of Medicine. Food and Nutrition Board (IMFNB, 1998):

Dietary Reference Intakes: Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline. Washington, DC: National Academy Press.

Jacob RA and Sotoudeh G (2002):

Vitamin C function and status in chronic disease. *Nutr Clin Care*; 5:66-74.

Keats EC; Das JK; Salam RA; Lassi ZS; Imdad A; Black RE and Bhutta ZA (2021):

Effective interventions to address maternal and child malnutrition: an update of the evidence. *Lancet Child Adolesc Health.*; 5 (5):367-384.

Kim DJ; Xun P; Liu K; Loria C; Yokota K; Jacobs DR and He K (2010):

Magnesium Intake in Relation to **Systemic** Insulin Inflammation. Resistance, the and Incidence of Diabetes. Diabetes Care 2010, 33, 2604-2610.

Kostov K (2019):

Effects of Magnesium Deficiency on Mechanisms of Insulin Resistance in Type 2 Diabetes: Focusing on the Processes of Insulin Secretion and Signaling. *Int. J. Mol. Sci.*, 20, 1351.

Kubiak K; Klimczak A; Dziki Ł; Modranka R and Malinowska K (2010):

[Influence of copper (II) complex on the activity of selected oxidative enzymes]. *Pol. Merkur. Lekarski.*, 28, 22–25.

Kumssa DB; Joy EJM; Ander EL; Watts MJ; Young SD;

Walker S and Broadley MR (2015):

Dietary calcium and zinc deficiency risks are decreasing but remain prevalent. Scientific Reports 5:1–11.

Leo BDA; Banca Dati and di Composizione (2007):

Degli Alimenti per studi Epidemiologici in Italia. Available online: <u>http://www.bda-ieo.it</u>. (accessed on 20 Sepetember 2007).

Levesque R (2007):

SPSS programming and data management. A guide for SPSS and SAS Users, Fourth Edition, SPSS Inc., Chicago, 3.

Li Y and Schellhorn HE (2007):

New developments and novel therapeutic perspectives for vitamin C. *J Nutr*; 137:2171-84.

Lopez A; Cacoub P; Macdougall IC and Peyrin-Biroulet L (2016):

Iron deficiency anemia. *Lancet*, 387,907– 916. Ma Y; He FJ; Sun Q; Yuan C; Kieneker LM; Curhan GC; MacGregor GA; Bakker SJL; Campbell NRC; Wang M; Rimm EB; Manson JE; Willett WC; Hofman A; Gansevoort RT; Cook NR and Hu FB (2022):

> 24-Hour Urinary Sodium and Potassium Excretion and Cardiovascular Risk. *N. Engl. J. Med.* 386, 252 – 263.

Manjeru P; Van Biljon Aand Labuschagne M (2017):

The development and release of maize fortified with provitamin Α carotenoids in developing countries. Critical Reviews in Food Science and Nutrition 59 (8):1284-1293.

Maverakis E; Fung MA; Lynch PJ; Draznin M; Michael DJ; Ruben B and Fazel N (2007):

Acrodermatitis enteropathica and an overview of zinc metabolism. *J Am Acad Dermatol.*;56(1):116-24.

Mazur A; Maier JAM; Rock E; Gueux E; Nowacki W and Rayssiguier Y (2007):

Magnesium and the inflammatory response: Potential physiopathological implications. *Arch. Biochem. Biophys*, 458, 48 – 56.

McHenry EW and Gavin G (1941):

The B vitamins and fat metabolism: iv. the synthesis of fat from protein. *JBC*, Volume 138, Issue 2, 471-475

National Institute of Diabetes and Digestive and Kidney Diseases (NIDDKD, 2020):

Liver Tox: Clinical and research information on drug-induced liver injury.

National Nutrition Institute for Egypt, NNI (2006):

Food composition table based on local food analysis and food composition table in the middle east.

National Research Council. US (2000):

Copper in Drinking Water, in N. R. C. Committee on Copper in Drinking Water (Ed.), USA.

Park JE and Park K (1979):

Textbook of preventive and social medicine, 7th edition, p. 81. Hesser's Barner sides, Publisher

Petry N; Olofin I; Hurrell RF; Boy E; Wirth JP; Moursi M; Donahue Angel M and Rohner F (2016):

> The Proportion of Anemia Associated with Iron Deficiency in Low, Medium, and High Human Development Index Countries: A Systematic Analysis of National Surveys. Nutrients. 2; 8 (11):693.

Pratt NS; Durham HA and Sherry CL (2014):

Nutrient Intakes from Food of Lactating Women Do Not Meet Many Dietary Recommendations Important for Infant Development and Maternal Health. *Food Nutr. Sci.* 5, 1644–1651.

Rivlin RS (2010):

Riboflavin. In: Coates PM, Betz JM, Blackman MR, et al., eds. Encyclopedia of Dietary Supplements. 2nd

ed. London and New York: Informa Healthcare:691-9.

Ross A (2014):

Vitamin A. In: Ross AC, Caballero B, Cousins RJ, Tucker KL, Ziegler TR, eds. Modern Nutrition in Health and Disease. 11th ed. Baltimore, MD: Lippincott Williams & Wilkins:260-77.

Roth DE; Abrams SA; Aloia J; Bergeron G; Bourassa MW; Brown KH; Calvo MS; Cashman KD; Combs G; De-Regil LM; Jefferds ME; Jones KS; Kapner H; Martineau AR; Neufeld LM; Schleicher RL; Thacher TD and Whiting SJ (2018):

> Global prevalence and disease burden of vitamin D deficiency: a roadmap for action in low- and middle-income countries. *Ann N Y Acad Sci.*; 1430 (1): 44-79.

Said HM and Ross AC (2014):

Riboflavin. In: Ross AC, Caballero B, Cousins RJ, Tucker KL, Ziegler TR, eds. Modern Nutrition in Health and Disease. 11th ed. Baltimore, MD: Lippincott Williams & Wilkins; 325-30.

Stevens G; Bennett JE; Hennocq Q; Lu Y; De-Regil LM; Rogers L; Danaei G; Li G; White R; Flaxman SR; Oehrle S; Finucane M; Guerrero R; Bhutta ZA; Then-Paulino A; Fawzi W; Black RE and Ezzati M (2015):

> Trends and mortality effects of vitamin Α deficiency in children in low-income and 138 middle-income countries between 1991 and 2013: A pooled analysis of population-based surveys. Lancet Glob. Health. 3. e528-e536.

Stone MS; Martyn L and Weaver CM (2016):

Potassium Intake, Bioavailability, Hypertension, and Glucose Control. *Nutrients*, 8, 444.

Timoneda J; Rodríguez-Fernández L; Zaragozá R; Marín MP; Cabezuelo MT; Torres L; Viña JR and Barber T (2018):

Vitamin A Deficiency and the Lung. *Nutrients*. 21; 10 (9):1132.

van Hagen P; de Jonge R; van Berge Henegouwen MI; Hötte GJ; van der Stok EP; Lindemans J; van Lanschot JJB and Wijnhoven BPL (2017):

> Vitamin B12 deficiency after esophagectomy with gastric tube reconstruction for esophageal cancer. *Dis Esophagus*. 1;30(12):1-8.

Vasara E; Marakis G; Breda J; Skepastianos P; Hassapidou M; Kafatos A; Rodopaios N; Koulouri AA and Cappuccio FP (2017):

> Sodium and Potassium Intake in Healthy Adults in Thessaloniki Greater Metropolitan Area-The Salt Intake in Northern Greece (SING) Study. *Nutrients*, 9, 417.

Waterlow JC (2006):

Measurement of wholebody protein turnover by the end-product method. In: Waterlow, J. C., editor. Protein turnover Wallingford (UK): CABI; p. 97–105.

World Health Organization, WHO (1991):

National strategies for overcoming micronutrient malnutrition. Geneva:

WHO, (2006):

Mean heights are (even) used as an indicator of nutritional status for today's populations. Medicine, 7th edition, p. 81.

World Health Organization (2012):

EffectofIncreasedPotassiumIntakeonBloodPressure,RenalFunction,BloodLipids,andOtherPotentialAdverseEffects;WorldHealthOrgani-zation:Geneva,Switzer-land,2012;ISBN978-92-4-150488-1.

World Health Organization (2012):

Guideline:PotassiumIntakeforAdultsandChildren;WHOGuidelinesInesApprovedbytheGuidelinesReviewCommittee;WorldHealthOrganization:Geneva,Switzerland;ISBN 978-92-4-150482-9.

World Health Organization WHO (2013):

Geneva, Switzerland; ISBN 978-92-4-150623-6.

World Health Organization (2014):

EuropeanFoodandNutritionActionPlan2015–2020;WorldHealthOrganization:Geneva,Switzerland, 2014.

WHO and UNICEF (2017):

ReportoftheFourthMeetingoftheWHO andUNICEFTechnicalExpertAdvisoryGrouponnutritionMonitoring(TEAM),Geneva,Switzerland.

World Health Organization (2020):

Accelerating Salt Reduction in Europe: A Country Support Package to Reduce Population Salt Intake in the WHO European Region 2020: World Health Organization: Geneva. Switzerland, 2020.

World Health Organization, 2020:

Centers for Disease Control and Prevention, WHO, Nutritional International, UNICEF. Micronutrient survey manual. Geneva:

World Health Organization WHO (2020):

Global Action Plan for the Prevention and Control of Noncommunicable Diseases 2013–2020.

Welch AA; Kelaiditi E; Jennings A; Steves CJ and Spector TD (2016):

MacGregor, A. Dietary Magnesium Is Positively Associated with Skeletal Muscle Power and Indices of Muscle Mass and May Attenuate the Association Between Circulating C-Reactive Protein and Muscle Mass in Women. J. Bone Miner. Res. 31, 317– 325.

Welch AA; Skinner J and Hickson M (2017):

Dietary Magnesium May Be Protective for Aging of Bone and Skeletal Muscle in Middle and Younger Older Age Men and Women: Cross-Sectional Findings from the UK Biobank Cohort. *Nutrients*, 9, 1189.

Wiseman EM; Bar-El Dadon S and Reifen R (2017):

The vicious cycle of vitamin a deficiency: A review. *Crit Rev Food Sci Nutr*; 57:3703-14.

Zhou SS and Zhou Y (2014):

Excess vitamin intake: an unrecognized risk factor for obesity. *World J Diabetes;* 5:1-13

Zhou SS; Li D; Zhou YM; Sun WP and Liu QG (2010):

B-Vitamin Consumption and the Prevalence of Diabetes and Obesity Among the US Adults: Population-Based Ecological Study. *BMC Public Health* 10: 746.

Zimmermann MB (2009):

Iodine deficiency. *Endocr Rev*; 30: 376–408.



Figure 1: Distribution of studied women according to demographic areas



Figure 2: Distribution of women according to age





Figure 3: The Percentage Distribution of women's social data according to the level of education



Salwa M. Saleh; Hanaa H El-Sayed; Hoda M El Gezery and Asmaa Mohamed

Figure 4: The Percentage Distribution of women's social data according to the level of Occupation.

Table (1): Distribution of w	omen's BMI related	to geographic d	listribution (n=
4694)			

Sample	Und ig	erwe ht	Nor	mal	Overweight		verweight Obese		X ²	P value
	No	%	No	%	No	%	No	%		vanae
Urban (2282)	14	0.6	438	19.2	671	29.4	1159	50.8	0.215	0.016
Rural (2412)	22	0.9	552	22.9	695	28.8	1143	47.4		
Total (4694)	36	0.8	990	21.2	1366	29.1%	2302	49.0		



Fig (5): household intake of consumed food items per day in gram

Table (2) Mineral consumption in the household/capita according toRDI of FDA in the studied sample

Minerals	Urban	Lower	Upper Equat	Frontier	P
	mean	mean	mean	mean	value
Sodium mg	7240.0	7682.0	7068.0	7356.0	0.000
D.V.		230	0 mg		
% Of D.V.	315	334	307	320	
Potassium	1992.0	2156.0	2119.0	1942.0	0.001
D.V.		470	0 mg		
% Of D.V.	42	46	45	41	
Calcium	427.0	469.0	445.0	439.0	0.059
D.V.		130	0 mg		
% Of D.V.	33	36	34.2	33.8	
Phosphorus	965.0	1008.0	957.0	1011.0	0.063
RDV		125	0 mg		
% Of D.V.	77	81	76.6	81	
Magnesium	163.0	182.0	167.5	165.0	0.064
D.V.					
% Of D.V.	39	43	40	39	
Iron	13.9	14.5	14.6	15.2	0.048
D.V.					
% Of D.V.	77	81	81	84	
Zinc	10.1	10.1	11.0	11.5	0.057
D.V.					
% Of D.V.	92	92	100	105	
Copper	1.19	1.38	1.18	1.07	0.005
D.V.					
% Of D.V.	132	153	131	119	

Reference of Recommended Daly value (D.V.) of (FDA, 2020)

Table 3: Vitamins consumption in households/capita according to RI	Э
of FDA in the studied sample	

Vitamins	Urban	lower Egynt	upper Egynt	Frontier	<i>P</i> value
	Mean	Mean	Mean	Mean	value
Vit. (A)	784.0	579.8	456.2	1096.3	0.001
mcg					
D.V.		900	mcg		
% Of D.V.	87	64	51	122	
Vit. (C) mg	67.7	83.1	68.8	55.2	0.000
D.V.		9()mg		
% Of D.V.	75	92	76	61	
Vit.(B1) mg	1.24	1.30	1.41	1.25	0.000
D.V.					
% Of D.V.	95	100	108	96	
Vit.B2 mg	0.98	0.98	0.95	1.02	0.594
D.V.	1.2 mg				
% Of D.V.	82	82	79	85	

Reference of Recommended Daly value (D.V.) of (FDA, 2020)

الملخص العربى

تواجه النساء في الدول منخفضة ومتوسطة الدخل قضابا صحبة عامة ر تبسبة تتعلق بنقص المغذبات الدقيقة. من كل منطقة في مصر، تم اختيار عينات من النساء في سن الإنجاب من غير الحوامل أو المرضعات بشكل عشوائي في عدة محافظات. شاركت 4694 امر أة تتراوح أعمار هن بين 18 و 50 عامًا في هذه الدراسة بين عامى 2015 و 2020. تم تقييم الحالة التغذوية للمغذيات الدقيقة باستخدام بيانات من العلاقات الاجتماعية للمشاركين ، والأطعمة اليومية المتناولة ، ومقاييس الجسم لهن. أظهرت النتائج أن معظم المشاركين في العينة تتراوح أعمارهم بين 25 و 35 عامًا. في مصر العليا ، شكلت النساء ما يقرب من 75.1٪ من خريجي الجامعات. سجلت المناطق الحدودية أعلى نسبة للأمية ، في حين سجلت مصر السفلي أدنى نسبة. تم العثور على أعلى نسب المهنيين العاملين في المناطق الحضرية والوجه القبلي. كانت بطالة الإناث هي الأعلى في المناطق الحدودية. وجد ايضا أن المصريات يعانين من زيادة الوزن(29.1%) والسمنة (49.0%). تستهلك الإناث معادن أكثر مما تحتاج إليه ، مثل الصوديوم والنحاس. بينما انخفض تناول البوتاسيوم والكالسيوم والمغنيسيوم إلى أقل من 50٪ من الكميات اليومية الموصبي بها. علاوة على ذلك ، انخفض استهلاك الفوسفور 81-77٪ والحديد 77-84٪ أما الزنك 92-105٪ ضمن النطاقات المسموح بها. وجد أن مناطق مصر العليا لديها أقل نسبة من فيتامين (أ) (51٪) في حين سجلت المنطقة الحدودية أعلى نسبة (122٪). كانت جميع الفيتامينات القابلة للذوبان في الماء ، باستثناء فيتامين ب 1 ، أعلى من 75٪ من المدخول اليومي الموصى به. اظهرت النتائج وجود قصور في تناول المغذيات الدقيقة لدى السيدات المصريات وحاجتهن إلى مزيد من التثقيف التغذوي.

الكلمات المفتاحية: المحافظات المصرية - النساء - المغذيات الدقيقة