

**Original Article** 

Title:

# Effectiveness of Specific Stabilization Exercise on Strength, Balance for Beginner Weightlifters

#### Author : Mohamed Saied Abdallah Eldeeb.

Waleed Mohamed Abo Bakr Mady

Affiliations:

E-mail address :

Doi: 10.21608/JAT.2023.281187

#### Abstract

The purpose of this study is to assess the effectiveness of using Specific Stabilizing Exercises on Strength and balance for beginner weightlifter. Balance is generally defined as the ability to maintain the body's centre of gravity within its base of support and may be categorized by either static or dynamic balance. Six novice weightlifters participated in the study. Mediolateral Maximum Strength and Explosive Power and Balance testing were performed a week before (Pre) and after (Intermediate) a 4-week and after (Post) a 8-week Specific Stabilizing Exercises program. Analysis of the results showed using Specific Stabilizing Exercises improvement dynamic balance and digital level for beginner weightlifter. where the value of (f) between (19.30 to 117.05) in dynamic balance test. The results also showed statistically significant differences in favor of the maximum strength test where the value of (54.36), while the explosive power did not show any statistically significant differences. All of these improvements lead to an improved digital level for beginners' weightlifters Where the value of (f) the two Lifts snatch and clean and jerk. (18.17) - (22.09). Such an exercise program can be incorporated into a training regime of Beginner weightlifters to improve core stability as well as dynamic balance and then their digital level.

#### **Keywords**:

: Specific Stabilizing Exercises, Dynamic balance, Core stability.

#### Introduction

Weightlifting is considered one of the Olympic sports that contribute to achieving an overall balance of the individual practitioner. In addition to the improvement in the level of multiple muscle strength types. the Olympic lifts are very complex and contain a high degree of technical difficulty .These exercises require also greater core stability and overall dynamic body balance



Many elite athletes continue to undertake core stability and core strength training as part of their training programme.(36),(11)

For weightlifters, the enhancement of strength and balance plays a critical role in their performance, since they must control extremely heavy weights above their head height for a couple of seconds, while their arms are fully extended. Any kind of muscle imbalance and inappropriate technique may hamper athletes' performance and lead to dangerous situation. (32

Olympic-style weightlifting are slightly different from other sports because weightlifters often consciously focus on bracing or stabilizing prior to initiating movement. Whereas in other sports, like tennis, basketball, or marathon running, stabilization is a complex process running in the background as the athlete focuses on external tasks. In each of these cases, the brain must continuously work to regulate intra- abdominal pressure (IAP) to preserve spinal stability for movement and function.(13)

Core muscles consist of the rectus abdominis, transverse abdominis, multifidus (back muscles), internal and external obliques, quadratus lumborum (low-back muscle), and spinal erectors (back muscles). To a greater extent, the glutes, hamstrings, and hip rotators may also be included because of their relationship to the hip joint.(27),(30)

The primary function of the core is for stabilization. Stability is the ability to control force or movement. Core stabilization is important because it provides a powerful link between lower and upper body strength. Generally, core development will consist of flexion-extension type exercises (sit-ups) that target the rectus abdominis. However, although the core is being strengthened, these exercises do not address the need for a stable spine or the transfer of power from the lower to upper body. Also, because of the body's position (lying) in most of these exercises, there is a low carryover to actual sport. The best core work is done in a sport-specific stance (standing), while maintaining the spine in an upright and erect position and allowing movement from the extremities in practical ways that place stress on the core (squat).(19),(20),(28)

Posture will be much more challenged during two lifts positions (clean- snatch). The weight or resistance is placed above the head at arm's length, causing a shift in the body's center of gravity. Remember, the center of gravity or point of stability resides in the core. If the core is off balance, the body is off balance. Because of this increased distance, the center of gravity shifts higher, thus requiring the muscles of the core to work harder in order to stabilize and support the spine in an upright position. Overhead squatting also puts the trunk in an elongated position, which in turn causes a natural activation of the core muscles. With the arms fully extended overhead, the deep abdominal muscles (transverse abdominals, internal/external obliques) along with the spinal erectors, must contract in order for both stabilization and thoracic extension to be maintained.(17)

Dynamic balance is one of the most important motor skills and is defined as the ability to perform a task while maintaining or regaining a stable position, or the ability to maintain or regain balance on an unstable surface with minimal extraneous motion. Improving the dynamic balance



can improve the overall physical performance in most sports, such as judo, gymnastics, weightlifting. (27),(26),(16)

In addition, exercise performed on unstable sur- faces rather than under stable conditions seems to more effectively increase muscle activation patterns in the trunk region and limbs, while decreasing perceived exertion. The overall effects of this approach, which is termed "core stability and strength enhancement", may possibly improve athletic performance and reduce the incidence of injury. As a result, it has a growing popularity not only among elite athletes but also the general population. (2),(7),(25),(24)

There is a lack of research looking at the effect of core stability on athletic performance. Although some studies have implied that there is an advantageous effect on performance by improving core stability and strength, these conclusions are largely assumptions based on basic testing. reported that core stability and balance are critical for good performance in almost all sports and activities. This is due to the 3-dimensional nature of many sporting movements, which demands that athletes must have good strength in the hip and trunk muscles to provide effective core stability. Some sports require good balance, some force production, and others body symmetry, but all require good core stability in all three planes of motion lack of core strength and stability is thought to result in an inefficient technique, which predisposes the athlete to injury.

The most predominant literature regarding balance has emphasized the physio- logical mechanisms controlling stability. Topics range from extrinsic factors (environment) to intrinsic factors (muscle coordination, vestibular response). Balance is achieved through an interaction of central anticipatory and reflexive actions as well as the active and passive restraints imposed by the muscular system. However, less research has attempted to document the effects of balance on performance measures (force, power). Furthermore, short- and long-term adaptations to unstable environments need more substantial research. While force and other performance measures can be adversely affected by a lack of balance, the transferability of instability training to activities of sport is not precisely known. The applicability of instability and resistance training using unstable platforms or implements may have strong relevance in athletic setting. Therefore, a comprehensive review of the literature in this area may possibly be of benefit to practitioners who deal with the athletes or persons debilitated by balance and/or stability disabilities. (3)

Over the past two decades, coaching professionals and athletes have shown increasing interest in training routines to enhance the physical prerequisites for strength performance in this regard. Exercise scientists have identified instability resistance training as a possible means to improve strength performance under conditions of instability with a special emphasis on the core muscles.(5)

# **Research objective:**

• Identify effective of using Specific Stabilizing Exercises on strength, dynamic balance for beginner weightlifter.

# **Research hypothesis:**

- Are there any statistically significant differences between pre and post measurement to experimental group, which implements the program of Specific Stabilizing Exercises on Maximum strength and Explosive power.
- Are there any statistically significant differences between pre and post measurement to experimental group, which implements the program of Specific Stabilizing Exercises on dynamic balance.
- Are there any statistically significant differences between pre and post measurement to experimental group, which implements the program of Specific Stabilizing Exercises on Digital level.

# **Material and Methods**

#### **Research procedures:**

The researchers used the experimental approach using the experimental design of a single experimental set in a method (Intermediate measurements).

### **Research domains:**

#### Spatial domain:

All pre, in-between and post measurements, and Implementing training program were conducted at in the weightlifting hall, the gymnasium, of physical Education at the University of Alexandria

#### Time domain:

- Premeasurements conducted in the period from April 10th to April 15th, 2021
- Program implementation conducted in the period from April 17th to June 15th, 2021.
- Post measurements conducted in the period from June 16th to June 20th, 2021.

#### **Research sample:**

- The sample was randomly selected from beginner weightlifters, with a total 6 players.

#### **Measuring devices:**

•	<b>Basic measurements:</b>	-	Age (Age) / total length of the body / the body weight
•	Physical measurements:	-	Sargent Test (cm) / Front squat test (kg)
•	Special measurements:	-	Y Balance Test (cm) / ON leg Squat (Rep)
•	<b>Record Level :</b>	-	Record Level to snatch / Record level to clean and jerk.

#### **Program design**

Training program, which consisted of 8 weeks with 4 training units per week, program with a combination of specific stability exercises, strength exercise, weightlifting exercises. The balance training interventions consisted of balance exercises on both a stable and unstable surface, with or without recurrent destabilization during performance. such as Overhead Pitcher's Squat on Ball - Overhead Lunge.



#### **Equipment and exercises:**

Training program included exercises on stable and unstable surfaces in anterior/posterior and mediolateral directions, with or without recurrent destabilization (strengthening exercises, the balance training programs included transitions from a double-leg stance to a single-leg stance on a stable or unstable surface For these types of exercises. wobble boards that allow for multiplanar movement, BOSUs inflated rubber discs and Swiss balls. These activities were also combined with resistance exercises while balancing with Hanging Band. Core stability training was also used to improve balance. Core stability exercises were also performed using Swiss balls.

#### **Program Components :**

1	Stability Snatch Exercises	4	Stability Clean Exercises
1.	Snatch Start position On unstable surface	1.	Clean Start position on unstable surface
2.	Single-Arm Dumbbell Snatch	3.	Single-Arm Dumbbell clean
3.	Single Leg Snatch	4.	Single Leg clean
4.	Dumbbell Hang Power Snatch	5.	Dumbbell Hang Power clean
5.	Single-leg dumb-bell snatch	6.	Single-leg dumb-bell clean
6.	Kettlebell One Arm Split Snatch	7.	Kettlebell One Arm Split clean
2	Stability Overhead Squat Exercises	5	Stability Squats Exercises
1.	Hanging band Overhead Squats	1.	Single Leg Squat
2.	Single-Arm Dumbbell Overhead Press	2.	F.S different weight on both sides
3.	Overhead Bulgarian split squat on ball	3.	Parallel squat on unstable surface
		4.	Hanging Band Squats
3	Weight Stability Exercises	6	Stability Core Exercises
1.	Single Leg Bent Over Barbell Row	1.	Russian twist on ball Twist on ball
2.	Single Leg RDL	2.	Stability ball hamstring curl
3.	Snatch Grip Romanian Deadlift one leg	3.	Single-leg stability ball hip thrust
4.	DB Lunge jump	4.	Stability ball V-pass
5.	Hanging Band RDL	5.	Stability ball deadbug
6.	Single Leg deadlift	6.	Skier crunch on ball
7.	Hanging Band Bench Press	7.	Bridge with heel raise on roller
8.	DB single arm chest press on ball	8.	Stability Ball Knee Tucks

# Table 2. Procedure of the 10-week specific stabilization exercise program

#### **Procedural research steps :**

- 1. Designing and codifying appropriate tests for measuring and analyzing research variables.
- 2. Designing a training program on a scientific basis for the experimental research group that includes specific stabilization exercises.
- 3. Conducting Pre-measurements for the research sample
- 4. Implementation of the trial program
- 5. Conducting the Intermediate measurements during the experiment.



- 6. Conducting Post measurements of the research variables.
- 7. Processing data statistically and describing it in tables for discussing the results.

# **Statistical methods:**

All statistical work has been done using SPSS program to find out the following:

- Minimum
- Maximum
- arithmetic mean.
- standard deviation.
- Skewness, variance, and kurtosis coefficient.
- Improvement Percentile
- Repeated Measures ANOVA
- Eta square coefficient
- Lowest significant difference L.S.D

# Results

Table (1)Descriptive Statistics

N=6					-					
Statistical indications Variables			Unit	Minimum	Maximum	Mean	Std.	Skewness	Kurtosis	
bles	Age			year	20.00	24.00	22.67	1.37	-1.93	4.55
varia	Height		cm	168.00	180.00	175.50	4.09	-1.38	2.82	
Basic	Body Weight		kg	62.00	91.00	75.50	9.89	0.45	0.54	
	Maximum Strength	Fro	ont Squat	kg	90.00	120.00	96.67	11.69	2.22	5.08
	Explosive Power	Sar	gent Test	cm	260.00	293.50	272.75	11.54	1.25	2.26
		Post	Left erolateral	cm	68.00	80.00	72.17	4.92	0.91	-0.64
e		] Poste	Right erolateral	cm	61.00	86.00	66.83	9.79	2.06	4.32
ıriabl	Y Balance	Post	Left eromedial	cm	55.00	89.00	74.83	11.82	-0.83	0.88
cal va	Test	] Poste	Right eromedial	cm	67.00	90.00	76.92	8.36	0.51	-0.27
hysic		Left Anterior		cm	40.00	56.00	47.50	5.39	0.30	0.81
4		Righ	t Anterior	cm	46.00	59.00	51.50	4.23	1.00	2.67
	Dynamic	Squat	Left	rep	3.00	13.00	7.33	3.93	0.37	-1.58
	Balance	ON leg	Right	rep	3.00	10.00	5.33	2.50	1.61	2.85



Statistical indicat	ions Variables	Unit	Minimum	Maximum	Mean	Std.	Skewness	Kurtosis
record level	Clean and Jerk	kg	55.00	100.00	67.50	16.66	2.02	4.20
	Snatch	kg	40.00	80.00	52.83	14.70	1.49	2.72

Table (1) results reveal of the statistical significance of the research sample in the basic variables, physical tests, and digital level under study before the experiment that the data for the total research sample are moderate, not scattered and characterized by the normal distribution of the sample, where the values of the torsion coefficient range between (-1.93 to 2.22). These values are close to zero, which confirms the moderation of the data for the research sample before the experiment.

 Table (2)

 Statistical indications of physical variables and the dynamic balance and digital Level of the experimental group before, during and after the experiment

<u>N=6</u>										
Statistical i	Statistical indications			Tr measu	ibal irement	Inter measure		Distance measurement		F
		Variables		Mean	Std.	Mean	Std.	Mean	Std.	•
Maximum Strength	From	nt Squat	kg	96.67	11.69	109.17	15.63	118.33	13.66	*54.36
Explosive Power	Sargent Test		cm	272.75	11.54	275.83	10.83	277.08	7.61	3.50
	Left Pos	sterolateral	cm	72.17	4.92	85.33	8.59	90.33	5.99	*25.02
	Right Posterolateral		cm	66.83	9.79	92.00	7.67	97.17	6.18	*117.05
Y Balance	Left Posteromedial		cm	74.83	11.82	90.00	6.03	94.33	7.09	*19.30
Test	Right Posteromedial Left Anterior		cm	76.92	8.36	90.50	7.66	95.00	7.07	*43.13
			cm	47.50	5.39	53.17	2.79	56.83	4.12	*31.09
	Right	Anterior	cm	51.50	4.23	55.17	3.87	58.00	3.16	*22.32
Dynamic	ON leg	Left	rep	7.33	3.93	9.50	5.32	18.00	6.81	*17.68
Balance	Squat	Right	rep	5.33	2.50	10.50	2.74	16.83	8.04	*12.77
Digital	Clean	and Jerk	kg	67.50	16.66	75.00	18.44	80.33	20.85	*18.17
level	Si	natch	kg	52.83	14.70	56.92	18.34	63.67	18.83	*22.09

Significant at the level (0.05) the tabular value of "F" (5.05) \*

Table (2) results reveal of the statistical significance of the physical variables (maximum strength - explosive power) that there are statistically significant differences at the 0.05 level) between the pre-measurement, inter-measurement, and post-measurement in the variable of



maximum strength, while there are no Statistically significant differences in the explosive power variable.

Table (2) results reveal of the statistical significance of the variables related to the dynamic balance (Y Balance Test- ON leg Squat) that there are statistically significant differences at the level of 0.05) between the pre-measurement, the inter-measurement, and the post-measurement, where the ranged The calculated (F) values are between (12.77: 117.05), and these values are greater than the tabular "F" value at the (0.05) level.

Table (2) results reveal of the statistical significance of the variables related to the numerical level (Clean and Jerk - Snatch) that there are statistically significant differences at the 0.05 level) between the pre-measurement, the inter-measurement, and the post-measurement, where the values of (F) calculated between (18.17: 22.09), and these values are greater than the tabular value of " F " at the level (0.05).



Figure (1) The arithmetic mean of the (maximum Strength – explosive power) for the experimental group before, during and after the experiment









Figure (3) Arithmetic mean of the dynamic balance test (On leg squat) of the experimental group before, during and after the experiment



Figure (4) The arithmetic means of the Digital level (Clean and jerk-Snatch) of the experimental group before, during and after the experiment

#### Table (3)

Contrast analysis of repeated measurements (**tribal- inter-dimensional**) of physical variables, dynamic balance and Digital level of the experimental group

Statist	ical indic	cations Variables	Type III Sum of SquaresdfMean Square		F	Sig.	Partial Eta	
u c		Intercept	210168.06	1	210168.06	*200 20	0.00	0.00
nu ngtl	Front	Error	2706.94	5	541.39	*388.20		0.99
axin Irer	Squat	Sphericity Assumed	1419.44	2	709.72	*54 36	0.00	0.92
St		Sphericity Assumed Error	130.56	10	13.06	*54.30		
e		Intercept	1363450.89	1	1363450.89		0.00	1.00
osiv ver	Sargent Test	Error	1456.61	5	291.32	*4080.22	0.00	1.00
pla Pow		Sphericity Assumed	59.69	2	29.85	3 50	0.07	0.41
Ex		Sphericity Assumed Error	icity Assumed Error 85.31 10 8.53		5.50	0.07	0.41	
t		Intercept	122842.72	1	122842.72	*1240.26	0.00	1 00
les	Left	Error	458.28	5	91.66	*1340.20	0.00	1.00
Ce	eral	Sphericity Assumed	1056.78	2	528.39	*25.02	0.00	0.83
Y Balano		Sphericity Assumed Error	211.22	10	21.12	*25.02	0.00	
		Intercept	131072.00	1	131072.00	*700.97	0.00	0.00
		Error	828.67	5	165.73	*/90.80	0.00	0.99



#### https://jat.journals.ekb.eg/ Faculty of Physical Education, University of Sadat City Journal of Theories and Applications of physical education sport sciences

Statist	ical indic	cations	Type III Sum					P
		Variables	of Squares	df	Mean Square	F	Sig.	artia Eta
	Right	Sphericity Assumed	3160.33	2	1580.17			
	Posterolat eral	Sphericity Assumed Error	135.00	10	13.50	*117.05	0.00	0.96
		Intercept	134334.72	1	134334.72			
	Left	Error	806.28	5	161.26	*833.05	0.00	0.99
	Postero	Sphericity Assumed	1258.11	2	629.06	*10.20	0.00	
	inculai	Sphericity Assumed Error	325.89	10 32.59		*19.30	0.00	0.79
		Intercept	137725.01	1	137725.01	*00 <b>5</b> 01	0.00	0.00
	Right	Error	769.40	5	153.88	*895.01	0.00	0.99
	Postero medial	Sphericity Assumed	1063.53	2	531.76	*12 12	0.00	0.00
	meului	Sphericity Assumed Error	123.31	10	12.33	*43.15		0.90
		Intercept	49612.50	1	49612.50	*1005 20	0.00	1.00
	Left	Error	226.50	5	45.30	*1095.20		
	Anterior	Sphericity Assumed	265.33	2	132.67	*21.00	0.00	0.86
		Sphericity Assumed Error	42.67	10	4.27	-31.09	0.00	0.00
	Right Anterior	Intercept	54230.22	1	54230.22	*1450 55	0.00	1.00
		Error	185.78	5	37.16	1459.55	0.00	1.00
		Sphericity Assumed	127.44	2	63.72	*22.32	0.00	0.82
		Sphericity Assumed Error	28.56	10	2.86	- 22.32		0.02
		Intercept	2426.72	1	2426.72	*25 28	0.00	0.88
e	ON leg	Error	342.94	5	68.59	-35.30	0.00	
lan	(Left)	Sphericity Assumed	381.44	2	190.72	*17 68	0.00	
Ba		Sphericity Assumed Error	ity Assumed Error 107.89 10		10.79	17.00	0.00	0.78
nic		Intercept	2134.22	1	2134.22	*15 76	0.00	0.00
nar	ON leg	Error	235.78	5	47.16	45.20	0.00	0.90
Dy	(Right)	Sphericity Assumed	398.11	2	199.06	*17 77	0.00	0.72
	Ϋ́Ο΄	Sphericity Assumed Error	155.89	10	15.59	12.77	0.00	0.72
		Intercept	99309.39	1	99309.39	*06 01	0.00	0.05
	Clean	Error	5123.61	5	1024.72	- 90.91	0.00	0.95
/el	and Jerk	Sphericity Assumed	498.78	2	249.39	*18 17	0.00	0.78
lev		Sphericity Assumed Error	137.22	10	13.72	10,17	0.00	0.70
ital		Intercept	60146.68	1	60146.68	*67 50	0.00	0.02
Dig	Spotsk	Error	4455.07	5	891.01	.0/.20	0.00	0.93
D	Snatch	Sphericity Assumed	359.19	2	179.60	*22.00	0.00	0.02
		Sphericity Assumed Error	81.31	10	8.13	** 22.09	0.00	0.82

Table (4) f-value at the level of 0.05 (10.97) and within the measurements (5.05)



#### Table (5)

# Lower moral difference (LSD) value at 0.05 between average physical variables, dynamic balance and the Digital level of the experimental group before, during and after the experiment

Statistical indication					Ν	Iean Diffe	rence (I-J)	
	Variables		Measurements	Mean	Intermeas	sure	Distance me	easurement
вц			Tribal measurement	96.67	↑*12.500	0.004	*21.667↑	0.000
ximur rength	Front Squat	ka	Intermeasure	109.17			*9.167↑	0.002
Aaxi Stre		кg	Distance measurement	118.33				
e			Tribal measurement	272.75	3.083	0.081	4.333	0.079
olosiv	Sargent Test	cm	Intermeasure	275.83			1.250	0.478
ixpl			Distance measurement	277.08				
			Tribal measurement	72.17	↑*13.167	0.008	*18.167↑	0.000
	Left Posterolateral	cm	Intermeasure	85.33			5.000	0.111
	1 osteroitterui		Distance measurement	90.33				
			Tribal measurement	66.83	↑*21.167	0.000	↑*30.333	0.000
	Right Posterolateral	cm	Intermeasure	92.00			<b>↑*5.167</b>	0.002
	1 osteroitterui		Distance measurement	97.17				
	Left Posteromedial	cm	Tribal measurement	74.83	↑*15.167	0.012	<b>↑</b> *19.500	0.005
ce Test			Intermeasure	90.00			↑*4.333	0.006
			Distance measurement	94.33				
alan	Right Posteromedial	cm	Tribal measurement	76.92	↑*13.583	0.003	<b>↑*18.083</b>	0.000
Y Ba			Intermeasure	90.50			<b>↑*4.500</b>	0.023
			Distance measurement	95.00				
	Left Anterior	cm	Tribal measurement	47.50	↑*5.667	0.016	↑*9.333	0.000
			Intermeasure	53.17			↑*3.667	0.003
			Distance measurement	56.83				
		cm	Tribal measurement	51.50	↑*3.667	0.018	↑*6.500	0.003
	Right Anterior		Intermeasure	55.17			↑*2.833	0.002
			Distance measurement	58.00				
			Tribal measurement	7.33	2.167	0.189	↑*10.667	0.008
ance	ON leg Squat (Left)	rep	Intermeasure	9.50			↑*8.500	0.003
Bal	()		Distance measurement	18.00				
mic			Tribal measurement	5.33	↑*5.167	0.003	↑*11.500	0.006
Jyna	ON leg Squat (Right)	rep	Intermeasure	10.50			6.333	0.083
	(8)		Distance measurement	16.83				
ıt			Tribal measurement	67.50	↑*7.500	0.017	↑*12.833	0.006
smer	Clean and Jerk	kg	Intermeasure	75.00			↑*5.333	0.004
neve			Distance measurement	80.33				
l act			Tribal measurement	52.83	4.083	0.063	↑*10.833	0.003
gita	Snatch	kg	Intermeasure	56.92			↑*6.750	0.002
Di			Distance measurement	63.67				

Significant at a level of significance less than  $0.05^*$ 

Table (6)



#### https://jat.journals.ekb.eg/ Faculty of Physical Education, University of Sadat City Journal of Theories and Applications of physical education sport sciences

# Improvement rates between average physical variables and the digital achievement of the experimental group before, during and after the trial

Statistical					Improvement rates		
indication	8 Variables	Unit	Measurements	mean	Intermeasure	Distance measurement	
	v al lables		Tribal measurement	96.67	%12.93	%22.40	
muı ngtl	Front Squat	kg	Intermeasure	109.17	7012.23	%8.39	
laxi tre	110nt Squat		Distance measurement	118 33		700.37	
e 3			Tribal measurement	272.75	%1.13	%1.59	
osiv ver	Sargent Test	cm	Intermeasure	275.83	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	%0.45	
∃xple Pov	~ 8	CIII	Distance measurement	277.08		700.15	
[+1			Tribal measurement	72.17	%18.23	%25.16	
	Left	cm	Intermeasure	85.33		%8.57	
	Posterolateral	•	Distance measurement	90.33			
	I		Tribal measurement	66.83	%37.66	%45.39	
	Right	cm	Intermeasure	92.00		%5.62	
	Posterolateral	•	Distance measurement	97.17			
			Tribal measurement	74.83	%20.27	%26.06	
est	Left	cm	Intermeasure	90.00		%4.81	
ce T	Posteromedial		Distance measurement	94.33			
lanc			Tribal measurement	76.92	%17.65	%23.50	
Ba	Right Posteromedial	cm	Intermeasure	90.50		%4.97	
Y	1 Ostel onieulai		Distance measurement	95.00			
		cm	Tribal measurement	47.50	%11.94	%19.64	
	Left Anterior		Intermeasure	53.17		%6.88	
			Distance measurement	56.83			
			Tribal measurement	51.50	%7.13	%12.62	
	<b>Right Anterior</b>	cm	Intermeasure	55.17		%5.13	
			Distance measurement	58.00			
e			Tribal measurement	7.33	%29.60	%145.57	
anc	ON leg Squat (Left)	number	Intermeasure	9.50		%89.47	
Bal	(100)		Distance measurement	18.00			
mic			Tribal measurement	5.33	%96.99	%215.75	
ynal	ON leg Squat (Right)	number	Intermeasure	10.50		%60.29	
Di	(8)		Distance measurement	16.83			
ent			Tribal measurement	67.50	%11.11	%19.00	
eme	Clean and Jerk	kg	Intermeasure	75.00		%7.11	
niev			Distance measurement	80.33			
acł			Tribal measurement	52.83	%7.74	%20.52	
jital	Snatch	kg	Intermeasure	56.92		%11.86	
Dig			Distance measurement	63.67			



#### Discussion

The primary objective of our study was to investigate the effects of Specific Stabilizing Exercises on Strength and balance for Beginner weightlifters preparing for Olympic lifts. The main findings are that the applied training program, which consisted of 8 weeks with 4 training units per week, significantly improved the maximum strength, dynamic balance and digital level for Beginner weightlifters.

#### Strength

The results of Strength showed a significant improvement in the maximum strength, which reached the value of F (54.36) while there were no significant differences in the explosive power test. These results are in agreement with studies Behm DG, 2002. Zemkova ,2017.

The researcher explains the improvement in maximum strength due to the effect of instability exercises on the core muscles, which have a key role in lifting maximum weights because of their importance in the stability of the spine and maintaining the position of the trunk erect. Strong correlation with the performance in weightlifters.

The results of strength in training program may be affected by some considerations, including weightlifted, subject's training back- ground, number of repetitions and sets, rest intervals, and so forth. This fact has to be taken into account when instability resistance exercises are implemented into the training program. These results are in agreement with studies Erika Zemkova, 2017.

#### Balance

The results of the dynamic balance showed a significant improvement in the balance for the quads resulting from the development of the Core muscle, and the results of the dynamic balance tests led to that. Where the F value of the (Y Balance Test- ON leg Squat) tests ranged between (12.77: 117.05). These results are in agreement with studies Imai, A., Kaneoka et al 2014. Hassan I, (2017). Watson, T,et al 2017. Rafał Szafraniec1, 2020. Szafraniec R, 2018. Stanton R, 2004.

A more pronounced activation of stabilizing muscles is assumed to be the main feature of instability resistance exercises. This assumption has been proven by EMG studies, which have highlighted significantly greater electromyographic activity of trunk-stabilizing muscles during exercises under unstable as compared to stable conditions. Intervention studies also demonstrated an enhanced improvement of trunk stability after training programs utilizing unstable devices as compared to floor exercises. Findings indicate that instability resistance training may facilitate the neural adaptation of trunk-stabilizing muscles, resulting in an improvement in trunk stability. Erika Zemkova, 2017 . Hassan I, 2017. Ozmen T, 2016. Sandrey M.A,2013.

The analyses including papers in which training protocols demonstrated positive effects on balance performance suggest that an efficient training protocol should last for 8 weeks, with a frequency of two training sessions per week, and a single training session of 45 min. This standard was established based on 36 reviewed studies. Anna Brachman et al ,2017. Bryanton ,2017.



Training under unstable and imbalanced conditions results in stability and that can be experienced during a variety of daily and sport activities. However, it is reasonable to assume that using unstable surfaces during training should be based on improvement of the neuromuscular adaptation process, particularly at the early stages of a resistance training pro- gram. In addition, it is a favorable to admit that enhancing balance ability produces greater muscular functionality. Sung Hwan et al 2013. Behm DG,2006. Behm DG,1995. Kibele A, 2009.

#### **Record level**

- The results of the digital level showed a significant improvement the variables related to the numerical level (Clean and Jerk Snatch) which reached the value of F (18.17: 22.09), and these values are greater than the tabular value of "F" at the level (0.05).
- This study focused more on muscular balance, strength, and strength development, and these three factors are leading to an improvement in the Digital level for weightlifters.
- Performance improves better if the training is specific to the type of activity practiced and includes the most important muscles working in this activity.
- The development in digital achievement occurs when completing the physical attributes of the lifter, because weightlifting is one of the sports that require high physical fitness, and lifters need a set of physical attributes, the most important of which are: maximum muscular strength, ability and balance, and there is also a positive correlation between strength and the digital level of the lifters. Vorobyev A.N ,1978.

#### **Conclusion**(s)

The applied short-term of Specific Stabilizing Exercises for weightlifting improved Strength, dynamic balance and Core muscle in Beginner weightlifters attempting to start training the Olympic lifts. Such a training program can be incorporated into a training regime of Beginner weightlifters to prepare them for technically difficult exercises using free weights.

Greater instability should challenge the neuromuscular system to a greater extent than stable conditions, possibly enhancing strength gains attributed to neural adaptations. Hence instability resistance training programs may both improve athletic performance.

# **Recommendations:**

- 1. The core muscles are one of the most important muscles that maintain the balance of the lifters, and therefore attention must be paid to their development.
- 2. Use the suggested Specific Stabilizing Exercises in the program to develop balance for Beginner weightlifters.
- 3. It is important to pay attention to the development of dynamic balance for beginner lifters because of its impact on the development of performance.
- 4. The exercises included in the lifters training programs must be varied between balance exercises strength explosive power because they are the basic elements for the development of the level.



5. Diversity in the use of tools in the exercise adds an atmosphere of joy and enjoyment of the exercise.

### **References:**

- 1. Akuthota V, Ferreiro A, Moore T, Fredericson M. Core stability exercise principles. Current sports medicine reports. 2008; 7:39–44.
- 2. Anderson K, Behm DG: Trunk muscle activity increases with unstable squat movements. Can J Appl Physiol, 2005, 30: 33–45.
- 3. Anderson, K, Behm, DG. The impact of instability resistance training on balance and stability. Sports Med. 2005;35(1):43-53.
- Anna Brachman, Anna Kamieniarz, Justyna Michalska1, Michał Pawłowski, Kajetan J. Słomka, Grzegorz Juras. Balance Training Programs in Athletes – A Systematic Review. Journal of Human Kinetics volume 58/2017, 45-64
- Armin Kibele, Lifting Loads on Unstable Platforms A Supplementary View on Stabilizer Muscles and Terminological Issues, The Open Sports Sciences Journal, 2017, 10, 114-121
- 6. **Behm DG, Anderson K.** The role of instability with resistance training. J Strength Cond Res, 2006, 20: 716–722.
- 7. **Behm DG, Leonard A,** Young W. Trunk muscle EMG activity with unstable and unilateral exercises. J Strength Cond Res, 2005, 19: 193–201.
- 8. **Behm DG, Muehlbauer T, Kibele A, Granacher U**. Effects of strength training using unstable surfaces on strength, power and balance performance across the lifespan: A systematic review and meta-analysis. Sports Med 2015; 45(12): 1645-69.
- 9. **Behm DG:** Neuromuscular implications and applications of resistance training. J Strength Cond Res, 1995.
- 10. Behm, D.G., K. Anderson, and R. S. Curnew. Muscle force and activation under stable and unstable conditions. J. Strength Cond. Res. 16(3):416–422. 2002.
- 11. **Bryanton M.A., Bilodeau M.** The Effect of Vision and Surface Compliance on Balance in Untrained and Strength Athletes. Journal of motor behavior. 2019; 51:75–82.
- 12. Erika Zemkova, Instability resistance training for health and performance .Journal of Traditional and Complementary Medicine 7 (2017) 245e250.
- 13. Hackett D, and Chow, C. The Valsalva maneuver: Its effect on intra-abdominal pressure and safety issues during resistance exercise. The Journal of Strength and Conditioning Research 27(8): 2338-2345, 2013.
- 14. **Hassan I.** The effect of core stability training on dynamic balance and smash stroke performance in badminton players. International Journal of Sports Science and Physical Education. 2017; 2:44–52.
- 15. **Hassan, I.** The effect of core stability training on dynamic balance and smash stroke performance in badminton players. International Journal of Sports Science and Physical Education 2(2017), 44-52.
- 16. **Hrysomallis C.** Balance ability and athletic performance. Sports medicine. 2011; 41:221–232.
- 17. **Ian Hasegawa, CSCS**. Using the Overhead Squat for Core Development. Volume 3 Number 6 NSCA's Performance Training Journal. <u>www.nsca-lift.org/perform</u>



https://jat.journals.ekb.eg/ Faculty of Physical Education, University of Sadat City Journal of Theories and Applications of physical education sport sciences

- 18. Imai, A., Kaneoka, K, Okubo, Y. and Shiraki, H. Effects of two types of trunk exercises on balance and athletic performance in youth soccer players. International journal of sports physical therapy 2014.
- 19. **Jin ZH, Kibler WB, Press J, Sciascia A.** The role of core stability in athletic function. Journal of Beijing Sport University. 2008;12:039.
- 20. **Kellie C, Barton E.** Core Stability Training for Injury Prevention. Sports Health. 2013 Nov; 5(6): 514–522.
- 21. **Kibele A, Behm DG.** Seven weeks of instability and traditional resistance training effects on strength, balance and functional performance. J Strength Cond Res 2009; 23(9): 2443-50.
- 22. **Kibler WB, Press J, Sciascia A**. The role of core stability in athletic function. Sports Med 2006; 36 (3): 189-98.
- 23. Lawrence MA, Carlson LA. Effects of an unstable load on force and muscle activation during a parallel back squat. J Strength Cond Res 2015; 29(10): 2949-53.
- 24. **Marshall P, Murphy B:** Changes in muscle activity and perceived exertion during exercises performed on a swiss ball. Appl 383. Physiol Nutr Metab, 2006, 31: 376–383.
- 25. **Marshall PW, Murphy BA**: Increased deltoid and abdominal muscle activity during Swiss ball bench press. J Strength Cond Res, 2006, 20: 745–750.
- 26. **Maszczyk A, Dobrakowski P, Żak M.** Differences in motivation during the bench press movement with progressive loads using EEG analysis. Biol Sport. 2019;36(4):351– 356.
- 27. **Maszczyk A, Wilk M, Krzysztofik M.** The effects of resistance training experience on movement characteristics in the bench press exercise. Biol Sport. 2020;37(1):79–83.
- 28. **Ozmen T., Aydogmus M.** Effect of core strength training on dynamic balance and agility in adolescent badminton players. Journal of bodywork and movement therapies. 2016; 20:565–570.
- 29. **Rafał Szafraniec1, Janusz Bartkowski1, Adam Kawczyński**. Effects of Short-Term Core Stability Training on Dynamic Balance and Trunk Muscle Endurance in Novice Olympic Weightlifters, Journal of Human Kinetics volume 74/2020, 43-50.
- 30. **Sandrey M.A., Mitzel J.G**. Improvement in dynamic balance and core endurance after a 6-week core-stability-training program in high school track and field athletes. Journal of sport rehabilitation. 2013; 22:264–271.
- 31. **Stanton R, Reaburn PR, Humphries B.** The effect of short- term Swiss ball training on core stability and running economy. J Strength Cond Res 2004; 18 (3): 522-8
- 32. Sung Hwan Kang, Cheol Woo Kim, Young Il Kim, Kwi Baek Kim, Sung Soo Lee, and Ki ok Shin1Alterations of Muscular Strength and Left and Right Limb Balance in Weightlifters after an 8-week Balance Training Program. J. Phys. Ther. Sci. Vol. 25, No. 7, 2013.
- 33. Szafraniec R., Baranska J., Kuczynski M. Acute effects of core stability exercises on balance control. Acta Bioeng Biomech. 2018a; 20:145–151.
- 34. Vorobyev A.N .Weightlifting, I.W.F, Pub, Budapest. (1978).
- 35. Watson, T., Graning, J., McPherson, S., Carter, E., Edwards, J., Melcher, I. and Burgess, T. Dance, balance and core muscle performance measures are improved



following a 9-week core stabilization training program among competitive collegiate dancers. International journal of sports physical therapy 12, 25,2017. (10)
36. Winwood P.W., Cronin J.B., Brown S.R., Keogh J.W.. A biomechanical analysis of the strongman log lift and comparison with weightlifting's clean and jerk. International Journal of Sports Science & Coaching. 2015;10:869–886.