

Molecular investigation of calf diarrhea in relation to changes in some immunological profiles

Nahed Saleh¹, Tamer Allam¹, Mohamed Nayel², Reem Ahmed³

(1) Department of Clinical Pathology, Faculty of Veterinary Medicine, University of Sadat City, Egypt.

(2) Department of Medicine and Infectious Diseases, Faculty of Veterinary Medicine, University of Sadat City, Egypt.

(3) Department of Veterinary Authority, Faculty of Agriculture, University of Menoufia, Egypt.

*Corresponding author: biomohamed2009@yahoo.com Received: 3/2/2022 Accepted: 1/3/2022

ABSTRACT

The aim of this study was to highlight the importance of using PCR to investigate the most important and common pathogens causing calf diarrhea. In addition, to characterize the profiles of some inflammatory cytokines, acute phase proteins, oxidant-antioxidant status, and some immune parameters to clarify their clinical importance during the course of the disease.

The study was conducted on 44 calves, 31 calves were suffering from diarrhea manifestations based on clinical examination (diseased group). The rest 13 calves were apparently healthy and were served as control group. Blood samples were collected from both groups and were assayed for the selected biochemical and immunological parameters. Fecal samples collected directly from the rectum of diarrhetic calves were subjected to PCR technique for detecting 5 major bovine enteric pathogens including bovine Coronavirus, bovine Rotavirus, *Salmonella spp.*, *E. coli*, and *Cryptosporidium parvum*. The results of PCR indicated that *E. coli*, *Salmonella* and *Cryptosporidium* were the most important pathogens isolated from diarrhetic calves at percentages of 38.7 %, 25.8 % and 12.9 % respectively. On the other hand, Rota and Corona viruses were not demonstrated in diarrhetic calves. Regarding the selected biochemical and immunological parameters, the results showed that compared to control group, the diseased calves demonstrated significant increases in IL-6, ceruloplasmin, haptoglobin, malondialdehyde and nitric oxide. Significant decreases were observed in total antioxidant capacity, IL-10 and IgA, while no significant changes were reported in serum levels of IgG and IgM. In conclusion, the presented results can document that *E. coli*, *Salmonella* and *Cryptosporidium* are among the most important pathogens resulting in calf diarrhea. The results also provide a confirmation about the importance of proinflammatory cytokines in signaling in the development of calf diarrhea probably through oxidative stress which may rise their values as therapeutic targets for treating diarrhea in calf.

Keywords: PCR, Calf, Diarrhea, Immunity. Oxidative stress.

INTRODUCTION

Calf diarrhea is a multifactorial disease entity that can have serious financial and animal welfare implications in both dairy and beef sucker herds. The disease was found as the most commonly reported disease in young animal and about 75% of early calf mortality in dairy herds has been estimated to be caused

by acute diarrhea in the pre-weaning period (Azzadeh *et al.*, 2012). Until this time, calf diarrhea is still a major cause of productivity and economic loss to cattle producers causing high morbidity and mortality in the cattle industry worldwide (Uhde *et al.*, 2008; Bartels *et al.*, 2010). Economic losses can be attributed to long-term effects on health and

productivity of surviving calves in addition to treatment and labor costs such as antibiotics, milk replacer and oral rehydration.

Infectious diarrhea in calves is mostly associated with enterotoxigenic *Escherichia coli*, *Cryptosporidium parvum*, *Coronavirus*, *Rotavirus*, or some combination of these pathogens (Foster *et al.*, 2009). Among these pathogens, *Escherichia coli* was found as the important cause of diarrhea in neonate. The prevalence of each of pathogen and disease incidence can vary by geographical location of the farms, farm management practices, and herd size (Cho and Yoon, 2014).

Different contributing factors that are involved in the occurrence and allow the emergence of the disease include dam, calf immunity, environmental conditions and management practices (Cho and Yoon, 2014).

Because the progression of calf diarrhea can be rapid and since different pathogens or factors may be implicated in the development of the disease, a quick accurate diagnosis is critical not only to confirm the cause but also help producers to implement appropriate treatment protocols. Many laboratory methods for identifying enteric pathogens have been widely used including pathogen isolation and characterization along with histopathology, direct visualization by light or electron microscopy of pathogens in feces or intestinal contents, as well as the detection of antigens by ELISA (Cho and Yoon, 2014). Molecular diagnostics, most often Polymerase Chain Reaction (PCR)-based tests, are becoming more common in diagnostic laboratories for detecting enteric pathogens and present their own challenges in interpretation (Espy *et al.*, 2006). PCR methods are usually more sensitive and may detect lower levels of pathogens that may not be causing disease due to age resistance or those arise from cross-contamination from one calf to another. PCR methods also can detect the dead or stressed bacteria or even in the presence of antibody-antigen complexes (Blanchard, 2012).

This study aimed to highlight the importance of using PCR to investigate the most important and common enteric pathogens causing calf diarrhea. In addition, to characterize the profiles of some inflammatory cytokines, acute phase proteins, oxidant-antioxidant status, and some immune parameters to clarify their clinical importance during the course of

the disease.

MATERIALS AND METHODS

Animals

The present study was conducted on up to 6 months old (7 days till 6 months) 44 bovine calves in different localities in Menufiya governorate, Egypt (in the period from last December 2018 to the first of January 2020). 31 calves were suffering from calf diarrhea. The rest 13 calves were apparently healthy and were considered as control group.

Samples

Fecal samples were collected directly from the rectum of diarrheic calves in sterile plastic bags. Fecal samples were subjected to PCR technique for detecting 5 major bovine enteric pathogens including bovine *Coronavirus*, bovine *Rotavirus*, *Salmonella spp.*, *E. coli*, and *Cryptosporidium parvum*. Blood samples were collected from both control and diseased groups in plain centrifuge tubes, allowed to clot, centrifuged at 3000 rpm for 15 minutes for separation of serum. Serum samples were stored at -20°C until assayed for the rest biochemical and immunological studies.

Analytical methods

Nucleic acid extraction:

Total DNA extraction was performed using QIAamp Fast DNA Stool Mini Kit, (Thermoscientific™, Cat No. /ID: 19593, USA) and following the manufacturer's recommendations. RNA extraction was performed using QIAamp Viral RNA Mini Kit (QIAGEN™) according to the manufacturer's instructions.

Polymerase Chain Reaction (PCR)

Individual PCR was carried out for each pathogen. Detailed information of the primers used were summarized in (Table 1). DNA samples were tested in 50 µl reaction volumes in a 0.2 ml Eppendorf tube containing 25 µl PCR Master Mix (10X buffer, 10mM d NTPs mixture, Taq polymerase), 2 µl of each primer and 2 µl target DNA, then sterile deionizer water was added to complete the volume to a final of 50 µl, following thermal cycling profile. The thermocycling format was established according to the target pathogen as shown in Table 2. A total of 10 µl of each PCR product was subjected to electrophoresis using a 1.5% ethidium bromide (Sigma-Aldrich, U.S.A.) agarose gel in TBE buffer

and was visualized under a UV Transilluminator.

Biochemical and immunological studies:

Serum level of haptoglobin was determined by ELISA method using commercial kits of Cell Biolabs, Inc. (USA) according to the method of Sadrzadeh and Bozorgmehr (2004). Serum ceruloplasmin concentration was measured by ELISA technique using test kits of Beckman Coulter, Inc. (USA) following the method described by Kazmierczak (1996).

Evaluation of oxidative-antioxidant status was done by colorimetric detection of total antioxidant capacity (TAC) according to the method of **Koracevic and Koracevic (2001)**. Serum level of malondialdehyde (MDA) was determined colorimetrically as described by **Satoh (1978)**. Nitric oxide (NO) levels were monitored according to the method of **Montgomery and Dymock (1961)**. All oxidant-antioxidant parameters were evaluated using commercial kits of Biodiagnostics (Egypt).

Serum levels of the pro-inflammatory cytokine interleukin- 6 (IL-6) and the anti-inflammatory interleukin- 10 (IL-10) were detected by ELISA technique using commercial kits supplied by Shanghai Coon Koon Biotech Company (China) and following the manufacturer's instructions. Concentrations of the immunoglobulins IgA, IgG and IgM in the serum were detected by ELISA using commercial kits supplied by DIMA (Germany) and following the manufacturer's instructions.

Statistical analysis procedures:

All the values were presented as mean \pm standard deviation (SD). Mean values of diseased group and control group were compared by student's t test at 0.05 level of probability (Snedecor and Cochran 1980).

RESULTS

Detection of various etiological agents of diarrhea by PCR:

Individual PCR assays were performed on 44 fecal samples (31 diseased and 13 control) for detecting 5 major bovine enteric pathogens including bovine coronavirus, bovine Rotavirus, *Salmonella spp.*, *E. coli*, and *Cryptosporidium parvum*. The prevalence rate of these enteric pathogens isolated from apparently healthy and diarrheic calves by PCR method is shown in Table 3.

The results of PCR indicated that *E. coli* were detected at the most higher prevalence rate whereas the microorganism was isolated from 12 out of 31 fecal samples at a percentage of 38.7% (Fig. 1). *Salmonella spp.* were isolated from 8 out of 31 fecal samples (Fig. 2) while *Cryptosporidium parvum* was isolated from 4 out of 31 fecal samples (Fig. 3) at percentages of 25.8% and 12.9% respectively. On the other hand, *Rota* and *Corona* viruses were not demonstrated in diarrheic samples.

Biochemical and immunological parameters:

The effect of diarrhea on acute phase response as shown in Table 4 clarified that serum levels of haptoglobin and ceruloplasmin demonstrated a significant ($p \leq 0.05$) increase in the diarrhetic calves when compared to the control. Affected calves also showed higher ($p \leq 0.05$) serum concentrations of both malondialdehyde and nitric oxide compared to the healthy animals while, serum level of total antioxidant capacity was significantly ($p \leq 0.05$) lower in the diseased group compared to the control (Table 4).

Evaluation of some immunologic parameters in the diarrhetic calves cleared that serum value of IL-6 significantly ($p \leq 0.05$) increased in affected calves compared to the control but on the other hand, significant ($p \leq 0.05$) decrease was observed in the serum level of IL-10 (Table 5). Compared to the control group, the mean serum value of IgA showed a significant ($p \leq 0.05$) decrease in the diseased calves, but no significant changes were reported in the serum concentrations of IgG and IgM.

Table 1: Primer sequence, amplicon size and references of the selected diarrhea-induced pathogens.

Target agent	Primer name and direction	Nucleotide sequence 5'-3'	Amplicon Size	Reference
<i>E.coli</i>	F5(F) F5(R)	TATTATCTTAGGTGGTATGG GGTATCCTTTAGCAGCAGTATTC	314 bp	Frank <i>et al.</i> (1998)
<i>Salmonella</i>	invA-F invA-R	TTGTTACGGCTATTTTGACCA CTGACTGCTACCTTGCTGATG	521 bp	Swamy <i>et al.</i> (1996)
<i>Cryptosporidium</i>	18S rRNA-F 18S rRNA-R	CCACATCTAAGGAAGGCAGC ATGGATGCATCAGTGTAGCG	1056 bp	Jellison <i>et al.</i> (2002)
<i>Rota virus</i>	VP4-F VP4-R	TGGCTTCGCTCATTATAGACA ATTCGGACCATTATAACC	880 bp	Park <i>et al.</i> (2006)
<i>Corona virus</i>	BCV-F BCV-R	GCCGATCAGTCCGACCAATC AGAATGTCAGCCGGGGTAT	407 bp	Tsunemitsu <i>et al.</i> (1999)

E. coli: Escherichia coli.

Table 2: PCR protocol for amplification conditions of PCR products for the detection of the selected diarrhea-induced pathogens.

Target agent	Denaturation			Annealing			Extension			Final extension		
	Temp(°C)	Time	No. of cycles	Temp(°C)	Time	No. of cycles	Temp(°C)	Time	No. of cycles	Temp(°C)	Time	No. of cycles
<i>E. coli</i>	94	30	25	50	45	25	70	90	25	70	600	1
<i>Salmonella</i>	93	60	30	42	60	30	72	120	30	72	600	1
<i>Cryptosporidium</i>	94	60	30	54	60	30	72	60	30	72	600	1
<i>Rota virus</i>	94	60	30	50	120	30	72	120	30	72	600	1
<i>Corona virus</i>	94	60	30	58	60	30	72	120	30	72	420	1

E. coli: Escherichia coli; Temp: Temperature.

Table 3: Prevalence rate of different enteric pathogens isolated from apparently healthy and diarrheic calves by PCR method.

Fecal isolate	No of +ve samples		Total	(%)
	Control	Diarrhetic		
<i>E. coli</i>	0	12	12	38.7
<i>Salmonella spp.</i>	0	8	8	25.8
<i>C. parvum</i>	0	4	4	12.9
Number of samples =44 (13 control, 31 diseased)				

E. coli: Escherichia coli; *C. parvum*: Cryptosporidium parvum

Table 4: Serum profile of acute phase proteins and Oxidant-antioxidant status of diarrhetic calves compared to the control healthy group. Values are mean \pm SD.

Parameters	Groups	
	Control (N =13)	Diseased (N =31)
Ceruloplasmin(mg/dl)	21.65 \pm 0.35	32.17 \pm 3.75*
Haptoglobin (mg/dl)	109.33 \pm 2.52	160.50 \pm 16.50*
TAC (mmol / ml)	3.0 \pm 0.62	1.70 \pm 0.10*
Malondialdehyde (nmol/ml)	2.90 \pm 0.40	4.57 \pm 0.60*
Nitric oxide (U/ml)	222.67 \pm 1.53	269.50 \pm 4.50*

TAC: Total antioxidant capacity.

Significant differences in the values between the diseased and the control groups were indicated by (*) at $P \leq 0.05$.

Table 5: Interleukins and immunoglobulin profiles of diarrhetic calves compared to the control healthy group. Values are mean \pm SD.

Parameters	Groups	
	Control (N =13)	Diseased (N =31)
IL6 (pg/ml)	74 \pm 3.40	112.27 \pm 4.25*
IL10 (pg/ml)	72.60 \pm 3.90	33.67 \pm 3.81*
IgA (mg/dl)	11.03 \pm 1.05	7.50 \pm 0.50*
IgG (mg/dl)	22 \pm 5.0	62.67 \pm 7.29
IgM (mg/dl)	30.37 \pm 1.52	37.13 \pm 4.007

IL6: Interleukin6; IL10: Interleukin10; IgA: Immunoglobulin A; IgG: Immunoglobulin G; IgM: Immunoglobulin M.

Significant differences in the values between the diseased and the control groups were indicated by (*) at $P \leq 0.05$.




Figure 1: Molecular diagnosis of *E. coli* by PCR procedure showing positive bands at 314 bp.

- Lane 1: DNA Ladder. Lane 2: control positive. Lane 3: control Negative. Lane 4-15: samples.




Figure 2: Molecular diagnosis of *Salmonella spp.* by PCR procedure based on *invA* gene sequences showing positive bands at 521 bp.

- Lane 1: DNA Ladder. Lane 2: control positive. Lane 3: control Negative. Lane 4-11: samples.




Figure 3: Molecular diagnosis of *Cryptosporidium* by PCR procedure based on 18s rRNA gene sequences showing positive bands at 1056 bp.

- Lane 1: DNA Ladder. Lane 2: control positive. Lane 3: control Negative. Lane 4-7: samples.

DISCUSSION

Newborn calves are susceptible to neonatal calf diarrhea (calf scours) especially during their first 28 days of life. Calf diarrhea results in decreases the absorption of essential nutrients from milk and leads to weight loss and dehydration. Severe cases may be fatal due to dehydration and acidosis that may result in anorexia and ataxia, however, even calves that survive will have poorer performance for the remainder of their lives when compared to healthy calves (Maes *et al.*, 2003 and Berchtold, 2009).

The etiology of calf diarrhea is complex involving both infectious and non-infectious factors (Prescott *et al.*, 2008). Multiple enteric pathogens including viruses, bacteria, and protozoa are involved in the development of the disease (Izzo *et al.*, 2011). Coronavirus, Rotavirus, bacteria (*E. coli*; *Salmonella* spp., *Clostridium perfringens* Type C) and parasites (*Cryptosporidium*, coccidia) were found to be the most common causes of neonatal calf diarrhea. Among these pathogens, *Escherichia coli* was found as the important cause of diarrhea in neonate (Cho and Yoon, 2014).

Because of the clinical and economic importance of the disease, laboratory testing is necessary for accurate diagnosis of the problem. Many laboratory methods for identifying enteric pathogens have been widely used including pathogen isolation and characterization, direct visualization by light microscopy or electron microscopy, fecal bacterial culturing, and ELISA (Cho and Yoon, 2014). More recently, detection of antigens by Polymerase Chain Reaction (PCR) has been widely accepted as alternative method for the rapid detection of various bacterial and viral pathogens in clinical specimens from diarrheic calves (Blanchard, 2012).

In the present study, fecal samples of both control and diarrhetic calves were subjected to PCR for detection of 5 major bovine enteric pathogens including bovine *Coronavirus*, bovine *Rotavirus*, *E. coli*, *Salmonella* spp., and *Cryptosporidium parvum*. The results showed that *E. coli* were isolated at the higher prevalence rate at a percentage of 38.7 % (12

out of 31 fecal samples) followed by *Salmonella* spp. and *Cryptosporidium parvum* at percentages of 25.8% and 12.9% respectively.

In agreement, the incidence of diarrheagenic *E. coli* in fecal samples of calves collected during November 2015 to March 2016 in Damietta governorate showed the highest incidence rates at percentages of 43% in calves aged 2-7 days, 33% in calves aged 8-14 days followed by 24% in calves aged 15-21 days (Rehab *et al.*, 2016). In the study of Aydin *et al.*, (2001) on diarrhetic calves, 92.07 % of cases were *E. coli* positive. In that study, *T. vitulorum*, *Eimeria* spp. *Cryptosporidium* and *Salmonella* spp. were isolated at percentages of 9.90 %, 5.94 %, 5.94 and %0.99 % respectively while 5.94 % of fecal samples revealed neither parasitic nor bacterial infection.

Additionally, Chaurasia, (2001) reported 87 % occurrence of *Escherichia coli* in the fecal samples of diarrhetic calves. These results suggest that *Escherichia coli* is the most important cause of diarrhea in neonatal calves (Cho and Yoon, 2014).

On the other hand, Seid *et al.*, (2020) found that out of 7 positive samples screened by RT-PCR for molecular characterization, 1 sample was positive for Coronavirus and 4 samples showed Rotavirus. The prevalence rates of enteric pathogen and disease incidence can vary by geographical location of the farms, farm management practices, and herd size (Cho and Yoon, 2014).

Prevention and control of calf diarrhea should be based on a good understanding of the disease pathogenesis and its impact on different body responses and functions. In this respect, the present study aimed to evaluate the effect of calf diarrhea on acute phase response, oxidant-antioxidant status and serum profile of interleukins and immunoglobulins.

Investigation of acute phase proteins in this study revealed a marked increase in serum concentrations of haptoglobin and ceruloplasmin of diarrhetic calves as compared to control group. Haptoglobin and ceruloplasmin are acute phase reaction proteins which have been reported to increase

in animals subjected to external or internal challenges such as infection, inflammation, surgical trauma, or stress (Murata *et al.*, 2004). They are considered integral to the non-specific innate immune response and can be valuable quantitative markers and useful indicators for stress and disease (Petersen *et al.*, 2004).

Previous studies revealed that haptoglobin significantly increased in infected calves with *Cryptosporidium*, as compared to control group (Al Zubaidi, 2014). Additionally, Hajimohammadi *et al.*, (2013) also found a significant increase in serum ceruloplasmin levels in calves that presented diarrhea, when compared to healthy animals. Also calves that presented diarrhea and tested positive for rotavirus in feces demonstrated higher ceruloplasmin levels in their serum compared to Rotavirus negative animals (Rocha *et al.*, 2016). Similar results of increased serum ceruloplasmin levels were reported by Silva (2011) in Holstein calves experimentally infected with *Salmonella Typhimurium* and *Salmonella Dublin*, respectively.

Inflammatory cytokines including IL-6 are known to stimulate the systemic acute phase response as stated by Moshage (1997). This hypothesis was supported by the significant elevation of serum levels of IL-6 seen in the present study.

The results of oxidant-antioxidant status revealed a significant decrease in TAC in diarrhetic group compared to control which is consistent with the previous studies (Osman *et al.*, 2020). In this work, serum concentrations of malondialdehyde and nitric oxide were significantly higher in the diarrhetic calves compared to their corresponding control values. These findings together with the reduction in TAC provide an evidence of the presence of oxidative stress in the diseased calves.

Biochemically, oxidative stress is associated with increased production of oxidizing species or a significant decrease in the effectiveness of antioxidant defenses with the severity depending on the duration and intensity of such stress.

Malondialdehyde is the major product of lipid peroxidation that reacts with biological structures such as proteins, lipids, carbohydrates, and DNA, causing tissue damage so it reflects the oxidative stress in

cells and tissues (Gurer *et al.*, 1998). Similar studies on calf diarrhea proved that serum concentration of MDA and NO increase significantly in *Cryptosporidium*-infected calves (Kojouri *et al.*, 2012 and Gultekin *et al.*, 2019).

Rachmilewitz *et al.*, (1995) suggested that in acute infectious gastroenteritis, the release of cytokines and inflammatory mediators from activated macrophages is likely responsible for production of NO by phagocytic leukocytes. Sowmyanarayanan *et al.*, (2009) added that it is well known that a large amount of NO is produced over a long period during the inflammatory process of infectious disease. Gultekin *et al.*, (2019) provided another explanation for increased plasma NO concentration in diarrhetic calves which occurs due to decreased renal perfusion related to hypovolemia with the resultant decrease in urinary nitrate excretion.

The current findings demonstrated a significant increase in serum levels of the pro-inflammatory IL-6 with a significant decrease in the concentration of the anti-inflammatory IL-10 in the diarrhetic calves compared to control. These results are consistent with Stephani *et al.*, (2016) who found greater IL-6 concentrations in the diarrhetic calves than in controls.

Further, increased IL-6 gene expression in blood mononuclear cells and in intestinal epithelial cells from scouring calves was observed in calves due to infection with bovine Rotavirus (BRV) (Aich *et al.*, 2007 and Qadis *et al.*, 2014). Pro-inflammatory cytokines were documented to be important in cell signaling that promote inflammatory process and were involved in the upregulation of the inflammatory reactions. They were found also to stimulate the liver to synthesize acute-phase proteins can be justified in three ways, namely its role in the inflammation pathway and inducing the production of acute-phase proteins (Murtaugh *et al.*, 1996).

IL-10 is well known as a cytokine with potent anti-inflammatory properties which counter-regulate production and function of pro-inflammatory cytokines and thus subsiding the inflammatory process. The significant decrease in IL-10 concentrations observed in diarrhetic calves compared to the control group suggesting an overwhelming inflammatory reaction and excessive intestinal

damage. Environmental conditions can play a role in the production of bovine IL-10.

Several in vitro studies have shown that high temperature (42°C) and temperature-humidity index (THI) value of > 72 can reduce the proliferation of IL-10-secreting cells compared to low temperature (38.5°C) and a THI value of < 72 (Elvinger *et al.*, 1991 and Lacetera *et al.*, 2005).

With respect to serum immunoglobulins evaluated in this study, the mean values of serum concentrations of IgA were markedly lower in diarrhetic group compared to healthy group. On contrary, no significant changes were demonstrated in IgG and IgM between the two groups.

Serum immunoglobulin concentration in dairy cattle is an important indication of immunity against pathogenic microorganism (Quigley, 2002 and Mendonsa, 2011). IgA is known to be the main immunoglobulin found in mucous membranes including those lining the gastrointestinal tract and is an antibody that plays a crucial role in the immune function of mucous membranes.

The obtained findings were nearly similar to those reported by Balikci and AL (2014) who observed significantly lower serum levels of IgG and IgA in calf diarrhea induced by *E. coli*, *Rotavirus* and *Coronavirus*. Youssef *et al.*, (2017) explored the modulation of the gene expression of some immune-inflammatory markers including IgG, and IgA in buffaloes demonstrating digestive disorders and they found downregulation of both IgG and IgA gene expression in blood of all diseased buffaloes compared with controls indicating a state of immunosuppression.

Mahmoud *et al.*, (2018) found a highly significant reduction in concentration of IgA in *Cryptosporidium*-diseased calves which was considered a sign of the suppressive effect of pathogens on the immune system. Further explanation was provided by Athanasiou *et al.*, (2019) who attributed the reduction of serum immunoglobulins in diarrhetic calves to the transfer of these immunoglobulins, to the intestinal lumen to counteract the infection.

Moreover, Hummelshoj *et al.*, (2006) have suggested that IL-10 could positively affect the production of IgA and IgG to support the regulation of B and T cell responses. Therefore, the significant reduction in serum levels of IL-10 observed in this study

might be a reasonable cause for the decreased IgA levels in diarrhetic calves.

In disagreement with the results of IgG and IgM presented in this work, some studies recorded that serum levels of IgG were significantly decreased in diarrhetic calves than that of healthy animals (Berge *et al.*, 2009; and Al-Alo *et al.*, 2018). Rocha *et al.*, (2016) found lower IgG concentrations in diarrhetic calves with Rotavirus-positive fecal samples while serum IgA did not show significant difference between diarrhetic and healthy groups.

Others documented significant decrease in serum IgG concentrations in the febrile diarrhetic calves in comparison with that of non-febrile and control animals (Ashgan *et al.*, 2020). The differences between studies can be related to type of pathogen, age of the calf, stage of the disease, strength of passive immunity transfer of immunoglobulins in addition to other environmental conditions.

From the results obtained in this study it can be concluded that *E. coli*, *Salmonella spp.* and *Cryptosporidium* appeared to be the important causes of diarrhea in buffalo calves with the higher prevalence rate was recorded for *E. coli*. Also, the changes observed in serum concentrations of haptoglobin, ceruloplasmin, IL-6, IL-10 and IgA could provide a good information about the disease process and therefore, rise their clinical importance as valuable diagnostic and prognostic markers for monitoring or treating the disease.

REFERENCES

- Al-Alo, K.Z.K.; Nikbakhtbrujeni, G.H.; Lotfollahzadeh, S.; Moosakhani, F. And Gharabaghi, A. (2018): Passive protective effect of anti-K99 antibodies against enterotoxigenic *E.coli* infection in neonatal calves. Iran J. Vet. Med., Vol. 12, No 2 19(1): 3-8.
- Al-Zubaidi, M.T.S. (2014): Acute phase proteins in calves naturally infected with cryptosporidium. AL-Qadisiya Journal of Vet. Med. Sci. Vol. 14, No. 2.
- Ashgan, F.E.; Abeer, S. H. and Atia, A. E. (2020): Evaluation of Immunological Status of Calves Suffered from Diarrhea under Field Condition. Journal of Applied Veterinary Sciences, 5 (2): 40-48.
- Athanasiou, V.L., Katsoulos, D.P.; Ziogas, C.; Kassab A. and Polizopoulou, Z. (2019): Serum

- protein electrophoretic profile in diarrheic neonatal calves Comparative Clinical Pathology volume 28, 685–688.
- Aydin, F.; Umur, S.; Gocke, G.; Geno, O. and Guler, M.A. (2001): The isolation and identification of bacteria and parasites from diarrhoeic calves in Kars district. Kafkas Universitesi Veteriner Fakultesi Dergisi, 7 (1): 7-14.
- Azizzadeh, M.; Shooroki, H.F.; Kamalabadi, A.S. and Stevenson, M.A. (2012): Factors affecting calf mortality in Iranian Holstein dairy herds. Prev. Vet. Med. 104:335–340.
- Balikci, E. and Al, M. (2014): Some serum acute phase proteins and immunoglobulins concentrations in calves with Rotavirus, Coronavirus, E. coli F5 and Eimeria species. Iran J. Vet. Res., 15: 397-401.
- Bartels, C.J.; Holzhauer, M.; Jorritsma, R.; Swart, W.A. and Lam, T.J. (2010): Prevalence prediction and risk factors of enteropathogens in normal and non normal faces of young Dutch dairy calves. Prev. Vet. Med. 93: 162-169.
- Berchtold, J. (2009): Treatment of calf diarrhea: intravenous fluid therapy. Vet. Clin. North Am Food Anim Pract. 2009 Mar; 25(1):73-99, vi.
- Berge, A.C.; Besser, T.E.; Moore, D.A. And Sicho, W.M. (2009): Evaluation of the effects of oral colostrum supplementation during the first fourteen days on the health and performance of preweaned calves. J. Dairy Sci., Lancaster, Vol. 92, p. 286-295.
- Blanchard, P.C. (2012): Diagnostics of dairy and beef cattle diarrhea, Veterinary Clinics of North America Food Animal Practice, 28(3), 443–64.
- Chaurasia, R. (2001): Characterization, enterotoxigenicity and transmissible drug resistant of E. coli associated with calf diarrhoea. M.V.Sc. Thesis submitted to Rajasthan Agriculture University, Bikaner, India.
- Cho, Y.I. and Yoon, K.J. (2014): An overview of calf diarrhea infectious etiology, diagnosis, and intervention. J. Vet. Sci., 15: 1-17.
- Elvinger, F.; Hansen, P.J. and Natzke, R.P. (1991): Modulation of function of bovine polymorphonuclear leukocytes and lymphocytes by high temperature in vitro and in vivo. Am J Vet Res 52, 1692-1698.
- Espy, M.J.; Uhl, J.R.; Sloan, L.M.; Buckwalter, S.P.; Jones, M.F. and Vetter, E.A. (2006): Real-time PCR in clinical microbiology: applications for routine laboratory testing. Clin. Microbiol. Rev. 19, 165–256.
- Fossati, P.; Prencipe, L. and Berti, G. (1980): Enzymatic determination of uric acid Clin. Chem. 26(2): 227-231.
- Foster, D.M. and Smith, G.W., (2009): Pathophysiology of Diarrhoea in Calves. Veterinary Clinics of North America: Food Animal Practice, 25: (1) 13–36.
- Frank, S.M.; Bosworth, B.T. and Moon, H.W. (1998): Multiplex PCR for enterotoxigenic, attaching and effacing, and Shiga toxinproducing Escherichia coli strains from calves. J Clin Microbiol; 36: 1795–7.
- Gultekin, M.; Voyvoda, H.; Ural, K.; Erdogan, H.; Balikci, C. and Gultekin, G. (2019): Plasma citrulline, arginine, nitric oxide, and blood ammonia levels in neonatal calves with acute diarrhea. J. Vet. Intern. Med. 33: 987–998.
- Gurer, H.; Ozgunes, H.; Neal, R.; Spitzand D.R. and Ereal, N. (1998): Antioxidant effects of N-acetyl cysteine and succimer in red blood cells from lead exposed rats. Toxicology, 128, 181-189.
- Hajimohammadi, A.; Saeed, N.; Maryam, A.; Mohammad, R.K. and Saeed, M. B. (2013): Identifying relationships among acute phase proteins (haptoglobin, serum amyloid A, fibrinogen, ceruloplasmin) and clinical findings in dairy calf diarrhea. Comparative Clinical Pathology volume 22, pages 227–232.
- Hummelshoj, L.; Ryder, L.P. and Poulsen, L.K. (2006): The role of the interleukin-10 subfamily members in immunoglobulin production by human B cells. Scand J. Immunol. 64:40–47.
- Izzo, M.M.; Kirkland, P.D.; Mohler, V.L.; Perkins, N.R.; Gunn, A.A. and House, J.K. (2011): Prevalence of major enteric pathogens in Australian dairy calves with diarrhea. Aust Vet J., 89(5):167-73.
- Jellison, K.L.; Hemond, H.F. and Schauer, D.B. (2002): Sources and Species of *Cryptosporidium* Oocysts in the Wachusett Reservoir Watershed. App. and env. Microbiol., 68 (2): 569-575.
- Kazmierczak, S.C.; Kaplan, L.A. and Pesce, A.J. (1996): Ceruloplasmin. In: Clinical chemistry theory, analysis, correlation. St Louis: Mosby, :966pp.
- Kojouri, G.A.; Hassanpour, H.; Taghavi, N. and Taghadosi, C. (2012): Nitric oxide metabolites status in calves with acute and

- chronic diarrhea. Comparative Clinical Pathology. 21, pages 1257–1261.
- Koracevic, D.; Koracevic, G.; Djordjevic, V.; Andrejevic, S. and Cosic, V. (2001): Method for the measurement of antioxidant activity in human fluids. J. clin. Pathol., 54: 356-361.
- Lacetera, N.; Bernabucci, U.; Scalia, D.; Ronchi, B.; Kuzminsky, G. and Nardone, A. (2005): Lymphocyte functions in dairy cows in hot environment. Int J Biometeorol 50, 105-110.
- Maes, R.K.; Grooms, D.L.; Wise, A.G.; Han, C.; Ciesicki, V.; Hanson, L.; Vickers, M.J.; Kanitz, C. and Holland, R. (2003): Evaluation of a human group A Rotavirus assay for on-site detection of bovine rotavirus. J. Clin. Microbiol. 41: 290–294.
- Mahmoud, A.E.; Haidy, G.; Abdel-Rahman, E.; Abouelhassan, M. and Nahla, H. S. (2018): Cryptosporidium Parvum In Clinically Examined Dairy Calves; Molecular And Biochemical Studies. J. Egypt. Soc. Parasitol. (JESP), 48(3), 2018: 535 – 542.
- Mendonsa, K.M. (2011): Factors affecting passive transfer in neonatal calves. Dairy Science Department. California Polytechnic State University, San Luis Obispo, December.
- Montgomery, H.A.C. Dymock, J.F., (1961): Colometric method for determination of nitrite. Analyst. (86) 414.
- Moshage, H. (1997): Cytokines and the hepatic acute phase response. Journal of Pathology; 181:257-66.
- Murata, H.; Shimada, N. and Yoshioka, M. (2004): Current research on acute phase proteins in veterinary diagnosis: An overview. The Veterinary Journal, 168, 28–40.
- Murtaugh, M.P.; Baarsch, M.J.; Zhou, Y.; Scamurra, R.W. and Lin, G. (1996): Inflammatory cytokines in animal health and disease. Vet. Immunol. Immunopathol. 54, 45-55.
- Osman, M.M.; Mohammed, H.R.; Lamiaa, A.H. and Elham, A.A. (2020): Oxidative Stress Biomarkers and Pathological Alterations Induced by *Cryptosporidium* Infection in Buffalo Calves at Assiut Governorate, Egypt. Journal of Advanced Veterinary Research. Volume 10, Issue 2. 111-116.
- Park, S.H.; Saif, L. and Jeong, J.C. (2006): Molecular characterization of novel G5 bovine rotavirus strains,” Journal of Clinical Microbiology, vol. 44, no. 11, pp. 4101–4112.
- Petersen, H.H.; Nielsen, J.P. and Heegaard, P.M.H. (2004): Application of acute phase protein measurements in veterinary clinical chemistry. Veterinary Research; 35:163-87.
- Prescott, Bueschel, D.M.; Keel, M. K.; Glock, R.D.; Cuneo, P.; DeYoung, D.W.; Reggiardo, C.; Trinh, H.T. and Songer, J.G. (2008): A possible role for *Clostridium difficile* in the etiology of calf enteritis. Vet. Microbiol. Mar. 18: 127(3- 4): 343-352.
- Qadis, A.Q.; Goya, S.; Yatsu, M.; Kimura, A.; Ichijo, T. and Sato, S. (2014): Immune-stimulatory effects of a bacteria-based probiotic on peripheral leukocyte subpopulations and cytokine mRNA expression levels in scouring Holstein calves. J. Vet. Med. Sci. 76: 677–684.
- Quigley, J.D.; Kost, C.J. and Wolfe, T.M. (2002): Absorption of protein and IgG in calves fed a colostrum supplement or replacer. J. Dairy Sci. 85:1243–1248.
- Rachmilewitz, D.; Stamler, J.S.; Bachwich, D.; Karmeli, F.; Ackerman, Z. and Podolsky, D.K. (1995): Enhanced colonic nitric oxide generation and nitric oxide synthase activity in ulcerative colitis and Crohn' disease. Gut; 36:718-723.
- Rehab, E.D.; Gihan, M.O.; Mohammed, and Helal, I.M. (2016): Some Bacteriological and Molecular Studies on *Escherichia Coli* as Causative Agent of Calves Enteritis at Damietta Governorate. Egypt. J. Chem. Environ. Health, 2 (2): 195 -210.
- Rocha, T.G.; Silva, F.D.F.; Bortoletto, C.; Silva, D.G.; Buzinaro, M.G.; Zafalon, L.F. and Fagliari, J.J. (2016): Serum concentrations of acute phase proteins and immunoglobulins of calves with rotavirus diarrhea. Arq. Bras. Med. Vet. Zootec., Vol. 68, No. 4, p.865-872.
- Sadrzadeh, S.M. and Bozorgmehr, J. (2004): Haptoglobin phenotypes in health and disorders. Am. J. Clin. Pathol., 121 Suppl: S97–104.
- Satoh, K. (1978): Serum lipoperoxides in cerebrovascular disorders determined by colorimetric method. Clin. Chim. Acta. 90: 37-43.
- Seid, U.; Dawo, F.; Tesfaye, A. and Ahmednur, M. (2020): Isolation and Characterization of Coronavirus and Rotavirus Associated with Calves in Central Part of Oromia, Ethiopi. Veterinary Medicine International. 10 pages.
- Silva, D.G.; Silva, P.R.L.; Silva, P.C. and Fagliari, J.J. (2011): Serum protein concentrations, including acute phase proteins, in calves

- experimentally infected with Salmonella Dublin. *Pesqui. Vet. Bras.*, v.31, p.551-554.
- Snedecor, G.W. and Cochran, W. G. (1980): *Statistical Methods*. 7th edn Iowa State University Press. USA.
- Sowmyanarayanan, T.V.;Natarajan, S.K.; Ramachandran, A.; Sarkar, R.; Moses, P.D.; Simon, A.; Agarwal, I.; Christopher, S. and Kang, G. (2009): Nitric oxide production in acute gastroenteritis in Indian children. *Trans. R. Soc. Trop. Med. Hyg.*, 103(8): 849-51.
- Stephani, F.; Rolf, B.; Claus, P. C. and Stephan, N. (2016): Serum Interleukin 6 as a Prognostic Marker in Neonatal Calf Diarrhea. *J. Dairy Sci.* 99:6563–6571. Vol. 99, No. 8.
- Swamy, S.C.; Barnhard, H.M.; Lee, H.D. and Dresen, D.W. (1996): Virulence determinant *invA* and *spvC* in salmonella isolated from poultry product, waste water, and human sources. *App Environ Microbiol*; 62:3768–71.
- Tsunemitsu, H.; Smith, D.R. and Saif, L.J. (1999): Experimental inoculation of adult dairy cows with bovine coronavirus and detection of coronavirus in feces by RT-PCR,” *Archives of Virology*, vol. 144, no. 1, pp. 167–175.
- Uhde, F.L.K.; Aufmann, T.; Sager, H.; Albini, S.; Zanoni, R.; Schelling, E. and Meylan, M. (2008): Prevalence of four enteropathogens in the faeces of young diarrheic dairy calves in Switzerland. *Vet. Rec.* 163: 362-366.
- Youssef, A.M.; El-Ashker, R.M. and Ouda, F.M. (2017): Modulation of immunity and inflammatory gene expression in buffalo (*Bubalus bubalis*) with some digestive disorders. *Comparative Clinical Pathology*. volume 26, pages1123–1128