

VITRIFICATION AND IN VITRO MATURATION AND FERTILIZATION OF BOVINE OOCYTES WITH OR WITHOUT CUMULUS CELL LAYER.

Hammad, M.E.*; Sh.A. Gabr.*; I.T. El - Ratel** and Eman A. Amin*

* Anim. Prod. Dept., Fac. Agric., Tanta Univ., Egypt.

**Biotech. Dept., Anim. Prod. Res. Inst., Agric. Res. Center, Egypt.



ABSTRACT

Aim of this study was to evaluate the effect of presence or absence of cumulus cells around oocytes, on post-vitrification survival rate, maturation and developmental capacity of bovine oocytes. Ovaries were collected from abattoirs and all visible follicles were aspirated and examined. Only, cumulus cell complex (COCs) and natural (NDOs) or mechanically denuded (MDOs) were vitrified/warmed, matured, fertilized and cultured for 7 days to calculate post-thawing survival, maturation, fertilization and blastocyst production rates. Results showed that the proportions of total and morphological normal oocytes recovered post-vitrification were greater ($P<0.05$) for COCs (79.0 and 76.0%) than those obtained for NDOs (56.0 and 51.4%). Also, post-vitrification recovery rate and proportion of morphological normal oocytes was greater ($P<0.05$) for MDOs (67.8 and 63.2%) than those for NDOs (56.0 and 51.4%). Proportion of damaged oocytes post-vitrification was not affected significantly by oocyte type, but leakage of cellular contents represented the highest frequency of damaged oocytes, while splitting in two halves was the lowest frequent abnormality observed. The COCs yielded the highest ($P<0.05$) percentages of oocytes at M II (maturation rate) and the lowest percentages of degenerated oocytes, followed by MDOs, while NDOs showed the lowest percentages, being 60; 10% for COCs, 46.9; 20.5% for MDOs and 38.9; 30.4% for NDOs, respectively. Cleavage rate was higher ($P<0.05$) for COCs than those of MDOs and NDOs (40.0 vs. 26.3 and 21.3%, respectively). Production rate of embryos at morula and blastocyst stages was not affected significantly by oocyte type. In conclusion, cumulus cell layer surrounding the bovine oocytes recovered from ovaries collected from abattoirs play very important roles for maintain viability of oocytes during vitrification, and successfully *in vitro* matured, fertilized and developed to blastocyst stage.

Keywords: Bovine oocytes, cumulus, denudation, vitrification, maturation, fertilization

INTRODUCTION

Achievement a high efficiency of bovine oocyte cryopreservation remains for years of great interest of researchers and practitioners working on *in vitro* production of bovine embryos and somatic cloning (Papis *et al.* 2013). Cryopreservation of oocytes from slaughtered animals has great value in increasing the availability of materials for basic research and their subsequent utilization for embryos production may provide an opportunity to replenish the endangered species gene banking and the genetic improvement of the livestock species (Nucharin Sripunya, 2011).

Several cryopreservation methods such as conventional (slow), equilibrium rapid freezing (vitrification) and ultra-rapid freezing have been used to preserve embryos and oocytes of many animal species resulting in the birth of live offspring (El-Shahat and Hammam, 2014). These methods

are highly dependent upon cryoprotectant agents (CPAs) that protect oocytes from damage during the freezing process (Sanchez-Partida *et al.* 2011). A higher number of competent oocytes for *in vitro* maturation and *in vitro* fertilization (IVM – IVF) to obtain superior transferable bovine embryos coupled with development of freezing technique through vitrification will entail more productivity from the non-descript animals (Dutta *et al.* 2013).

One of the factors that could affect oocyte quality following vitrification is the presence or absence of cumulus cells around the oocyte prior to cryopreservation (Karima *et al.* 2014). It have been reported that presence of cumulus cells is beneficial to the oocyte survival after cryopreservation (Li *et al.* 2006); as it may minimize the release of cortical granules and prevent premature zona reaction, thereby cumulus cell improve the *in vitro* fertilization rates after cryopreservation (Vincent *et al.* 1990). Cumulus cell removal prior to *in vitro* maturation or vitrification have shown to have a detrimental effect on oocyte morphology for immature vitrified buffalo (Gasparrini *et al.* 2007), equine (Tharasanit *et al.* 2009), mouse (Suo *et al.* 2009), bovine (Zhou *et al.* 2010), and goat (Purohit *et al.* 2012) oocytes. In goat, Purohit *et al.* (2012) found that cumulus compact oocytes are less vulnerable to cryo-injuries compared to their denuded counterparts.

The immature oocyte should be considered as a one functional unit with cumulus cells communicating each other via cellular projections penetrating across a zona pellucida. Undisturbed oocyte-cumulus communication and co-operation seem essential for adequate maturation process (Gilchrist, 2011), so an optimum cryopreservation method should provide protection for oocyte, cumulus cells and their intercommunication system.

Although immature oocytes vitrified without cumulus were matured and fertilized *in vitro* with acceptable efficiency (Modina *et al.* 2004; Luciano *et al.* 2009; Zhou *et al.* 2010), their further development as embryos was compromised (Modina *et al.* 2004). However, less information is known about the efficiency and the consequences of cryopreservation on immature and *in vitro* matured oocytes (Fausta *et al.* 2013).

Therefore, the present study aimed to evaluate the effect of, presence or absence of cumulus cells on immature bovine oocytes, on post-vitrification survival rate and developmental capacity.

MATERIALS AND METHODS

This study was carried out at the International Livestock Management Training Center (ILMTC), belonging to the Animal Production Research Institute, Agriculture Research Center, Ministry of Agriculture, in cooperation with Department of Animal Production, Faculty of Agriculture, Tanta University.

Oocyte collection:

Bovine ovaries were collected from abattoirs and transported within 3 h to the laboratory in normal saline (0.9% NaCl) containing gentamicin (50 µg/ml) at 27-30°C. In the laboratory, extraneous tissues on the ovarian

surface were removed and the ovaries were washed three times in phosphate buffer saline (PBS, pH 7.3). All visible follicles (3-8 mm in diameter) were aspirated using 18-gauge needle attached to a 5 ml syringe containing 2 ml of DPBS with 20% fetal calve serum (FCS, sigma) and antibiotics (50 µg/ml gentamicin).

Oocytes were examined under stereomicroscopy and classified according to their compaction, number of cumulus cell layers and homogeneity of ooplasm according to Ravindranatha *et al.* (2003) into 4 categories namely cumulus oocytes-complexes (COCs), expanded cumulus cells oocytes, natural denuded oocytes (NDOs), partial denuded oocytes. Only, COCs and NDOs (natural or mechanically denuded) were used in this study.

Mechanical denudation of oocytes:

Natural COCs were mechanically denuded (MDOs) by repeated pipetting in PBS supplemented 0.025% hyaluronidase solution (SIGMA, St. Louis, MO) till the complete separation of the cluster of cumulus cells according to Papis *et al.* (2013).

Vitrification of oocytes:

Different type of oocytes (COCs, NDOs or MDOs) were vitrified by open-pulled straw cryodevice (OPS). The vitrification procedures employed throughout this experiment were based on the methods originally designed by (Shayegh and Barati, 2011) with minor modifications. TCM-199 medium (Sigma) supplemented with 20% (v: v) of FCS as a basic medium (BM) as well as ethylin glycol (EG) and dimethyl sulfoxide (DMSO) as cryoprotectants were used. Different types of oocytes were vitrified by placing them in the first vitrification solution (V1, 10% EG+10% DMSO in BM) for 5 min, then they were transferred into the second vitrification solution (VS2: 20% DMSO, 20% EG, and 0.5M sucrose in BM) for 30 s, instantly oocytes were loaded in OPS and plunged in liquid nitrogen (LN2).

Thawing and evaluation of oocyte viability:

After storage for at least 2 weeks in LN2, all types of vitrified oocytes (COCs, NDOs and MDOs) were warmed by holding the OPS for 6 s in air and then agitating them in water bath at 20 °C for at least 10 s. The contents of OPS were expelled into Petri dish. To remove of intracellular cryoprotectants effects, oocytes were transferred in BM plus 0.25M sucrose for 5 min and then transferred to buffer solution (BS) plus 0.125M sucrose solution for 5 min and finally, the oocytes were washed twice in BS without sucrose for 5 min according to Hajarian *et al.* (2011) with minor modifications.

Oocyte viability (survival) was evaluated morphologically based on the integrity of the oolemma and zona pellucida; loss of membrane integrity (lysis) was obvious upon visual inspection as the sharp demarcation of the membrane disappeared and the appearance of the cytoplasm changed. The criteria used for assessing the post-thaw morphology of vitrified/warmed oocytes were as follows: Normal oocytes with spherical and symmetrical shape with no sing of lysis, and damaged oocytes (abnormal) with crack in zona pellucida, split in two halves, change in shape and leakage of contents.

The survival rate was calculated as the proportion of normal morphology oocytes against the total number of vitrified oocytes. Thereafter, morphologically normal oocytes were matured and fertilized *in vitro*.

In vitro maturation:

All types of oocytes (COCs, NDO and MDO), morphologically normal post-thawing, were cultured in 100 µl of TCM-199+20% FCS+1 µg/ml oestradiol-17β (Sigma) and 50 µg/ml gentamicin covered with mineral oil (Sigma) in four-well culture plates (8-10 oocytes per droplet) for 24 h in a CO₂ incubator (5% CO₂ and in humidified air at 38°C). Oocytes were washed using phosphate buffer solution (PBS) containing 1 mg /ml hyaluronidase to remove the cumulus cells. Then, oocytes were washed two times with PBS supplemented with 2% bovine serum albumin (BSA), and loaded on clean slide. Slides were placed into fixation solution (3 ethanol: 1 glacial acetic acid) for 24 h and stained with 1% orcein in 45% glacial acetic acid. The nuclear status of oocytes was evaluated under a microscope and considered to be matured if they were at metaphase II stage with reduced number of chromatin, metaphase plate and extrusion of the 1st polar body (Purohit *et al.* 2012).

Sperm preparation and in vitro fertilization:

Spermatozoa were recovered from frozen semen by swim-up separation in Tyrode's Albumin Lactate Pyruvate medium (TALP) previously described by (Parrish *et al.* 1988) with some modifications. One straw of frozen Holstein bull (0.25 ml) was thawed in a water bath at 37.5 °C for 1 min. The contents were washed twice in 5 ml of sperm-TALP supplemented by 6 mg/ml BSA (fraction V, A- sigma) by centrifugation at room temperature for 5 min to remove extender and cryoprotectants. The sediment of spermatozoa was resuspended in 5 ml of IVF-TALP containing 10 µg/ml heparin and then centrifuged for 5 min. The supernatant was removed leaving 0.25-0.5 ml of IVF-TALP and sperm pellet. The concentration of sperm was adjusted by adding IVF-TALP medium to reach 1x10⁶ sperm/ml as tested by a haemocytometer.

Fertilization Process:

After *in vitro* maturation, all matured oocytes from each type were washed two times in sperm-TALP, followed by final washing in fertilization medium (IVF-TALP). *In vitro* matured oocytes were transferred into Petri dish containing 100 µl droplets of fertilization medium at the rates of (10 oocytes per drop). Aliquots of the sperm suspension (8 µl) were added to each droplet containing matured oocytes. The oocytes and spermatozoa were co-cultured in an CO₂ incubator at 38.5°C in 5% CO₂ in air, with saturated humidity for 24 h.

In vitro culture and embryo development:

After 24 h co-incubation of spermatozoa and oocytes from each type, the presumptive embryos were washed in sperm-TALP medium. The final washing was done in a culture medium consisting of TCM-199 supplemented with 3 mg/ml BSA, 20 µg/ml Na-pyruvate and 50 µg/ml gentamycin.

After co-incubation, presumptive embryos were placed in petri dish in the culture medium covered with mineral oil and incubated at 38.5°C under 5% CO₂ in humidified air for 7 days. The frequency of morula and/or

blastocyst was recorded. Medium was replaced with fresh medium after every 48 h of culture (Dutta *et al.* 2013).

Statistical analysis:

Data were analyzed by analysis of variance using computer program of SAS (2000). The significant differences among group means were performed using Duncan Range Test (Duncan, 1955).

RESULTS AND DISCUSSION

Effect of oocyte type on:

Post-thawing recovery rate and quality of vitrified oocytes:

Data in Table (1) showed that the proportions of total and morphological normal oocytes found to be recovered post-vitrification were significantly ($P<0.05$) greater for COCs (79.0 and 76.0%) than those obtained for NDOs (56.0 and 51.4%). Also, post-vitrification recovery rate and proportion of morphological normal oocytes was significantly ($P<0.05$) greater for MDOs (67.8 and 63.2%) than those for NDOs (56.0 and 51.4%). However, proportion of damaged oocytes post-vitrification showed an opposite trend.

Table: (1): Effect of oocyte type on recovery and quality of vitrified oocytes.

Oocyte type	Vitrified oocytes (n)	Recovery rate (%)	Oocytes viability (%)	
			Viable	Damage
COCs	100	79.00±1.00 ^a	76.00±1.00 ^a	24.00±1.00 ^c
NDOs	125	56.00±0.00 ^c	51.43±1.43 ^c	48.57±1.43 ^a
MDOs	115	67.80±0.80 ^b	63.17±1.98 ^b	36.82±1.98 ^b

^{a, b} and ^c: Means within the same column with different superscripts are significantly different at $P<0.05$.

In accordance with the present results, Prentice *et al.* (2012) reported that the survival and developmental rate of bovine oocytes were higher when vitrified with enclosed cumulus cells than partially denuded cells. They showed that the survival rate and normal morphologically immature oocytes of goats was significantly higher for COCs than MDOs (86.73 and 89.22% vs. 80.31 and 94.12%, respectively). In buffalo, Karima *et al.* (2004) found that percentage of morphological normal buffalo oocytes (COCs) after vitrifying, was non-significantly differing with partially DOs, while there were significant ($P<0.01$) difference between COCs and NDOs (91.6 vs. 78.0%). Dhali *et al.* (2000) reported that the percentage of buffalo COCs found to be morphologically normal which varied from 89 to 96% for the two equilibration solutions and the two exposure times was not significantly different.

In this respect, Zhou *et al.* (2010) studied the effects of cumulus cells on bovine oocyte vitrification at the GV stage. They reported that the survival rate of vitrified oocytes was significantly higher for COCs than partially-denuded vitrified and control oocytes. Also, Babaei *et al.* (2006) reported a

high proportion of morphologically normal bovine oocytes (90%) were recovered after vitrification-warming using glass capillary micropipette.

Moreover, Nikseresht *et al.* (2015) showed that the survival rate of mice oocytes in the stepwise cumulus oocytes complexes and denuded oocytes group were significantly higher than those for the single-step NDOs. Mahmoud *et al.* (2013) reported that the rate of morphologically intact oocytes following vitrification / warming was high, ranging from 87.7% in straws to 90.8% in cryotops using a mixture of 3 M DMSO + 3 M EG.

It was suggested that the presence of the cumulus cells can reduce the adverse effects of DMSO on the oocytes (Johnson and Packer, 1987). Also, the presence of cumulus cells can minimize the release of cortical granules and premature zona reaction for zona hardening resulting in low fertilization rates (Vincent *et al.* 1990).

On the other hand, Chian *et al.* (2004) reported that bovine oocytes survival rates following vitrification are not affected by the presence or absence of cumulus cells. Also, Zhang *et al.* (2009) observed no difference in the survival rate of vitrified mature ovine oocytes with or without cumulus cells. The difficulty, in obtaining acceptable rates of survival and functionality for oocytes after cryopreservation, is due to the size of this cell and its unique morphologic characteristics (Martins *et al.* 2005).

Classification of damaged oocytes post-vitrification:

Various degree of damage were observed in abnormal oocytes post- vitrification included cracking zona pellucida, split into two halves, change in shape and leakage of cellular content (Table 2). Similar findings were observed by Dhali *et al.* (2000) for morphological abnormalities observed in buffalo oocytes after vitrification-thawing.

Proportion of damaged oocytes post-vitrification was not affected significantly by oocyte type, but leakage of cellular contents represented the highest frequency of damaged oocytes, while splitting in two halves was the lowest frequent abnormality observed. In agreement with the present results, Dhali *et al.* (2000) showed that morphological abnormalities observed in buffalo oocytes after vitrification-thawing were, crack in zona pellucida (69%), split into two halves (3%), change in shape (5%) and leakage of cellular contents (36%). Also in buffalo, Hammam and El-Shahat (2005) found that types of damages observed after vitrification-thawing were, 16.6% crack in zona pellucida, 50% shrinkage of cytoplasm and 33.3% leakage of cellular contents. However, Babaei *et al.* (2006) found that the percentage of morphologically damaged bovine oocytes was 10% and mostly cracking zona pellucid. Recently, El-Shahat and Hammam (2014) reported that among the damaged oocytes, cracking of zona pellucida was the most frequent abnormality observed compared to shrinkage of cytoplasm and leakage of cellular content (53.3 % vs. 20.0 and 26.6%).

Table: (2): Effect of oocyte type on classification of damaged oocytes post-vitrification.

Oocyte type	Damaged oocytes (n)	Oocyte classification (%)			
		Crack in zona pellucida	Splitting into two halves	Change in shape	Leakage of cellular contents
COCs	19.00	20.00±5.00	15.00±6.12	21.67±5.65	43.33±8.07
NDOs	34.00	26.67±3.23	17.62±2.77	29.52±0.95	26.19±2.38
MDOs	29.00	32.10±4.96	13.05±3.42	18.05±2.02	36.80±3.42

***In vitro* maturation of vitrified oocytes:**

Results in Table (3) revealed that COCs yielded significantly ($P<0.05$) the highest percentages of oocytes at M II (maturation rate) and the lowest percentages of degenerated oocytes, followed by MDOs, while NDOs showed the lowest percentages, being 60; 10% for COCs, 46.9; 20.5% for MDOs and 38.9; 30.4% for NDOs, respectively.

Table: (3): Effect of type of vitrified on *in vitro* maturation of vitrified oocytes post-thawing.

Item	N	Oocyte stage (%)				
		GV	GVB	M I	M II	Degenerated
COCs	60	8.3±0.00	8.3±0.00	13.3±2.04	60.0±1.66 ^a	10.0±1.66 ^c
NDOs	36	11.4±2.85	11.1±2.78	8.2±3.36	38.9±2.78 ^c	30.4±1.78 ^a
MDOs	49	10.3±0.38	12.1±1.54	10.26±0.38	46.9±1.29 ^b	20.5±0.76 ^b

^{a, b} and ^c: Means within the same column with different superscripts are significantly different at $P<0.05$. N: Number of viable oocytes. GV: Germinal vesicle. GVB: Germinal vesicle breakdown. M I: Metaphase I. M II: Metaphase II.

In harmony with the present results, Karima *et al.* (2014) showed the highest maturation rate of COCs, followed by partial DOs (73.7 vs. 64.8%), while NDOs showed the lowest maturation rate (53.0%). In goat, Purohit *et al.* (2012) found that immature COCs had significantly higher maturation rates compared to their MDOs (41.25 vs. 27.48%, respectively). In mice, Nikseresht *et al.* (2015) found that maturation rate in the stepwise COCs was significantly higher than in single step NDOs (73.2 vs. 50.8%).

Cumulus cells are closely related to oocytes, forming in fact one functional unit reflecting each other on auto- or paracrine manner (Gilchrist, 2011). Cumulus-oocyte interrelationship during oocyte maturation process seems crucial for achievement of optimum oocyte maturation and developmental capacity (Papis *et al.* 2013). The interactions between the oocyte and its surrounding cumulus cells at this stage are crucial for development of a matured oocyte (MII), the only cell type that can be fertilized to initiate a new organism (Sutton *et al.* 2003). It was reported that cumulus cell removal prior to *in vitro* maturation or vitrification has a detrimental effect on oocyte morphology for both immature and mature vitrified buffalo (Gasparini *et al.* 2007), equine (Tharasanit *et al.* 2009), mouse (Suo *et al.* 2009) and bovine (Modina *et al.* 2004) oocytes. Cumulus cell removal increases the maturation promoting factor activity and accelerates the transition to metaphase stage and the redistribution of cortical

granules (Ge *et al.* 2008). Therefore, according to these findings and those reported in the present study, immature bovine oocytes can be vitrified after cumulus cell removal, and that they successfully mature (Modina *et al.* 2004). Finally, It has been reported that the absence of cumulus cells could provoke a possible shortcoming in protein synthesis and could reflect the levels of molecules involved in the regulation of meiotic and mitotic cell cycles (Combelles *et al.* 2005).

In vitro fertilization of vitrified-matured oocytes:

Data presented in Table (4) showed that cleavage rate was significantly ($P<0.05$) higher for COCs than those of MDOs and NDOs (40.0 vs. 26.3 and 21.3%, respectively). However, production rate of embryos at morula and blastocyst stages was not affected significantly by oocyte type.

In accordance with the present results, Purohit *et al.* (2012) showed significantly ($P<0.05$) higher proportion of fertilized oocytes for immature COCs of goat than immature MDOs (31.7 vs. 25.0%, respectively).

Table: (4): Effect of category of vitrified oocytes on cleavage and development rates.

Item	Type of oocytes		
	COCs	NDOs	MDOs
Vitrified oocytes, n	80	75	80
Cleaved oocytes, n	32	16	21
Cleaved oocytes, %	40.00±0.00 ^a	21.33±1.25 ^b	26.25±1.25 ^b
Embryonic stage:			
2-cell embryos, n	25	14	20
2-cell embryos, %	78.13±3.12	87.50±5.00	95.24±6.57
Morulae, n	4	1	1
Morulae, %	12.50±0.00	6.25±5.00	4.76±5.00
Blastocysts, n	3	1	0
Blastocysts, %	9.37±3.12	6.25±6.25	0

^{a, b and c:} Means denoted within the same row with different superscripts are significantly different at $P<0.05$.

Also, Modina *et al.* (2004) observed a lower percentage of fresh NDOs reached the blastocyst stage in comparison with intact COCs (23.9 vs. 35.4, $P<0.05$). The blastocyst rate was 4.3%, being lower than those reported in the current study (6.25-9.37%). In fact, this is one of the evidences of successful oocyte cryopreservation for which, to date, only controversial and sporadic data are available for the bovine species (Hochi, 2003). Moreover, Zhou *et al.* (2014) achieved cleavage (63.5%) and blastocyst development (20.0%) after parthenogenetic activation of vitrified-warmed bovine oocytes similar to that from oocytes vitrified by the open-pulled straw method (57.0% cleavage and 23.0% blastocyst development, respectively (Hou *et al.* 2005). Low blastocyst development rates (less than 10%) was reported by Martins *et al.* (2005). In buffalo, Hammam and El-Shahat (2005) found that oocytes vitrified at the immature cleaved and developed into morula and blastocyst stage after thawing were 20.0, 3.3 and 2.0%, respectively). The developmental capacity of the vitrified-thawed immature buffalo oocytes was

significantly lower compared to control (Hammam and El-Shahat., 2014; Yadav *et al.* 2008). Furthermore, Mahmoudi *et al.* (2005) reported that intact mouse oocytes had a higher developmental competence than denuded oocytes. In this respect, Hoshi *et al.* (1998) vitrified immature bovine oocytes in straws by using a mixture of 40% EG, ficoll and sucrose as a vitrification medium. They reported 47.5% fertilization rate from the vitrified bovine oocytes. Abe *et al.* (2005) reported developmental rates to blastocyst of bovine GV-COCs, using Nylon-Mesh and exposer with stepwise cryoprotectant, being significantly higher than with the single-step vitrification.

In this respect, Zhou *et al.* (2010) reported that cleavage and blastocyst rates of cumulus-enclosed vitrified bovine oocytes of GV were significantly higher than those of partially-denuded vitrified and control oocytes.

Studies have shown that GV-stage oocytes which are stripped of cumulus cells have a reduced developmental capacity compared with that of cumulus-enclosed GV-stage oocytes (Mahmoudi *et al.* 2009). Therefore, Cumulus cells play an important role in oocyte maturation since they provide and transfer several known and unknown factors that are essential for normal nuclear and cytoplasmic maturation of oocytes and subsequent embryonic development after fertilization (Mahmoudi *et al.* 2012).

CONCLUSION

Based on the foregoing results in the present study, immature bovine oocytes can be vitrified after cumulus cell removal, and that they successfully mature and develop up to the blastocyst stage after *in vitro* fertilization as reported by Modina *et al.* (2004). Recently, El-Shahat and Hammam (2014) observed that the immature germinal vesicle stage oocytes tolerate the cryopreservation damage more efficiently compared to oocytes at metaphase-II and cumulus compact oocytes are less vulnerable to cryo-injuries compared to their denuded counterpart.

In conclusion, cumulus cell layer surrounding the bovine oocytes recovered from ovaries collected from abattoirs play very important roles for maintain viability of oocytes during vitrification, and successfully *in vitro* matured, fertilized and developed to blastocyst stage.

REFERENCES

- Abe, Y.; Hara, K.; Matsumoto, H.; Kobayashi, J.; Sasada, H. and Ekwall, H. (2005). Feasibility of a nylon-mesh holder for vitrification of bovine germinal vesicle oocytes in subsequent production of viable blastocysts. *Biol. Reprod.* 72: 1416-20.
- Gasparini, B.; Attanasio, L.; De Rosa, A.; Monaco, E.; Di Palo, R. and Campanile, G.(2007). Cryopreservation of *in vitro* matured buffalo (*Bubalus bubalis*) oocytes by minimum volumes vitrification methods. *Anim. Reprod. Sci.*, 98:335–342.

- Babaei, H.; Nematollahi, M. S. N; Kheradmand, A. and Ayen, E. (2006). *In vitro* survival rate of bovine oocytes following vitrification in glass capillary micropipette (GCM). Iranian Journal of Veterinary Research, University of Shiraz, Vol. 7, No. 1, Ser. No. 14.
- Chian, R.C.; Kuwayama, M. Tan, L.; Tan, J.; Kato, O. and Nagai, T. (2004). High survival rate of bovine oocytes matured *in vitro* following vitrification. J. Reprod. Dev. 2004; 506: 685-96.
- Combelles, M.H.; Fissore, R.A.; Albertini, D.F. and Racowsky, C., (2005). *In vitro* maturation of human oocytes and cumulus cells using a coculture three-dimensional collagen gel system. Hum. Reprod., 20:1349–1358.
- Dhali, A.; R.S. Manik, S.K.; Das, S.K. and Singla, P. (2000). Post-vitrification survival and *in vitro* maturation rate of buffalo (*Bubalus bubalis*) oocytes: effect of ethylene glycol concentration and exposure time. Animal Reproduction Science 63 (2000) 159–165.
- Duncan, D. B. (1955). Multiple Range and Multiple F-Test. Biometrics, 11:1-42.
- Dutta, D. J.; Hiramoni, D. and Himangshu, R. (2013). *In vitro* blastocyst development of post-thaw vitrified bovine oocytes. Veterinary World, EISSN: 2231-0916.
- Fausta, B.; Maria, C. G.; Giovanni, C.; Mario Mignini, R.; Mariabeatrice, D. C. and Rubens, F. (2013). The current challenges to efficient immature oocyte cryopreservation. J Assist Reprod Genet (2013) 30:1531–1539.
- Ge, L.; Sui, H.S.; Lan, G.C.; Liu, N.; Wang, J.Z. and Tan, J.H. (2008) Coculture with cumulus cells improves maturation of mouse oocytes denuded of the cumulus oophorus: observations of nuclear and cytoplasmic events. Fertil Steril. 90(6):2376-88.
- Gilchrist, R.B. (2011). Recent insights into oocyte-follicle cell interactions provide opportunities for the development of new approaches to *in vitro* maturation. Reproduction Fertility and Development 23, 23-31.
- Hajarian, H. Wahid, H.; Rosnina Y.; Daliri, M.; Dashtizad, M.; Karamishabankareh, H. and Abas, M. O.. (2011). Cryotop and development of vitrified immature bovine oocytes. Arq. Bras. Med. Vet. Zootec., v.63, n.1, p.67-73.
- Hammam, A. M. and El-Shahat, K. H. (2005). vitrification of immature and mature buffalo oocytes in glycerol solution by a simple method. ISAH - Warsaw, Poland-Vol 1. 252-255.
- Hochi, S. (2002). Cryopreservation of follicular oocytes and preimplantation embryos in cattle and horses. J Reprod Dev 2003;49:13-21.
- Hochi, S.; Ito, K.; Hirabayashi, M.; Ueda, M. and Kimura, K. (1998). Effect of nuclear stages during IVM on the survival of vitrified warmed bovine oocytes. Theriogenology, 49: 787-796.
- Hou, Y.; Dai, Y.; Zhu, S.; Zhu, H.; Wu T.; Gong, G.; Wang, H.; Wang, L.; Liu, Y. and Li, R. (2005). Bovine oocytes vitrified by the open pulled straw method and used for somatic cell cloning supported development to term. Theriogenology. 2005;64:1381–1391.
- Johnson, M.H. and Pickering, S.J. (1987). The effect of dimethylsulphoxide on the microtubular system of the mouse oocyte. Development. 100:313–324.

- El-Shahat, K.H. and Hammam, A.M. (2014). Effect of different types of cryoprotectants on developmental capacity of vitrified-thawed immature buffalo oocytes. *Anim. Reprod.*, v.11, n.4, p.543-548.
- Karima, G.h .; Mahmoud,M.; Al-Shimaa, Al-H.; El-Naby,H.; Ahmed Y. F.; Eldebaky, H. A. A.; Abou El-Roos, M.E. A. and Abd- El-Ghaffar, A. E. (2014). Effect of Cumulus Cells and Meiotic Stages on Survivability and Meiotic Competence in Vitrified Buffalo Oocytes. *Pakistan J. Zool.*, vol. 46(5), pp. 1185-1192.
- Li, G.P.; Bunch, T.D.; White, K.L.; rickords, L.; LIU, Y. and Sessions, B.R. (2006). Denuding and centrifugation of maturing bovine oocytes alters oocyte spindle integrity and the ability of cytoplasm to support parthenogenetic and nuclear transfer embryo development. *Mol. Reprod. Dev.*, 73: 446–451.
- Luciano, A.M.; Franciosi, F.; Lodde, V.; Perazzoli, F.; Slezakova, M.and Modena, S. (2009). Cryopreservation of immature bovine oocytes to reconstruct artificial gametes by germinal of early stage bovine *in vitro* produced embryos with the open pulled straw (OPS) method. *Cryoletters*. 18: 191-195.
- Mahmoud, K.G.M.; El-Sokary, M.M.M.; Scholkamy Abou El-Roos M.E.A.; Sosa, G.A.M.; Nawito, M.F. (2013). The effect of cryodevice and cryoprotectant concentration on buffalo oocytes vitrified at MII stage. *Anim Reprod*, 10:689-696.
- Mahmoudi, R.; Abbasi, M.; Amiri, I.; Ragardi Kashani, I.; Pasbakhsh, P.; Saadipour, Kh. et al., (2009). Cumulus cell role on mouse germinal vesicle oocyte maturation, fertilization, and subsequent embryo development to blastocyst stage *in vitro*. *Yakhteh Medical Journal*. 11: 299-302.
- Mahmoudi, R.; Rajaei, F.; Ragardi Kashani, I.; Abbasi, M.; Amidi, F.; Sobhani, A. et al(2012). The rate of blastocysts production following vitrification with step-wise equilibration of immature mouse oocytes. *Iran. J. Reprod. Med.*,10:453-458.
- Mahmoudi, R.; Subhani, A.; Pasbakhsh, P.; Abolhasani, F.; Amiri, I.; Salehnia, M. *et al.* (2005). The Effects of cumulus cells on *in vitro* maturation of mouse germinal vesicle stage oocytes. *IJRM*; 3:74-78.
- Modina, S.; Beretta, M.; Lodde,V.; Lauria, A. and Luciano, A.M. (2004). Cytoplasmic changes and developmental competence of bovine oocytes cryopreserved without cumulus cells. *Eur. J. Histochem.*, 48: 337-346.
- Nikseresht, M.; Mehdi, A. T.; Tahere, R.,; Iraj, R.and RezaMahmoudi k. (2015). The Nuclear Maturation and Embryo Development of Mice Germinal Vesicle Oocytes with and without Cumulus Cell after Vitrification. *Journal of Clinical and Diagnostic Research*. 2015 Jan, Vol-9(1): AF01-AF04.
- Nucharin Sripunya. (2011). cryopreservation of geminal vesicle and metaphase-II stages bovine oocytes by cryotop and solid surface vitrification. Master of Science in Biotechnology ,Suranaree University of Technology.

- Papis, K.; Manabu, Sh.; Sukumar, S.; Yoshiaki, I. and Jacek, A. (2013). Effects of vitrification of partially denuded bovine immature oocytes. *Animal Science Papers and Reports* vol. 31.no. 1, 5-14.
- Parrish, J. J.; Susko-Parrish, J. L. and First, N. L. (1988). Capacitation of bovine sperm by heparin. *Biol. Reprod.*, 38: 1171-1180.
- Prentice-Biensch, J.R.; Singh. J.; Mapletoft. R.J. and Anzar, M. (2012): Vitrification of immature bovine cumulus oocytes complexes: effects of cryoprotectants, the vitrification procedure and warming time on cleavage and embryo development. *Reproductive Biology and Endocrinology*, 10:73.
- Purohit, G. N.; Harikesh, M. and Kanika, S. (2012). Effects of Vitrification on Immature and *in vitro* Matured, Denuded and Cumulus Compact Goat Oocytes and Their Subsequent Fertilization. *J Reprod Infertil*. 2012;13(1):53-59.
- Martins, R. D.; Costa, E. P.; Chagas, J. S. C.; Ignácio, F. S.; Torres, C. A. A. and McManus, C. (2004). Effects of vitrification of immature bovine oocytes on *in vitro* maturation. *Anim. Reprod.*, v.2, n.2, p.128-134.
- Ravindranatha, B.M.; Nandi, S.; Gupta, P.S.P. and Sarma, P.V. (2003). *In vitro* effects of different levels of commercially available PMSG on oocyte maturation. *Buffalo J*. 1:101-107.
- Sanchez-Partida, L.G.; Kelly, R.D.W.; Sumer, H et al (2011). Then Generation of Live Offspring from Vitrified Oocytes. *PLoS ONE*. 6 (6): e21597. doi: 10.1371/journal.pone.0021597.
- SAS . (2001). SAS/Stat. User's Guide Static's, Ver., 6.06 4th Ed. SAS Institute Inc. Cary, NC.
- Shayegh, M. and Barati, F (2011). The effects of pre and post vitrification taxol treatment on bovine immature oocyte. *Iranian Journal of Veterinary Research, Shiraz University*, Vol. 12, No. 3, Ser. No. 36.
- Suo, L.; Zhou, G. B.; Meng, Q.G.; Yan, C. L.; Fan, Z.Q. and Zhao, X. M., (2009). OPS vitrification of mouse immature oocytes before or after meiosis: the effect on cumulus cells maintenance and subsequent development. *Zygote*, 17:71-77.
- Sutton, M.L.; Cetica, P.D.; Beconi, M.T.; Kind, K.L.; Gilchrist, R.B. and Thompson, J.G. (2003). Influence of oocyte-secreted factors and culture duration on metabolic activity of bovine cumulus cell complexes. *Reproduction*; 126 .127–134.
- Tharasanit, T.; Colleoni, S.; Galli, C.; Colenbrander, B. and Stout, T.A., (2009). Protective effects of the cumulus-corona radiate complex during vitrification of horse oocytes. *Reproduction*, 137:391-401.
- Vincent, C.; Pickering, S.J. and Johnson, M.H. (1990). The hardening effect of dimethyl sulphoxide on the mouse zona pellucida requires the presence of an oocyte and is associated with a reduction in the number of cortical granules present. *J. Reprod. Fertil.*, 89: 253– 259.
- Yadav, R.C.; Sharma, A.; Garg, N. and Purohit, G.N. (2008). Survival of vitrified water buffalo cumulus-oocytes complexes and their subsequent development *in vitro*. *Bulg J Vet Med*. 11 (1):55-64.

- Zhang, J.; Nedambale, T.L. and Yang, M.; Li, J. (2009) Improved development of ovine matured oocyte following solid surface vitrification (SSV): effect of cumulus cells and cytoskeleton stabilizer. Anim Reprod Sci.110(1-2):46-55.
- Zhou, X.L.; AL Naib, A.; Sun, D.W. and Lonergan, P. (2010). Bovine oocyte vitrification using the Cryotop method: Effect of cumulus cells and vitrification protocol on survival and subsequent development. Cryobiology, 61: 66-72.
- Zhou, Y.; Xiangwei, F.;Guangbin, Z.;Baoyu, J.; Yi, F.; Yunpeng, H. and Shien, Z. (2014). An efficient method for the sanitary vitrification of bovine oocytes in straws. J Anim Sci Biotechnol. 2014; 5(1): 19.

التجميد بالتزجج والانضاج والاصحاب المعملی لبويضات الابقار مع أو بدون طبقات الخلايا الركامية

محمد الفاتح حماد*، شريف عبد الو نيس جبر*، إبراهيم طلعت الرطل** و ايمان عادل أمين*

* قسم الإنتاج الحيواني- كلية الزراعة - جامعة طنطا.

** قسم التكنولوجيا الحيوية - معهد بحوث الإنتاج الحيواني - مركز البحوث الزراعية.

تهدف هذه الدراسة الى تقييم تأثير وجود أو غياب طبقات الخلايا الركامية المحيطة بالبويضة على معدل الحيوية والانضاج والقدرة التنموية لبويضات الابقار بعد التجميد. تم جمع المبايض من المجازر وشفط البويضات من الحويصلات المرئية الموجودة على المبيض ثم فحصها. تم فقط تجميد البويضات الجيدة والبويضات المعراة كليا طبيعيا والبويضات المعراة كليا ميكانيكيا ثم عمل اسالة لها ثم انضاجها وخصابها و زراعتها معمليا لمدة ٧ ايام لحساب معدلات الحيوية والانضاج والاصحاب وانتاج البلاستوسيسست . وقد أظهرت النتائج أن العدد الكلى للبويضات المستردة ونسبة البويضات الطبيعية بعد التجميد بالتزجج كانت اعلى معنويا للبويضات الجيدة (٧٩.٠ و ٧٦.٠%) عن البويضات المعراة كليا طبيعيا (٥٦.٠ و ٥١.٤%). ايضا وجد ان معدل الاسترداد بعد التجميد بالتزجج ونسبة البويضات الطبيعية كانت اعلى معنويا للبويضات المعراة كليا ميكانيكيا (٦٧.٨ و ٦٣.٢%) عن البويضات المعراة كليا طبيعيا (٥٦.٠ و ٥١.٤%). وجد ان نسبة البويضات المضمحلة بعد التجميد بالتزجج لم تتأثر بشكل معنوى بنوع البويضات ولكن لوحظ ارتفاع نسبة تسرب المحتويات الخلوية من البويضات المضمحلة و انخفاض نسبة البويضات المضمحلة المنقسمة الى نصفين. وقد اسفرت البويضات الجيدة عن ارتفاع معدل الانضاج المعملی لها مع انخفاض نسبة البويضات المضمحلة (٦٠.٠ و ١٠.٠%) تليها البويضات المعراة كليا ميكانيكيا (٤٦.٩ و ٢٠.٥%) بينما اظهرت البويضات المعراة كليا طبيعيا ادنى نسب لها (٣٨.٩ و ٣٠.٤%) على التوالي . وكان معدل الانقسام للبويضات الجيدة اعلى معنويا عن كلا من البويضات المعراة كليا ميكانيكيا والبويضات المعراة كليا طبيعيا فكانت (٤٠.٠ مقابل ٢٦.٣ و ٢١.٣%). ولم يتأثر معدل انتاج الاجنة فى مرحلة الموريولا والبلاستوسيسست معنويا بنوع البويضات . وتشير هذه النتائج الى ان طبقات الخلايا الركامية المحيطة ببويضات الابقار المستردة من المبايض التى تم جمعها من المجازر تلعب ادوارا هامة جدا للمحافظة على حيوية البويضات اثناء التجميد ونجاح انضاجها وخصابها وتطورها معمليا حتى مرحلة البلاستوسيسست.