

EFFECT OF rbST ON PUBERTY AGE, HORMONES AND BLOOD METABOLITES IN FRIESIAN HEIFERS.

Hammad, M.E.*; M.A. Abu El-Hamd; Sh.A. Gabr*; Sh.M. Shamiah** and A. Alam***

* Anim. Prod. Dept., Fac. Agric., Tanta University.

** Anim. Prod. Res., Institute. Agric., Res. Center, Egypt.

ABSTRACT

To study the influence of rbST on puberty incidence of Friesian heifers in relation to progesterone (P4), growth hormone (GH), insulin like growth factor (IGF-I), glucose and urea-N in blood plasma, total of 16 Friesian heifers having 187.63 ± 2.56 kg live body weight and 8.23 ± 0.59 months of age were divided into two similar groups, 8 animals each. Animals in the first group (G1) were subcutaneously (s.c.) injected with saline (0.9% NaCl) while, animals in the second group (G2) were s.c. injected with 250 mg rbST 14 d-interval for five times pre-puberty. Feeding system and management were the same for both groups. Blood samples were collected to determine concentration of P4, GH, IGF-I, glucose and urea-N in blood plasma throughout the experimental period from 8.2 mo of age up to puberty. Results show insignificant differences in average P4 concentration during 3 wk pre-puberty, P4 peak pre-puberty and interval from P4 peak to puberty. The interval from 1st rbST injection to puberty was earlier (83.67 vs. 136.0 d, $P < 0.05$) and LBW was lighter (254.6 vs. 279.1 kg) in G2 than in G1. Concentration of P4 at puberty was not affected by treatment. Average age at puberty was 342 and 391d in G2 and G1, respectively. Overall concentration of P4 during the experimental period was nearly similar in G1 and G2, being 0.460 and 0.472 ng/ml, respectively. Concentration of IGF-I and GH was higher ($P < 0.05$) in G2 than in G1 as overall mean or pre- and at puberty. This increase was about 20 and 19% for IGF-I and 61 and 41% for GH pre- and at puberty, respectively. The differences in IGF-I and GH between pre- and at puberty for each group were not significant. Average concentration of IGF-I showed sharp increase post the 1st rbST injection by about 24.5%. Overall concentration of glucose during the experimental period increased ($P < 0.05$) in rbST group (85.9 mg/dl) as compared to the control group (79.1 mg/dl), although glucose concentration pre- and at puberty was not affected significantly by rbST treatment. Concentration of urea-N was lower ($P < 0.001$) in G2 than in G1 as overall mean (27.5 vs. 32.2 mg/dl). Also, concentration of urea-N reduced ($P < 0.05$) in G2 compared with G1 pre- and at puberty by 15 and 19%, respectively. Pre-pubertal urea-N concentration in heifers showed marked reduction by increasing number of injections, in particular post-1st injection. The strongest positive correlation was between concentration of GH and IGF-I ($r = 0.695$, $P < 0.001$).

In conclusion, rbST treatment at a level of 250 mg at 14- day interval for five times pre-puberty is strongly in relation to concentration of GH and IGF-I and in less extend to glucose and urea-N concentration to induce precocious puberty in Friesian heifers.

INTRODUCTION

Administration of exogenous somatotropin (ST) influences the somatotrophic axis, improves body weight gain and feed efficiency (Moseley *et al.*, 1992 and Rausch *et al.*, 2002), increases the number and size of ovarian follicles (Lucy, *et al.*, 1994) and growth of corpus luteum, and progesterone (P4) production (Gallo and Block, 1991). The somatotrophic axis is closely associated with pubertal development in heifers. Chandrashekar *et al.* (2004)

suggested a vital role for GH and IGF-I in the control of pituitary and gonadal functions in animals and humans. In this respect, Lucy (2000) found that ST and the IGF are important hormones for ovarian follicular growth. Receptors of ST and IGF are present in follicular cells. In addition, the granulosa and theca cells of the follicle are sites of IGF-I and IGF-II synthesis, respectively. Somatotropin can increase ovarian IGF-I synthesis. The IGF are important for follicular growth because both IGF-I and IGF-II are synergistic with gonadotropins for growth and differentiation of ovarian follicles.

The possible involvement of growth hormone (GH) in the regulation of ovarian follicular growth and development in mammals has been suggested by several observations. Therefore, circulating concentrations of GH increase rapidly during pubertal ages (Ojeda and Jameson, 1977). The effects of ST and IGF-I on the onset of puberty of heifers (195 kg) were studied by Simpson *et al.* (1991). They proposed that ST and IGF-I are important metabolic mediators involved in the initiation of puberty in heifers. Jones *et al.* (1991) determined changes in growth hormone (GH) and insulin-like growth factor I (IGF-I) before puberty in heifers. Frequency of GH release was greater at day -40 to -17 from puberty. Concentrations of IGF-I (measured every 2 wk) increased linearly ($P < 0.07$) from day -56 to 0 day from puberty. Mejia *et al.* (1999) suggested that enhanced pre-pubertal IGF-I levels in conjunction with increased pre-pubertal LH levels and pubertal LH pulse amplitude might be involved in the accelerated somatic maturation and in puberty advancement observed in heifers. Somatotropin increased circulating concentration of glucose an effect that may be attributed to increased hepatic gluconeogenesis and reduced uptake of glucose by adipose tissue (Hall *et al.* 1994). It was concluded that the changes in the follicular growth of rbST are not necessary induced by IGF-I but may be caused by changes in blood concentration of other hormones or metabolites, i.e. glucose (Oldick *et al.*, 1997). Consistent with the improved efficiency of amino acid in rbST-treated animals, a decrease in blood concentrations of urea nitrogen is consistently observed by Eisemann *et al.* (1986) and Eisemann *et al.* (1989).

Aim of the present study was to investigate the effect of rbST treatment in relation to some hormones (IGF-I and GH) and metabolites (glucose and urea-N) on puberty of Friesian heifers.

MATERIALS AND METHODS

This experiment was carried out at Sakha Experimental Station, belonging to Animal Production Research Institute (APRI), Agricultural Research Center (ARC), Ministry of Agriculture, during the period from September 2006 to Jun 2007.

Animals:

Total of 16 Friesian heifers having 187.6 ± 2.56 kg live body weight and 8.2 ± 0.59 months of age were divided into two similar groups, 8 animals in each. Animals in the first group were injected subcutaneously with 2 ml sterilized physiological saline (0.9% NaCl) at the same time; animals in the second group were s.c. injected with 250 mg rbST (Somatech® of Elilly) at

intervals of 14 days (five injections) from 252 days (8.2 mo) of age up to puberty. All heifers were free of any diseases with healthy appearance and they were housed in separated two groups under semi-open sheds, partially roofed with asbestos.

Feeding system and management:

Heifers in both groups were fed on equal amounts of diet containing the CFM, rice straw and fresh berseem (during winter season) or berseem hay (during summer season) according to the recommendation of the NRC (2001) allowances for growing dairy heifers based on live body weight. Diets were fed to both groups twice daily at 8 a.m. and 3 p.m, while fresh water was available all daytime.

Representative monthly samples of feedstuffs were chemically analyzed for CP, CF, EE, NFE and ash on DM basis according to the official methods of the A.O.A.C (1995). Chemical composition of CFM, rice straw, fresh berseem and berseem hay used in feeding heifers in both groups is shown in Table (1).

Table (1): Chemical analysis of different feed stuffs (on DM basis).

Item	Chemical composition (%)			
	CFM	Rice straw	Fresh berseem	Berseem hay
Dry matter, DM	90.22	89.24	15.26	88.23
Organic matter, OM	89.76	83.22	86.15	88.58
Crude protein, CP	16.04	1.59	14.71	14.41
Crude fiber, CF	10.96	37.21	24.9	24.67
Other extract, EE	4.91	1.47	2.90	6.04
Nitrogen free extract	56.38	42.85	43.64	43.16
Ash	10.24	16.78	13.85	11.42

Detection of puberty:

At the beginning of 9 mo of age, vasoectomized male was introduced to heifers of each group for 20 min three times daily at 6 and 12 a.m. and 6 p.m. to detect heifers exhibiting the 1st oestrous activity. The onset of 1st oestrus was used as an indicator for the onset of puberty. In addition, ovulatory activity of heifers at puberty was also indicated as P4 concentration exceeded one ng/ml for two consecutive sampling days (Jones *et al.*, 1991 and Simpson *et al.*, 1991) in blood plasma collected from heifers during the experimental period. Interval from rbST treatment to heifers reaching puberty was recorded.

Experimental procedures:

Blood sampling:

Blood samples were collected from the jugular vein of each animals in both groups starting before rbST treatment and 3-4 day-interval throughout an experimental period of 170 days. Blood samples were centrifuged at 3000 rpm for 10 minutes to separate blood plasma which stored at -200C until analysis.

Concentration of P4 was 3-4 day-interval determined at the beginning of the experiment up to puberty incidence. Throughout the experimental

period (170 days), concentrations of GH and IGF-I were weekly determined and concentration of glucose and urea-N were monthly determined in plasma.

Hormonal assay:

Direct radioimmunoassay technique (RIA) was performed for determination of plasma P4 concentration using antibody-coated tubes kit (Diagnosis systems, laboratories Texas, USA) according to the procedure outlined by the manufacture. According to the manufacture's information, the radioimmunoassay of progesterone is a competition assay. Sample and standards are incubated with I25labeled progesterone, as tracer, in antibody-coated tubes. After incubation the content of tubes is aspirated and bound radioactivity is measured. A calibration curve is established and unknown values are determined by interpolation from the curve. The standard curve of progesterone concentration ranged from 0 to 2.4 ng/ml. The intra-and inter-assay coefficient of variation were 5.4 and 9.1%, respectively.

Plasma GH and IGF-I were analyzed by radioimmunoassay. Measurements of GH and IGF-I and analyses were done according to the procedures previously described by Van Wyk (1983) and Daughaday and Rotwein (1989), respectively. The GH used for analysis was radioimmunoassay-grade. The inter- and intra-assay coefficients of variation for the GH analysis were 11.8 and 5.1%, respectively. Radioimmunoassay grade IGF-I was purchased from Gropep (Thebarton, SA, Australia). The inter- and intra-assay coefficients of variation for the IGF-I analysis were 10.6 and 7.7%, respectively.

Blood biochemical analysis:

Concentration of glucose (Trinder 1969) and urea-N (Patton and Crouch, 1977) in blood plasma were estimated using commercial kits (Diagnostic System Laboratories, Inc USA) and spectrophotometer.

Statistical analysis:

The obtained data were statistically analyzed according to Snedecor and Cochran (1982) and correlation coefficients were determined using SAS (1990)

RESULTS AND DISCUSSION

Progesterone profile:

Results presented in Table (2) show insignificant group differences in average P4 at 3 wk prior to puberty, P4 peak determined prior to puberty and interval from P4 peak to puberty incidence. However, the interval from first rbST injection to puberty in treated group was significantly ($P<0.05$) earlier than in control group (83.67 vs. 136.0 d), reflecting significantly ($P<0.05$) earlier incidence of puberty and insignificantly lighter weight of heifers in treated than in control group. Also, P4 concentration at puberty was not affected by treatment. The heavier weight at puberty of heifers in control group was attributed to the early puberty in treated group as compared to the controls.

Table (2): Effect of rbST treatment on concentration of P4 during pre-pubertal age and at puberty, and on P4 concentration, age and live body weight at puberty of Friesian heifers.

Item	Control	Treatment
During pre-pubertal stage:		
Average P4 (ng/ml) during 3 wk pre-puberty	0.644±0.039	0.585±0.036
P4 peak (ng/ml) prior to puberty	0.854±0.105	0.711±0.105
Interval from P4 peak to puberty (day)	14.0±1.86	11.5±1.86
Interval from first injection to puberty (day)	136.0±3.467 ^a	83.67±3.467 ^b
At puberty:		
P4 concentration (ng/ml)	0.360±0.043	0.414±0.043
Age (day)	391.0±3.5 ^a	341.7±3.5 ^b
Live body weight (kg)	279.1±18.6	254.6±12.9
Mean of P4 during the experimental period	0.460±0.03	0.472±0.04

a and b: Means having different superscripts within the same row are significantly different at (P<0.05).

Results illustrated in Figure (1) revealed that plasma P4 concentrations were almost higher in treated than in control group at all sampling pre-pubertal days. Average age of all heifers to attain puberty (progesterone greater than 1 ng/ml for two consecutive sampling days) was 342 and 391 days (11.4 and 13.0 mo of age) in treated and control group, respectively, being lower (P<0.05) in heifers injected with rbST than in those control group by 49 days (1.6 mo). It is of interest to note that overall concentration of P4 during the experimental period was nearly similar in both of treated and control group, being 0.472 and 0.460 ng/ml, respectively.

This proposed that rbST is important to induce the initiation of puberty in heifers. Similar results were obtained by Simpson *et al.* (1991) in heifers injected with growth hormone-releasing factor (GRF) and human serum albumin (HAS). Effects of rbST on puberty may be mediated by IGF-I, which prevents follicular atresia by hindering apoptosis of granulosa cells (Billig *et al.*, 1996). In this respect, Gallo and Block (1991) found that the combined effects of rbST and IGF-I have measurable effects on the ovarian function. Treatment with exogenous rbST increased the number of ovarian follicles with a distinct class of follicular diameter and increased concentration of P4 in the blood of treated cows. Results of Chadio *et al.* (2002) showed that rbST treatment did not significantly affect progesterone concentrations, although a tendency for higher levels in treated heifers (Gong *et al.*, 1991). Recently, Aboul-Wafa (2009) found an increase in P4 concentration in blood serum of ewes treated with rbST as compared to the controls.

In addition to its known metabolic effects, GH has been shown to have direct effects on function of ovarian granulosa cells in rats and pigs (Hsu and Hammond, 1987 and Adashi *et al.*, 1989). Some studies have shown that GH treatment in vivo can stimulate growth of small follicles in prepubertal pigs (Spicer *et al.*, 1990) and increase ovulation rates in cyclic gilts (Kirkwood *et al.*, 1988). The mechanism(s) for the increase in follicular growth and ovulation rate is (are) unclear. Furthermore, the action of r-bst may be indirectly mediated by increased IGF-I which stimulate ovarian function by acting synergistically with gonadotrophin to promote growth and steroidogenesis of ovarian cells (Lucy, 2000).




Fig. (1): Changes in plasma P4 concentration in treated and control group from rbST injection up to puberty.

Concentration of IGF-I and GH:

Data in Table (3) show that overall concentration of IGF-I and GH during the experimental period was significantly ($P < 0.001$) higher in treated than in control group (248.3 vs. 207.3 ng/ml for IGF-I and 29.9 vs. 21.8 ng/ml for GH). Also, concentration of IGF-I and GH pre- and at puberty was significantly ($P < 0.05$) higher in treated than in control group. This increase was about 20 and 19% for IGF-I and 61 and 41% for GH pre- and at puberty, respectively. It is of interest to note that concentration of IGF-I increased at puberty as compared to pre-pubertal values while, GH concentration showed different trend in both groups. However, the differences in IGF-I and GH between pre- and at puberty for each group were not significant.

Table (3): Average concentration of IGF-I and GH in blood plasma of treated and control heifers during the experimental period.

Item	Time	Control	Treatment	Sign.
IGF-I (ng/ml)	Pre-puberty ⁽¹⁾	189.3±5.4	226.7±5.4	*
	At puberty	204.3±12.0	242.7±12.0	*
	Overall mean	207.25±1.9	248.33±2.4	***
GH (ng/ml)	Pre-puberty	19.80±0.57	31.80±0.57	*
	At puberty	21.8±1.5	30.8±1.5	*
	Overall mean	21.80±0.52	29.98±0.32	***

• Significant at $P < 0.05$ *** Significant at $P < 0.001$. (1): average values of treated and control group were 8 and 10 wk, respectively.

Results illustrated in Figure (2) revealed that plasma IGF-I concentrations were almost higher in treated than in control group at all sampling pre-pubertal days. Average concentration of IGF-I showed sharp increase post the 1st injection in treated heifers, resulting in large differences

from that in control group by about 24.5%. These differences continued to be present after the subsequent injections, indicating higher pre-pubertal IGF-I concentrations in treated than in control group as affected by rbST injection.




Fig. (2): Change in plasma IGF-I concentration in heifers of treated and control groups throughout the experimental period.

It is worthy noting that GH concentration showed the same trend of change as IGF-I as affected by rbST injection, but the differences between treated and control group were reduced by increasing number of rbST injections from the 1st one (Fig. 3).




Fig. (3): Change in plasma GH concentration in treated and control heifers throughout the experimental period.

In similarity with the present study, Stelwagen *et al.* (1993) and Slaba *et al.* (1994) found a 3-7 fold-increase in plasma bST concentrations during the first three post-injection hours in cows treated with rbST. In the next 8 days, the rbST concentration in rbST treated cows was significantly higher than the controls. Plasma concentration of IGF-I increased nearly 2 folds as

early as 24 h following rbST treatment and then continued to increase by 48 h post 48 h injection (4 times higher than controls). From 48 h after rbST treatment, IGF-I concentrations remained at a plateau till day 11, then decreased slowly but still remained higher on day 14 than those in controls.

Administration of exogenous bovine GH stimulates growth in growing cattle, including pre-pubertal heifers (Sandles and Peel, 1987; McShane *et al.*, 1989 and Vestergaard *et al.*, 1993). Growth hormone exerts some of its action via IGF-I, and, consistent with this, the concentration of IGF-I in blood is increased by GH treatment (Crooker *et al.*, 1990). Also, Santos *et al.* (1999) found that level of IGF-I in blood serum of Zebu bulls significantly increase by rbST treatment. Similar trend was reported in Holstein Friesian bull calves (Holzers *et al.*, 2000).

The biological effects of IGF-I are modulated by IGF binding proteins (IGFBP), which have been characterized by legend blotting in calves (Skaar *et al.*, 1994).

Effects of exogenous ST on reproductive function have been studied during pre-pubertal ages in gilts (Bryan *et al.*, 1989) and heifers (Gong *et al.*, 1991, 1993 a&b). The somatotrophic axis is closely associated with pubertal development in heifers and Chandrashekar *et al.* (2004) suggested a vital role for GH and IGF-I in the control of pituitary and gonadal functions in animals and humans. In accordance with the present results, Schams *et al.* (1991) observed an increase of plasma bST at 7 d after treatment of cows with recombinant Methionyl bST (500 mg/14 d), and then decreased at 14 d. Also, concentration of IGF-I increased at 7 d and decreased by 14 d, but the concentration was still higher ($P<0.01$) at any point in treated than in the controls.

To identify metabolic hormones that serve as metabolic cues for onset of puberty, Jones *et al.* (1991) determined changes in GH, IGF-I and LH before puberty in heifers. Frequency of GH release was greater at day -40 than at day -17 from puberty in Angus heifers; however, in Braford and Charolais heifers frequency of GH release was greater at day -17 than at day -40. Concentrations of IGF-I (measured every 2 wk) increased linearly ($P<0.07$) from day -56 to 0 day from puberty in Angus but not in other breeds.

It has been suggested that gonadotropin concentrations are transiently increased prior to 22 wk of age in heifer calves (Schams *et al.* 1981 and Evans *et al.* 1992), but not all investigators have seen this early rise (Dodson *et al.*, 1988). Serum IGF-I levels increased from birth to 22 wk of age and then reached a plateau. Enhanced pre-pubertal IGF-I levels in conjunction with increased pre-pubertal LH levels and pubertal LH pulse amplitude might be involved in the accelerated somatic maturation and in puberty advancement observed in heifers (Mejia *et al.*, 1999). Plasma IGF-I concentrations were significantly higher in the rbST treated than in the control heifers at pre-pubertal stage. Similar trend was reported by Hodate *et al.* (1991), who observed that plasma IGF-I concentrations were significantly higher in the rbST treated cows than in control ones for 14 days after the treatment. Treatment with rbST significantly increased IGF-I concentration by 36.7% (121.2 and 77.6 ng/dl for rbST and control group, respectively, Gallo

and Block, 1990). Tripp *et al.* (1998) found that in heifers and steers, serum ST and IGF-I concentrations increased ($P<0.05$) by ST administration.

Lucy, (2000) found that ST and the IGF are important hormones for ovarian follicular growth. Receptors of ST and IGF are present in follicular cells. In addition, the granulosa and theca cells of the follicle are sites of IGF-I and IGF-II synthesis, respectively. Somatotropin can increase ovarian IGF-I synthesis. The IGF are important for follicular growth because both IGF-I and IGF-II are synergistic with gonadotropins for growth and differentiation of ovarian follicles. In Holstein growing steers, serum IGF-I concentrations increased ($P<0.01$) by 151% from d 7 through 35 in rbST-treated animals (Schlegel *et al.*, 2006).

Concentration of glucose and urea-N:

Data in Table (4) show that overall concentration of glucose during the experimental period significantly ($P<0.05$) increased in rbST group (85.9 mg/dl) as compared to the control group (79.1 mg/dl), although glucose concentration pre- and at puberty was not affected significantly by rbST treatment, but it showed different trend of change pre- and at puberty in each group.

On the other hand, concentration of urea-N was significantly lower in treated than in control group as overall ($P<0.001$) or pre and at puberty ($P<0.05$). The reduction in urea-N concentration was about 15 and 19% pre- and at puberty, respectively.

As affected by rbST injection, glucose concentration (Fig. 4) showed marked increased up to post 2nd rbST injection in treated group versus marked reduction in control group. However, glucose concentration was higher at puberty in treated than control group, then it increased in control group and still stable in treated group to be similar in both groups at puberty of control group. Such trend suggested incidence of puberty in control group when glucose level reached the same level in treated group. Similar trend of change in glucose level was reported by Gallo and Block (1990), who found that rbst-treatment, resulted in sharp increase in glucose level post-rbst-injection of Friesian cows.

Table (4): Average concentration of glucose and urea-N pre- and at puberty in blood plasma of heifers in treated and control group.

Item	Time	Control group	Treatment group	Significance
Glucose (mg/dl)	Pre-puberty	75.21±3.92	86.23±3.92	NS
	At puberty	81.68±3.25	80.73±3.25	NS
	Overall mean	79.10±1.2	85.86±1.3	*
Urea-N (mg/dl)	Pre-puberty	33.23±1.08	28.33±1.08	*
	At puberty	31.70±1.54	25.53±1.54	*
	Overall mean	32.16±0.86	27.48±0.92	***

NS: Not significant * Significant at $P<0.05$ *** Significant at $P<0.001$




Fig. (4): Plasma glucose concentration before (0 day) and after rbST treatment in treated and control group.

In accordance with the present results, Stevens, *et al.* (1980) reported that glucose values in lactating Holstein cows were 65.2, 68.1, 65.1, and 68.6 at bST levels of 0, 10.3, 20.6 and 41.2 mg/dl, while pre-treatment value was 57.1. Also, Pocius and Herbein (1986) stated that glucose concentration was not affected in plasma of Holstein-Friesian cows in mid lactation injected with GH (50 IU/day) for 11 consecutive days. Furthermore, Morbeck *et al.* (1991) reported that glucose was not influenced by dose of rbST.

In contrast, Molento *et al.* (2002) found that bST induced insulin peripheral resistance and increased liver gluconeogenesis, or both. Also, Hall *et al.* (1994) stated that ST increased circulating concentration of glucose, an effect that may be attributed to increased hepatic gluconeogenesis and reduced uptake of glucose by adipose tissue.

Concerning the results illustrated in Figure (5), plasma urea-N concentrations were almost lower in treated than in control group at all sampling days pre- and post puberty. It is of interest to note that pre-pubertal urea-N concentration in heifers showed marked reduction by increasing number of injections, in particular post-1st injection, then urea-N concentration showed slight rise post-puberty. This reduction in urea-N concentration in treated heifers may be a reflection of decreased amino acid degradation and utilization of these amino acids for increased protein synthesis (Marcek *et al.* (1989), suggesting that treated animals utilized protein more effectively (Whitaker *et al.* 1989 and Hodate *et al.* 1991). Also, in ST treated ruminants, a more efficient use of absorbed amino acids was accompanied by a reduction in circulating urea-N concentration (Boisclair *et al.*, 1994), demonstrating that whole body oxidation of amino acids is reduced with ST treatment (Eisemann *et al.*, 1989).




Fig. (5): Change in plasma urea-N concentration in heifers of treated and control groups throughout the experimental period.

In agreement with the present results, treatment with rbST (640 mg rbST/14d) significantly decreased plasma urea-N concentrations in Holstein heifers (Whitaker *et al.* 1989 and Hodate *et al.* 1991). Also, Early *et al.* (1990) reported that concentrations of serum urea was lower ($P < 0.05$) in steers (initially 9 mo of age and 231 ± 18 kg) receiving daily injections of rbST (20.6 mg/d) for 112 d. Moreover, Marcek *et al.* (1989) found that blood urea-N concentration decreased in pregnant lactating Holstein cows treated with 430 mg/d of rbST as compared to control cows. On the other point of view, several authors observed insignificant effect of rbST treatment on urea-N in blood of cows (Schams *et al.*, 1991 and West *et al.*, 1991) and primiparous Holstein cows Morbeck *et al.* (1991).

Correlation coefficients between each of all parameters studied are presented in Table (5). Results show that GH showed the strongest positive correlation with concentration of IGF-I ($r = 0.695$, $P < 0.001$) and negatively correlated with urea-N ($r = 0.188$, $P < 0.05$). Schams *et al.* (1991) found that concentrations of IGF-I were positively correlated with changes in ST.

Such results may indicate the important role of exogenous ST on increasing concentration of IGF-I as a metabolic mediator involved in the initiation of puberty in heifers (Simpson *et al.*, 1991).

It is of interest to note that the correlation between concentration was positive between IGF-I and glucose ($r = 0.314$, $P < 0.001$) and negative with urea-N ($r = 0.314$, $P < 0.001$). The present results indicated poor positive correlation between P4 concentration and each of IGF-I, GH, and urea-N, while P4 concentration negatively correlated ($P < 0.01$) with glucose concentration.

In conclusion, rbST treatment at a level of 250 mg at 14-day interval for five times pre-puberty is strongly in relation to concentration of ST and IGF-I and in less extend to glucose and urea-N concentration to induce precocious puberty in Friesian heifers.

Table (5): Pearson correlation coefficients between different parameters studied in both treated and control groups

Item	IGF-I	GH	P4	Urea-N
GH	0.69560***			
P4	0.06175 ^{NS}	0.15344 ^{NS}		
Urea-N	-0.21743*	-0.18888*	-0.08411 ^{NS}	
Glucose	0.31459***	0.16080 ^{NS}	-0.27330**	-0.22963*

NS: Not significant * Significant at P<0.05 ** Significant at P<0.01 *** Significant at P<0.001.

REFERENCES

- A.O.A.C. (1995). Official Methods of Analysis 15th ed. Association of Official Analytical Chemists. Arlington, Virginia, USA.
- Aboul-Wafa, A. Heba (2009). Improving reproductive performance of sheep using rbST. M. Sc. Thesis, Fac. Agric. Mansoura Univ., Egypt.
- Adashi, E.Y.; C. Resnick; E.R. Hemsndcz; M.E. Svoboda; E. Hohf; D.R. Clemmm; P.K. Lund and I.J. Van Wyk (1989). Rodent studies on the potential relevance of insulin-like growth factor (IGF-1) to ovarian physiology. In: A. N. Hirschfield (Ed.) Growth Factor and the Ovary Plenum Press, New York.
- Billig, H.; S.Y. Shun; K. Eisenhauer and A.J.W. Hsueh (1996). Gonadal cells apoptosis, hormone regulated cell demise. Hum. Reprod. Uptat, 2:103-117.
- Boisclair, Y.R.; D.E. Bauman; A.W. Bell; F. Bunshed and N. Harkins (1994). Nutrient utilization and protein turn over in the hin limb of the cattle treated with bovine somatotripin. J. Nutr., 124: 664-673.
- Chadio, S.A.; J.Menegatos; J. Zervas; C. Goules; S. Deligeorgis and D. Kalogiannis (2002). Pituitary responsiveness to gonadotropin and thyrotropin releasing hormone in goat treated with recombinant bovine somatotropin. Small Ruminants, 46: 149-157.
- Chandrashekar, V.; D. Zaczek and A. Bartke (2004). The consequences of altered somatotropic system on reproduction. Biol. Reprod., 71:17–27.
- Crooker, B.A., M.A. McGuire, W.S. Cohick, M. Harkins, D.E. Bauman, and K. Sejrson (1990). Effect of dose of bovine somatotropin on nutrient utilization in growing dairy heifers. J. Nutr., 120:1256.
- Daughaday W.H. and P. Rotwein (1989). Insulin-like growth factor-I and II, peptide, messenger ribonucleic acid and gene structures, serum and tissue concentrations. Endocrine Rev., 10 (1): 68-91.
- Dodson, S.E.; B.J. Mcleod; W. Haresign; A.R. Peters, and G.E. Lamming, (1988). Endocrine changes from birth to puberty in the heifer. J. Reprod. Fertil., 82: 527-538.
- Early, R.J., B.W. McBride, and R.O. Ball (1990). Growth and metabolism in somatotropin-treated steers: I. Growth, serum chemistry and carcass weights. J. Anim. Sci., 68:4134–4143.
- Eisemann, J.H.; A.C. Hammond; B.H. Reynolds; S.N. McCutcheon; H.F. Tyrrell and C.L. Halaanab (1986). Effect of bovine growth hormone administration on metabolism of growing Herford heifers: Protein and lipid metabolism and plasma concentration of metabolites and hormones. J. Nutr., 116: 2504-2515.

- Eisemann, J.H.; A.C. Hammond; T.S. Rumsey and B.H. Baumman (1989). Nitrogen and protein metabolism and metabolites in plasma and urine of beef steers treated with somatotropin. *J. Anim. Sci.*, 67: 105-115.
- Evans, A.C.O.; G.P. Adams and N.C. Rawlings (1992). Effects of naloxone on circulating gonadotropin concentrations in pre-pubertal heifers. *J. Reprod. Fertil.*, 96:847-855.
- Gallo, G.F. and E. Block (1990). Effect of recombinant bovine somatotropin on nutritional status and liver function of lactation dairy cows. *J. Dairy Sci.*, Vol. (11): 3276.
- Gallo, G.F. and E. Block. (1991). Effects of recombinant bovine somatotropin on hypophyseal and ovarian functions of lactating dairy cows. *Can. J. Anim. Sci.*, 71:343–353.
- Gong, J.G.; T.A. Bramley and R. Webb. (1991). The effect of recombinant bovine somatotropin on ovarian function in heifers: Follicular populations and peripheral hormones. *Biol. Reprod.*, 941-949.
- Gong, J.G.; T.A. Bramley and R. Webb. (1993a). The effect of recombinant bovine somatotropin on ovarian follicular growth and development in heifers. *J. Reprod. Fertil.* 97:247–254.
- Gong, J.G.; D. McBride; T.A. Bramley and R. Webb (1993b). Effects of recombinant bovine somatotrophin, insulin-like growth factor-I and insulin on the proliferation of bovine granulosa cells *in vitro*. *J. Endocrinol.*, 139:67–75.
- Hall, J.B., K.K. Schillo, B.P. Fitzgerald and N.W. Bradley (1994). Effects of recombinant bovine somatotropin and dietary energy intake on growth, secretion of luteinizing hormone, follicular development, and onset of puberty in beef heifers. *J. Anim. Sci.*, 72:709.
- Hodate, K.; A. Ozawa, T. Johke (1991). Effect of a prolonged release formulation of recombinant bovine somatotropin on plasma concentrations of hormones and metabolites and milk production in dairy cows. *Endocrinol. Jpn.*, 38 (5):527-532.
- Holzer, Z.; Y. Aharoni; A. Brosh; A. Orlov; J. Veenhuizen and T.R. Kasser (2000). The influence of recombinant somatotropin on dietary energy level-related growth of Holstein Friesian bull calves. *J. Anim. Sci.*, 78: 621-628.
- Hsu, C.J. and J.M. Hammond (1987). Concomitant effects of growth hormone on secretion of insulin-like growth factor-I and progesterone in cultured porcine granulosa cells. *Endocrinology*, 121:1343.
- Jones, E.J.; J.D. Armstrong and R.W. Harvey (1991). Changes in metabolites metabolic hormones, and luteinizing hormone before puberty in Angus, Braford, Charolais, and Simmental heifers. *J. Anim. Sci.* 69: 1607.
- Kirkwood, R.N.; P.A. Thacker; A.D. Gooneratne; B.L. Guedo and B. Laarveld (1988). The influence of exogenous growth hormone on ovulation rate in gilts. *Can J. Anim. Sci.*, 68:1097–103.
- Lucy, M.C. (2000). Regulation of ovarian follicular growth by somatotropin and insulin-like growth factors in cattle. *J. Dairy Sci.*, 83:1635–1647.
- Lucy, M.C.; J.C. Byatt; T.L. Curran; D.F. Curran and R.J. Collier (1994). Placental lactogen and somatotropin hormone binding to the corpus luteum and effects on the growth and functions of the ovary in heifers. *Biol. Reprod.*, 50:1136.

- Marcek, J.M.; W.J. Seaman and J.L. Nappier (1989). Effects of repeated high dose administration of recombinant bovine somatotropin in lactating dairy cows. *Vet. Hum. Toxicol.*, 31:455.
- McShane, T.M.; K.K. Scillo; J.A. Boling; N.W. Bradley and J.B. Hall (1989). Effects of recombinant-DNA derived somatotropin and dietary energy intake on development of beef heifers: I. Growth and puberty. *J. Anim. Sci.*, 67:2230–2236.
- Mejia, M.; A. Gonzalez-Iglesias; G.S. Diaz-Torga; P. Villafane; N. Formia; C. Libertun, D. Becu-Villalobos and I.M. Lacau-Mengido (1999). Effects of continuous ivermectin treatment from birth to puberty on growth and reproduction in dairy heifers. *J. Anim. Sci.*, 77:1329-1334.
- Molento, C.F.M.; E. Block; R.I. Cue and D. Peticlerc (2002). Effects of insulin, recombinant bovine somatotropin, and their interaction on insulin-like growth factor-I secretion and milk production in dairy cows. *J. Dairy Sci.*, 85:738–747.
- Morbeck, D.E.; J.H. Britt and B.T. McDaniel (1991). Relationships among milk yield, metabolism, and reproductive performance of primiparous Holstein cows treated with somatotropin. *J. Dairy Sci.*, 74:2153–2164.
- Moseley, W.M.; J.B. Paulissen; M.C. Goodwin; G.R. Alaniz and W.H. Clafin (1992). Recombinant bovine somatotropin improves growth performance in finishing beef steers. *J. Anim. Sci.*, 70:412–425.
- NRC (2001). *Nutrient Requirement of Dairy Cattle*. (7th Ed), National Academy Press. Washington. D.C.
- Ojeda S.R. and H.E. Jameson (1977). Developmental patterns of plasma and pituitary growth hormone in the female rat. *Endocrinology*, 100:881-889.
- Patton, C.J. and S.R. Crouch (1977). Determination of urea-N in blood. *Anal. Chem.*, 49: 464
- Pocius, P.A. and J.H. Herbein (1986). Effects of *in vivo* administration of growth hormone on milk production and *in vitro* hepatic metabolism in dairy cattle. *J. Dairy Sci.*, 69:713–720.
- Rausch, M.I.; M.W. Tripp; K.E. Govoni; W. Zang; W.J. Weber; B.A. Crooker; T.A. Hoagland and S. A. Zinn (2002). The influence of level of feeding on growth and serum insulin-like growth factor-1 and insulin-like growth factor binding proteins in growing beef cattle supplemented with somatotropin. *J. Anim. Sci.* 80:94–100.
- Sandles. L.D. and C.I. Peel (1987). Growth and carcass composition of pre-pubertal dairy heifers treated with bovine growth hormone. *Anim. Prod.*, 44: 21.
- Santos, R.L.; A.F.C. Riberio; A.C. Vasconcelos; J.L. Persquere; S.J. Coelho; R. Serakides and S.R. Reis (199). Effect of growth hormone and induced release in germ cell population and apoptosis in the bovine testis. *Vet. School and Institute of Biological Sci.* , Fedral Univ. Minas Gerais Belo Horizonte, Brazil, *Theriogenology*, 15: 975-984.
- SAS (1990). *SAS user's guide*. Statistical Analysis System. Institute, Inc. Cary., NC.
- Schams D.; E. Schallenberger; S. Gombe and H. Karg (1981). Endocrine patterns associated with puberty in male and female cattle. *J. Reprod. Fertil. Suppl.*, 30: 103-110.

- Schams, D.; F. Graf; B. Graule; M. Abele and S. Prokopp (1991). Hormonal changes during lactation in cows of three different breeds. *Livest. Prod. Sci.*, 27:285–296.
- Schlegel, M.L.; W.G. Bergen; A.L. Schroeder; M.J. Vande Haar and S.R. Rust (2006). Use of bovine somatotropin for increased skeletal and lean tissue growth of Holstein steers. *J. Anim. Sci.*, 84:1176–1187.
- Simpson, R.B.; J.D. Armstrong; R.W. Harvey; D.C. Miller; E.P. Heimer and R.M. Campbell (1991). Effect of active immunization against growth hormone-releasing factor on growth and onset of puberty in beef heifers. *J. Anim. Sci.*, 69:4914-4924.
- Skaar, T.C.; C.R. Baumrucker; D.R. Deaver and J.W. Blum (1994). Diet effects and ontogeny of alterations of circulating insulin-like growth factor binding proteins in newborn dairy calves. *J. Anim. Sci.*, 72: 421.
- Slaba, J.; P. Krejci; L.M. Huybrechts; E. Decuypere and H. Herrmann (1994). Plasma profiles of somatotropin and IGF-I in dairy cows following application of two preparations of recombinant bovine somatotropin in a sustained release vehicle. *Physiol. Res.*, 43 (1):37-43.
- Snedecor, G.W. and Cochran, W.G. (1982). *Statistical Methods*. 7th Ed. Iowa Univ. Press, Ames. Iowa, USA.
- Spicer, L.J.; J. Klindt; R. Maurer; F.C. Buonomo and S.E. Echemkamp. (1990). Effect of porcine somatotropin (pST) on number of granulosa cell LH/hCG receptor, oocyte viability, and concentration of progesterone (P) and insulin-like growth factor-I (IGF-1) in follicular fluid (FFL) of lean and obese gilts. *J. Anim. Sci.*, 68:410.
- Stelwagen, K.; D.G. Grieve; B.W. McBride and J.D. Rehman (1992). Growth and subsequent lactation in primigravid Holstein heifers after prepartum bovine somatotropin treatment. *J. Dairy Sci.*, 75:463–471.
- Stelwagen, K.; D.G. Grieve; J.S. Walton; J.L. Ball and B.W. McBride (1993). Effect of pre-partum bovine somatotropin in primigravid ewes on mammogenesis, milk production, and hormone concentrations. *J. Dairy Sci.*, 76:992-1001.
- Stevens, J.B.; J.F. Anderson; W.G. Olson and J.C. Schlatthauer (1980). Metabolic profile testing. Page 597, in *Bovine Medicine and Surgery*. H. E. Amstutz, ed. Am. Vet. Publ. Inc., Santa Barbara, CA, USA.
- Trinder, P. (1969). Determination of blood serum glucose. *Ann. Cl. Biochem.*, 6:24.
- Tripp, M.W.; T.A. Hoagland; G.E. Dahl, A.S. Kimrey and S.A. Zinn (1998). Methionine and somatotropin supplementation in growing beef cattle. *J. Anim. Sci.*, 76:1197–1203.
- Van Wyk J.J. (1983). Diagnosis of growth hormone deficiency and other causes of growth failure. *The Endocrine Society, Syllabus for the 35th Annual Postgraduate Assembly October*, 90-106.
- Vestergaard, M.; K. Sejrsen; J. Foldager; S. Kastrup and D.E. Bauman (1993). The effect of bovine growth hormone on growth, carcass composition, and meat quality of dairy heifers. *Acta Agric. Sci. and Sect. A: Anim. Sci.* 43:165–172.
- West, J.W.; B.G. Mullinix and T.G. Sandifer (1991). Effects of bovine somatotropin on physiologic responses of lactating Holstein and Jersey cows during hot, humid weather. *J. Dairy Sci.*, 74: 480.

Whitaker, D.A.; E.J. Smith and J.M. Kelly (1989). Milk production, weight changes and blood biochemical measurements in dairy cattle receiving recombinant bovine Somatotropin. Vet. Record, 124:83- 86.

تأثير مستحضر هرمون النمو علي عمر البلوغ وهرمونات ومواد تمثيلية في الدم في عجلات الفريزيان

محمد الفاتح حماد*، محمد عوض أبو الحمد**، شريف عبد الوهيد جبر*، شريف مغاوري شامية** و أحمد علام*
* قسم الإنتاج الحيواني- كلية الزراعة - جامعة طنطا.
** معهد بحوث الإنتاج الحيواني - مركز البحوث الزراعية - مصر.

لتقييم هرمون البروجسترون وهرمون النمو وهرمون IGF-1 والجلوكوز واليوريبا في بلازما الدم المتعلقة بحدوث البلوغ في العجلات الفريزيان المعاملة بمستحضر هرمون النمو rbST. واشتملت الدراسة علي ١٦ عجلة فريزيان وزنها ١٧٨,٦٣ وعمرها ٨,٢٣ شهر من العمر ووضعت في مجموعتين متشابهتين ٨ حيوانات لكل مجموعة. المجموعة الأولى حققت تحت الجلد بمحلول فسيولوجي (٠,٩% كلوريد الصوديوم)، بينما حيوانات المجموعة الثانية حققت تحت الجلد ٢٥٠ ملليجرام مستحضر هرمون النمو (rbST) كل ١٤ يوم خمس مرات قبل البلوغ. وبدأ الحقن من عمر حوالي ٨,٢٣ شهر وحتى ١٦ شهر من العمر. وكان نظام التغذية والرعاية متشابهة في المجموعتين. وجمعت عينات الدم لتقدير تركيز هرمون البروجسترون وهرمون النمو وهرمون IGF-1 والجلوكوز واليوريبا في بلازما الدم. وأظهرت النتائج عدم وجود اختلافات معنوية في متوسط تركيز البروجسترون خلال ٣ أسابيع قبل البلوغ وأعلى تركيز البروجسترون قبل البلوغ والفترة من أعلى تركيز البروجسترون قبل البلوغ وحتى البلوغ. كانت الفترة الفاصلة من أول حقنة حتى البلوغ أقصر (٨٤ يوم) في المجموعة المعاملة عن الكنترول (١٣٦ يوم). أنخفض وزن الجسم الحي في المجموعة المعاملة عن الكنترول (٢٥٧,٦ و ٢٧٩,١ علي الترتيب). كان متوسط عمر البلوغ ٣٤٢ و ٣٩١ في في المجموعة المعاملة والكنترول علي الترتيب. لم يختلف تركيز البروجسترون عند البلوغ معنويًا في المجموعتين. كان تركيز البروجسترون متساوي تقريبًا في المجموعتين خلال الفترة التجريبية. ارتفع تركيز هرموني IGF-1 وهرمون النمو قبل وأثناء البلوغ وخلال الفترة التجريبية في المجموعة المعاملة والكنترول وكانت الزيادة في حدود ٢٠ و ١٩% في IGF-1 و ٦١ و ٤١% قبل وإثناء البلوغ علي الترتيب. ارتفع معنويًا تركيز هرمون النمو و IGF-1 في المجموعة مقارنة بالكنترول في كل العينات قبل البلوغ. وقد أظهر تركيز IGF-1 زيادة حادة بعد الحقنة الأولى و كانت في حدود ٢٤,٥% بينما كانت الزيادة كبيره في هرمون النمو بعد الحقنة الثانية. ارتفع تركيز الجلوكوز معنويًا خلال الفترة التجريبية بينما لم يختلف تركيزه معنويًا قبل أو أثناء البلوغ. انخفض تركيز نيتروجين اليوريا في الدم معنويًا في المجموعة المعاملة قبل و أثناء البلوغ وكان معدل الأنخفاض ١٥ و ١٩% على التوالي. أظهر تركيز نيتروجين اليوريا قبل البلوغ انخفاضًا ملحوظًا بزيادة عدد مرات الحقن وخصوصًا بعد الحقنة الأولى. وكان هناك ارتباط موجب ($r=0.695$) بين تركيز هرمون النمو و IGF-1.

و قد أوضحت نتائج الدراسة المقدمه أن الحقن بمستحضر هرمون النمو (٢٥٠ ملليجرام) لخمس مرات بفترة بيئية ١٤ يوم قبل البلوغ كان له علاقة قوية لرفع مستوى هرمون النمو و IGF-1 وبدرجه اقل مع مستوى الجلوكوز واليوريبا في التبيكر بحدوث البلوغ المبكر للعجلات الفريزيان.