

EVALUATION OF THE GROWTH PERFORMANCE OF BROILER CHICKS FED ON PLANT DIETS SUPPLEMENTED WITH SOME FEED ADDITIVES.

Osman, A. M. A.¹; M. A. H. Soliman¹; H. H. Hassanein² and A. A. A. Abd EL-Warh²

1- Animal Prod. Dept., Fac. of Agric., Minia Univ., Minia

2- Animal Prod. Dept., Fac. of Agric., South Valley Univ., Qena

ABSTRACT

A total number of 360, unsexed one-day old Hubbard broiler chicks were randomly distributed into 12 groups with 3 replicates of 10 chicks each and used in a factorial experiment [3 diets x 4 feed additives (probiotics)]. Two control diets (one starter and another grower) were formulated from plant-origin ingredients to meet the major nutrient requirements of broiler chicks. Other two starter and two grower diets were formulated to have 2% less CP but adequate ME (low-CP diets) or less ME by 200 kcal/kg but adequate CP (low-ME diets).

Both starter and grower diets were supplemented with or without 0.1% Bio-Min (as a microbial growth promoter) or 0.03% Ronozyme (as enzymatic growth promoter) or 0.1% Super-Bio-Action (as a microbial and enzymatic growth promoter). Birds received the experimental diets up to 6 weeks of age. The criteria of response were the performance of chicks for weight gain and feed conversion, some carcass and digestive tract measurements, and economic efficiency.

The results revealed the following

- Broiler chicks fed on the low-ME diets achieved heaviest body weight at 6 weeks of age and body weight gain during the period from 3 to 6 and 0 – 6 weeks of age.
- Addition of probiotics into broiler chick diets improved ($P \leq 0.05$) the chicks, body weight gain, during starting period (0-3 weeks of age) and during the whole experimental period (0 to 6 weeks of age).
- Broiler chicks fed on the low ME diet consumed the greatest ($P \leq 0.05$) amount of feed during the growing period (3 to 6 weeks of age) and entire experimental period (0 to 6 weeks of age).
- Birds fed on the low-ME or low-CP diets exhibited a significantly ($P \leq 0.05$) inferior feed conversion during the starting period compared to their control counterparts, but this significant difference was disappeared during the growing and entire experimental period.
- Probiotics supplementation improved ($P \leq 0.05$) the feed conversion during starting, growing and entire experimental periods.
- Neither diet nor probiotics had significant effect on viability of chicks during the entire experimental period.
- Broiler chicks fed on the low-ME diets achieved a lower ($P \leq 0.05$) cost of feed/kg gain compared to those fed on the low-CP diets. However, selling price/bird and economic efficiency were the highest ($P \leq 0.05$) by feeding the low-ME diets. Economic efficiency was also improved by feed supplements used.
- Dietary treatments had insignificant effect on dressing % , while broiler chicks fed on the low-CP diets had the highest ($P \leq 0.05$) abdominal fat % at marketing age.
- Probiotics supplementation improved ($P < 0.01$) carcass weight, while dressing, abdominal fat and giblets percentages were not significantly influenced.
- Broiler chicks fed on the low-ME diets recorded the lowest ($P \leq 0.05$) relative length of small intestine, cecum, large intestine and relative weight.
- Probiotics had negligible effect on some digestive tract measurements.

INTRODUCTION

Feeding cost represents the major component of the cost of poultry production. Many attempts have been made to decrease the cost of feeding of different species of poultry. These attempts focused on replacing the more expensive and less available feedstuffs by cheaper and more abundant ones, and the use of more balanced diets supplemented with synthetic amino acids or growth promoters, and thus increasing the efficiency of feed utilization. Recently, many countries have begun to prevent the application of chemical antibiotics; as growth promoters, because of their side effect on both birds and humans. Therefore, most of poultry producers tended to use probiotics as safe alternative growth promoters.

Nowadays, there is a tendency to compose poultry diets of plant materials, mainly cereals and vegetable protein. Most of the feed ingredients contain less-or non-digested part (cellulose, xylose, arabinose,...) and some anti-nutritive factors, which inhibit feed utilization and birds performance. Part of this adverse effect can be overcome by dietary enzyme supplementation which has been shown to lower the viscosity of intestinal contents and thus improving digestibility of starch, protein and fat by broiler chicks fed wheat-containing diets (Annison and Choct, 1991; Bedford, 1995). Moreover, the beneficial effect of dietary supplementation with probiotics which contain *Lactobacillus acidophilus*, on the performance of broiler chicks may be mediated by a direct antagonistic effect against pathogenic micro-organisms, resulting in a decrease in their population (Hentges, 1983) or by stimulation of immunity (Umesb, 1999).

The present work was carried out to study the extent to which the growth performance, carcass traits and/or some digestive tract measurements in broiler chicks can be affected by dietary supplementation of plant-protein diets with some probiotics.

MATERIALS AND METHODS

This study was carried out at the Poultry Farm, Department of Animal Production, Faculty of Agriculture, South Valley University. A total number of 360, one-day old Hubbard broiler chicks were used in this study. Chicks were purchased from Qena Poultry Station and brooded in two-tier wire-floored battery, located in windowless house under same managerial conditions. Chicks were randomly distributed into 12 groups (3 diets X 4 feed additives) with 3 replicates of 10 chicks each. Average initial body weights of chicks in all treatments were almost similar. Two control diets (one starter and another grower) were formulated from plant-origin ingredients to meet the major nutrient requirements of broiler chicks, as recommended by the National Research Council (NRC, 1994). Other two starter and two grower diets were formulated to have 2% less protein, but adequate level of metabolizable energy (ME) and termed as low-crude protein diets (low CP) or less ME by about 200 Kcal/kg diet, but adequate level of CP and assigned as low-ME diets. All diets were formulated to have similar levels of lysine and sulphur

amino acids (Table 1). All chicks had a full access to feed and water during the experimental period.

Table (1) Composition of the experimental diets:

INGREDIENTS %	STARTER DIETS			GROWER DIETS		
	Contro	Low Protein	Low Energy	Control	Low Protein	Low Energy
Yellow corn	53.20	60.00	57.15	57.15	63.94	61.60
Soybean meal,44%CP	31.10	25.23	31.10	26.00	20.13	26.00
Corn gluten meal	8.00	8.00	7.50	8.00	8.00	7.30
Vegetable oil	3.83	2.59	----	4.74	3.50	1.02
Dicalcium phosphate	1.70	1.70	1.65	1.75	1.76	1.70
Limestone	1.50	1.60	1.50	1.50	1.60	1.52
Common salt	0.10	0.10	0.10	0.10	0.10	0.10
Vit&min.mix	0.25	0.25	0.25	0.25	0.25	0.25
Lysine	0.14	0.29	0.13	0.27	0.42	0.27
Methionine	0.18	0.24	0.18	0.24	0.30	0.24
Total	100,00	100,00	100,00	100,00	100,00	100,00
Calculated analysis						
Crude protein	23.01	21.00	23.08	21.09	19.09	21.05
Metabolizable energy; ME(kcal ME/ kg diet)	3110	3105	2902	3209	3203	3004
Calcium; %	1.00	1.03	1.00	1.00	1.00	1.00
Avail. P; %	0.45	0.45	0.45	0.45	0.45	0.45
Lysine; %	1.20	1.20	1.20	1.20	1.20	1.20
Methionine+cys; %	0.93	0.93	0.93	0.93	0.93	0.93

Each diet was supplemented with Vit. & Min. Mix : Each 2.5 kg contains, Vit. A 10 MIU; Vit. D₃ 2 MIU; Vit. E 10000 mg ; Vit. K₃ 1000 mg ; Vit. B₁ 1000 mg ; Vit. B₂ 5000 mg ; Vit. B₆ 1500 mg ; Biotin 50 mg ; BHT 10000 mg ; Pantothenic acid 10000 mg ; Folic acid 1000 mg ; Nicotinic acid 30000 mg ; Mn 60 g ; Zn 50 g ; Fe 30 g ; Cu 4 g ; I 3 g ; Se 0.1 g ; and Co 0.1 gm.

Both starter and grower diets were supplemented with or without Bio-Min (1 kg/ton feed); as a microbial growth promoter, which contain Enterococci faecium, fructo-oligosaccharide, piece of cell wall of bacteria and a mixture of algae or Ronozyme (300g/ton feed), as an enzymatic growth promoter containing glucanase and pectinase or Super Bio-Action (1 kg/ton feed), as a microbial and enzymatic growth promoter. The applied level of such probiotics and enzyme products was recommended by the manufacturer. The composition of Super Bio- Action is presented in Table 2.

Birds in each replicate were weighed at 21 and 42 days of age and the feed consumed was also recorded on a replicate group basis. Feed conversion (grams feed /gram of gain) was calculated for the different experimental periods. Mortality was monitored and recorded daily throughout the entire experimental period. Cost of one kilogram feed for different diets, cost of feed/kg gain and the cost of feed/bird were calculated on the basis of the prevailing price of feed ingredients in the local market. The absolute net return was calculated as selling price/bird minus cost of feed/bird. The relative economic returns for the experimental birds were calculated relative to their control counterparts.

At 6 weeks of age (end of the experiment), three birds from each treatment with an average body weight of their respective treatment were slaughtered. Eviscerated carcasses were individually weighed and dressing

percentage was calculated (weight of carcass + giblets + abdominal fat/pre-slaughter weight x 100).

Table (2) Composition of Super Bio – Action / kg of the product.

Items		
Saccharomyces cerevisiae	1.25	Trillion CFU
Bacillus subtilis	500	million CFU
Protease	125	Unit
Amylase	125	Unit
Cellulase	10	Unit
Lipase	5	Unit
Pure calcium	180	Mg
Active dry yeast	125	Mg
Malted Barley	50	Mg
Aspergillus oryzae	25	Mg
Cane molass	50	Mg
Fermented corn	200	Mg
Fermented corn extract	25	Mg
Oyster shells	500	Mg

Percentages of giblets (liver + gizzard + heart) and abdominal fat were also calculated relative to carcass weight. Some digestive tract measurements of the empty small and large intestines, duodenum and cecum were estimated relative to the starved body weight. The density of small and large intestine, duodenum and cecum were calculated as weight of part or organ (g) divided by its length in cm. Data were subjected to analysis of variance using General Linear Model described in SAS User's Guide (SAS, 1994). The differences among the means of different treatments were compared using Duncan's multiple range test (at 5% level) only when F-value was significant (Duncan, 1955).

RESULTS AND DISCUSSION

1- Body weight and body weight gain:-

1-1-Effect of nutrient level:-

Average weekly body weight and body weight gain of broiler chicks as affected by diet (energy and protein levels) and feed additives at different age intervals are presented in Table 3. At 6 weeks of age and during the period from 3 to 6 and 0 to 6 weeks of age, the highest ($P \leq 0.05$) body weight and body weight gain were attained by birds fed on the low-ME diets. However, those fed on the control or the low-CP diets achieved similar body weight and body weight gain during the same previously mentioned periods. The superiority in body weight and body weight gain achieved by birds fed on the low-ME diet represented about 29.4% and 50.8% and 30.5% over those of their control counterparts respectively, at marketing age (at 6 weeks of age). This superiority in body weight and body weight gain achieved by birds fed the low-ME diets was mainly related to their higher feed intake (Table, 4). In

agreement with the present results, by Babu *et al.* (1986) who indicated that the reduction of protein by 2% units in the finisher diet did not adversely affect body weight. On the other hand, the present results disagree with those reported by Attia (1999), who indicated that body weight of broiler chicks responded positively with the increase of ME in the diets.

1-2-Effect of feed additives:

The addition of probiotics into broiler chick diets improved ($P < 0.05$) body weight at 3 and 6 weeks of age. Also, probiotics supplementation improved ($P < 0.05$) body weight gain of broiler chicks during the starting period (0-3 weeks of age) and the whole experimental period (0-6 weeks of age), but their insignificant improving effects on body weight gain during the growing period (3 - 6 weeks of age) were insignificant compared to those of the control birds (without probiotics). The relative improvements in 6-week-old body weight and body weight gain as a result of the addition of probiotics into broiler chicks diets were 14.92%, 12.04%, and 11.73%, and, 15.47%, 12.57% and 12.17% for Ronozyme, Biomin and Super Bio-Action for these two measurements, respectively. The beneficial effects of probiotics on body weight were associated with improvements in feed conversion ratios (Table, 4). Noy and Sklan (1994) indicated that the beneficial effects of dietary enzyme supplementation are likely related to an increase in the rate of diet digestibility and production of simple sugars (from fiber degradation). Also, Sklan (2001) suggested that the young bird might be limited in the types and amounts of enzymes necessary to utilize the high-carbohydrate and vegetable-protein diets at an early age, thus improving the nutrient digestibility. The dietary supplements used in the present study; as previously mentioned, either have enzymatic activity (such as Ronozyme) or have the ability to synthesize certain enzymes which can enhance the digestibility of nutrients.

Moreover, Watkins and Kratzer (1984) and Haddadin *et al.* (1997) found that *Lactobacillus* implantation in broiler chicks improved their weight gain and feed efficiency. Similarly Haddadin *et al.* (1997) indicated that the most likely explanation for the enhanced weight gain is that selected strain of *L. acidophilus* colonizes the intestine of the chicks to varying degrees and once established, improves the bioavailability of essential nutrients such as the minerals, calcium and zinc. Also, the same organism causes a massive decline in the population of undesirable Gram-negative species like *E. coli*. Yeo and Kim (1997) showed that dietary probiotic suppressed the growth of harmful bacteria and increased the useful intestinal bacteria, which metabolize lactic acid and synthesize some vitamins. This effect along with a slight increase in feed intake may be responsible for the increased weight gain of birds fed such supplements under the condition of the present study.

2-Feed intake and feed conversion

2-1-Effect of nutrient level:-

Average cumulative feed intake and feed conversion ratio of broiler chicks as affected by diet and probiotics during starting (0-3 weeks of age) and growing (3-6 weeks of age) periods and the whole experimental period (0-6 weeks of age) are illustrated in Table, 4. During the starting period, broiler chicks consumed similar amounts of feed.

Table (3): Means of body weight gain of broiler chicks as affected by nutrient level and type of feed supplements during the experimental period.

TREATMENT	Body weight		Body weight gain		
	3 Age in wks	6 Age in wks	0-3 Age in wks	3-6 Age in wks	0-6 Age in wks
Nutrient level	NS	**	NS	**	**
Control	543.6	1296b	496.41	753.75b	1250b
Low-CP	537.5	1262b	491.58	741.25b	1216b
Low-ME	540.4	1677a	494.75	1136.75a	1632a
SE	8.169	24.82	0.013	65.77	99.25
Feed additives	**	**	**	NS	**
Control	485.1b	1287b	440.77c	822.88	1241b
Ronozyme	579.4a	1479a	533.66a	900.00	1433a
Biomin	548.9a	1442a	503.11b	893.33	1397a
Super Bio-Action	547.6a	1438a	499.44b	892.77	1392a
SE	9.432	28.66	8.201	10.22	28.75
Interaction	NS	NS	NS	NS	NS
Control-Without	475.0	1150	429.33	675.00	1104
Control-Ronozyme	571.7	1332	526.33	760.00	1287
Control-Biomin	573.3	1343	528.33	770.00	1298
Control-Superbio action	554.3	1358	501.66	810.00	1312
Low-C.P-Without	503.3	1160	458.00	723.33	1115
Low-C.P-Ronzyme	590.0	1357	543.33	766.66	1308
Low-C.P-Biomin	523.3	1252	477.00	728.33	1205
Low-C.P Superbio action	533.3	1280	488.00	746.66	1235
Low-M.E-Without	480.0	1550	435.00	1070.33	1506
Low-M.E-Ronozyme	576.7	1750	531.33	1173.33	1705
Low-M.E-Biomin	550.0	1732	504.00	1181.66	1685
Low-M.E-Superbio action	555.0	1677	50.66	1121.66	1630
SE	16.34	49.64	121.10	101.7	49.80

For each main factor, means within the same column bearing different letter (s) differ significantly ($P \leq 0.05$)

However, significantly higher ($P < 0.01$) feed intakes were recorded for broiler chicks fed on low-ME diets during both the growing and the whole experimental periods. The relative increases in feed intake of broiler chicks fed on the low-ME diets were estimated to be 46.70 and 33.46% during the growing period and the entire experimental period, respectively, compared to those of the control birds. These results are in full agreement with the results reported early by Arram (1984) and Sohair (1984) who indicated that, broiler chicks fed the low energy diets achieved the highest feed intake. Moreover, Sailer (1985) indicated that feed intake was not significantly affected with reducing protein level up to 6 weeks of age in broiler chicks. On the other hand, the present results disagree with those reported by Garcia Neto *et al.* (2000) who indicated that feed intake increased as protein content in the diet was decreased.

Diet had significant effect ($P \leq 0.05$) on feed conversion ratio only during the starting period; broiler chicks fed on the low CP or ME diets recorded inferior ($P < 0.05$) feed conversion ratio compared to those fed on the control diet at the previously mentioned period. From these results, it could be noticed that lowering the level of protein content or metabolizable energy in diets (by about 2% or 200 kcal ME/kg) less than the recommendations of NRC (1994)

had no deleterious effect on feed conversion ratio of broiler chicks reared in windowless houses. These results coincide with the results reported by Moran (1979), who found insignificant differences in cumulative feed conversion ratio from 0 to 7 weeks of age in broiler chicks fed isocaloric grower rations of 24, 22 and 20% CP for males and 24, 20 and 16% CP for females. Also, Dagher (1983) showed that, feed efficiency was not reduced by feeding either 15 or 12 % CP in the roaster diets, provided that these diets contain the recommended levels of methionine and lysine. Bougon and Menec (1986) fed broiler chicks on four diets containing 20.5, 21.7, 22.4 and 22.9 CP% from 1 to 45 days of age, found similar feed conversion ratios among all groups. Moreover, Stilborn *et al.* (1990) found that birds fed on diets containing 18 or 20% CP had similar feed conversion ratios. Daster and Golian (2003) indicated that broiler chicks fed on corn-soy diet containing 19% CP or higher and supplemented with appropriate amounts of indispensable amino acids had similar feed conversion ratio compared to those fed a diet with 23% CP.

2-2-Effect of feed additives:-

Probiotics had significant effect ($P \leq 0.05$) on feed intake of broiler chicks only during the first 3 weeks of age. Broiler chicks fed on the diets supplemented with Ronozyme consumed insignificantly higher ($P < 0.05$) amounts of feed compared to those fed on the control diets. The increase in feed intake during the first 3 weeks of age in response to the addition of Ronozyme into broiler chick diets may be due to that its composite enzymes reduced the residence time of digesta in the gastrointestinal tract coincided with an improved nutrient digestibility (Adams, 1995). Those results agree with these reported by Friesen *et al.* (1992), Esteve garcia *et al.* (1997) and Alam *et al.* (2003) who indicated that enzymes and probiotics can increase feed intake in broiler chicks.

On the other hand, addition of probiotics into broiler chicks diet improved ($P < 0.05$) feed conversion ratios during the starting, growing and whole experimental periods. The improvements in feed conversion ratios in response to the addition of Ronozyme, Biomin or Super Bio Action were 14.62%, 14.57% and 17%, respectively, during the entire experimental period. The beneficial effect of feed supplements (enzymes or probiotic) on feed conversion ratio may be due to a lower viscosity of intestinal contents and to an improvement in nutrients digestibility in the plant-protein diets (Annison and Choct, 1991; Bedford, 1995). According to Bedford (2001) much of performance response is thought to be related to the change in the intestinal microflora, rather than to a direct effect of enzyme on diet digestibility. Moreover, the beneficial effect of probiotic which contain *Lactobacillus acidophilus* on the performance of broiler chicks may be mediated by a direct antagonistic effect against specific groups of pathogenic micro-organisms, resulting in a decrease in their number (Hentges, 1983) or by increasing the immunity (Umesb, 1999). Noy and Sklan (1994) indicated that beneficial effects of enzyme addition to broiler diets are likely related to an increase in the rate of diet digestibility and production of simple-sugars (from fiber degradation). As a result of such improvements in the diet digestibility there is a significant change in the substrate quality and quantity available to the intestinal microflora in both the upper and lower gut. Hussein and El-Ashry

(1991) indicated that the improvement in feed conversion ratio observed by Lactobacillus supplementation was mainly due to the improvement in the gastrointestinal conditions against the pathogenic bacteria which invade the digestive tract.

Table (4): Means of feed intake and feed conversion ratio of broiler chicks as affected by nutrient level and type of feed supplement during the experimental period.

Treatments	Feed intake (g)			Feed conversion		
	Age (wks)			Age (wks)		
	0-3	3-6	0-6	0-3	3-6	0-6
Nutrient level	NS	**	**	*	NS	NS
Control	855.00	1374.42c	2142.7b	1.730b	1.862	1.723
Low C.P.	880.00	1435.41b	2174.1b	1.786a	1.925	1.803
Low M.E.	887.50	2016.26a	2859.7a	1.800a	1.850	1.756
SE	0.123	0.0158	40.13	0.015	0.029	0.038
Feed additive:	**	NS	NS	**	**	**
Control	846.66b	1584.11	2447.2	1.925a	2.096a	1.990a
Ronozyme	928.33a	1621.27	2441.3	1.739b	1.821b	1.699b
Biomin	875.00b	1639.28	2379.7	1.741b	1.815b	1.70b
Super-Bio-Action	846.66b	1590.13	2300.4	1.696b	1.785b	1.65b
SE	12.48	0.0152	46.34	0.0158	0.030	0.042
Interaction	NS	*	NS	NS	NS	NS
Control-without	806.66	1428.33b	2156.7	1.884	2.133	1.952
Control-Ronozyme	911.66	1376.83b	2166.7	1.731	1.819	1.684
Control-Biomin	905.00	1350.86a	2147.7	1.715	1.755	1.654
Control-Superb	796.66	1341.66c	2100.0	1.590	1.752	1.601
Low-C.P-Without	893.33	1616.66b	2333.3	1.955	2.183	2.123
Low-C.P-Ronozyme	930.00	1393.33b	2196.7	1.711	1.812	1.676
Low-C.P-Biomin	850.00	1398.33b	2063.3	1.782	1.840	1.712
Low-C.P-Superbio	846.66	1333.33c	2013.0	1.735	1.866	1.703
Low-M.E-Without	840.00	1707.33b	2850.7	1.935	1.983	1.900
Low- M.E -Rononzy	943.33	2093.66b	2960.7	1.775	1.832	1.740
Low- M.E -Biomin	870.00	2168.66b	2928.0	1.725	1.849	1.736
Low- M.E Superbio	896.66	2095.40b	2698.3	1.764	1.738	1.656
SE	98.32	101.7	80.26	0.124	0.142	0.073

For each main factor, means within the same column bearing different letter (s) differ significantly ($P < 0.05$)

3-Viability:

Neither diet nor probiotics had significant effect on viability of broiler chicks during the entire experimental period. These results coincide with those reported by Abd El-Fattah (2002) who illustrated that diet density had insignificant effect on viability rate of broiler chicks from one-day up to 6 weeks of age. Also, Alam *et al.* (2003) and Huang *et al.* (2004) found that probiotic had insignificant effect on the mortality rate of broiler chicks.

4-Cost of feed/kg gain and economic returns

4-1-Effect of nutrient level:

Cost of feed (LE) per Kg gain and economic returns of broiler chicks as affected by diets and probiotics at 6 weeks of age are illustrated in Table 5. Broiler chicks fed on the low-ME diets achieved lower ($P < 0.05$) cost of feed/kg gain than those fed on the low-CP diets. The highest feed cost/bird was achieved by birds fed on the low-ME diets, however, they attained the best

($P < 0.05$) economic return compared to those fed on the control or low-CP diets. The economic efficiency for birds fed the low-ME diets at marketing age were 34.4% and 40.56% greater than those fed on the control or low-CP diets, respectively. This could be attributed to the higher body weight gain (Table, 3) for broiler chicks fed the low-energy diets compared to those fed the control or low-CP diets. Also, broiler chicks fed on the control diets recorded insignificant better economic efficiency by about 4.4% compared to those fed the low-CP diets.

Table (5): Cost of feed (LE) /Kg gain and economic return of broiler chicks as affected by nutrient level and type of feed supplement

TREATMENT	PRICE OF DIETS (LE /Kg)		COST OF DIETS PER		PERICE / BIRD	ECONOMIC EFFICIENCY	
	Starter	Grower	Kg gain	Bird		Return	%
Nutrient level	---	--	*	**	**	**	--
Control	1.890	1.899	3.24ab	4.12b	9.38b	5.26b	100
Low-C.P	1.850	1.914	3.32a	4.08b	9.12b	5.03b	95.6
Low-M.E	1.770	1.960	3.05b	5.17a	12.24a	7.07a	134.4
SE	----	----	0.07	0.08	0.25	0.21	---
Feed additives:		---	**	NS	**	**	---
Control	1.838	1.850	3.58a	4.50	9.30b	4.81b	100
Ronozyme	1.853	1.861	3.08b	4.52	10.75a	6.23a	129.5
Biomin	1.904	1.907	3.16b	4.52	10.47a	5.94a	123.5
Superbio-action	1.856	1.864	2.99b	4.27	10.44a	6.17a	128.3
SE	---	----	0.027	0.23	0.56	0.33	---
Interaction	----	----	NS	NS	NS	NS	----
Control-Without	1.895	1.899	3.62	4.09	8.28	4.18	100
Control-Ronozy	1.910	1.914	3.17	4.14	9.64	5.50	131.5
Control-Biomin	1.961	1.960	3.19	4.21	9.74	5.53	132.3
Control-Superbi	1.914	1.913	2.97	4.01	9.83	5.81	139.3
Low-C.P-Without	1.850	1.860	3.87	4.33	8.35	4.02	96.2
Low-C.P-Ronzym	1.860	1.870	3.07	4.09	9.82	5.72	136.8
Low-C.P-Biomin	1.910	1.930	3.22	3.96	9.04	5.07	121.8
Low-C.PSuperbi	1.864	1.880	3.11	3.49	9.26	5.31	127.0
Low-M.E-Without	1.770	1.790	3.26	5.08	11.29	6.20	148.3
Low-M.E-Ronozy	1.790	1.800	3.01	5.32	12.78	7.46	178.5
Low-M.E-Biomin	1.840	1.850	3.06	5.40	12.64	7.23	173.0
Low-M.E-Superbio	1.790	1.800	2.88	4.84	12.23	7.37	176.3
SE	--	---	0.021	0.027	0.052	0.051	--

For each main factor, means within the same column bearing different letter (s) differ significantly ($P < 0.05$)

4-2-Effect of feed additives:-

Probiotics supplementation into broiler chicks diet decreased ($P < 0.05$) the cost of feed/kg gain and improved ($P < 0.05$) the economic return by about 29.5%, 23.5% and 28.3% due to supplemental Ronozyme, Biomin and Super Bio-Action respectively, as compared to the control diets. This could be attributed to the higher body weight gain (Table, 3) and better feed conversion ratio attained by broiler chicks fed the supplemented diets compared to those fed the control diets. These results are coincided with the results reported by Kamaran *et al.* (2002), who indicated that enzyme supplementation into broiler chick diets improved their economic efficiency.

Moreover, Hussein and El-Ashry (1991) found that dietary supplementation with *Lactobacillus* decreased the cost per kg live weight of chicks by 17.3% CP compared to a control group.

5-Carcass traits:

5-1-Effect of nutrient level:

Means of carcass traits in broiler chicks as affected by diet and probiotics are illustrated in Table 6. Broiler chicks fed the low-ME diets had higher ($P < 0.05$) carcass weight compared to those fed the control or low-CP diets. This could be attributed to the close relationship between fasting body weight and carcass weight. Krapoth (1987) and Saleh (1992) found a strong correlation coefficient which fluctuated from $r = 0.92$ to $r = 0.98$ between body weight and carcass weight. Diet had insignificant effect on dressing% where the values of dressing% ranged between 75.79% to 78.55%. Similar trend was noticed in giblets %. As it was expected, broiler chicks fed on the low-ME diets had lower ($P < 0.05$) abdominal fat % compared to those fed on the low-CP diets. Similar results were reported by Mahapatara *et al.* (1984). Diamabra and McCartney (1985) who found that male broilers fed 9 and 15 % CP diets had significantly more fat than those fed 12 and 18% protein-diets. Choudhary and Aggarwal (1987) reported a positive relationship between dietary protein level and carcass weight of broiler chicks, but dressing % was not significantly affected by early protein restriction. Fancher and Jensen (1989) observed higher abdominal fat deposition when broiler chicks were fed lower levels of dietary crude protein during the growing period. Similar results were reported by Bunchasak *et al.* (1996) during the starting period. Also, Hai and Blaha (1998) indicated that, the low protein diets had little effect on dressing %, but increased the abdominal fat content when offered to broiler chickens during the finishing period (from 21 to 42 days of age). Moreover, Barbour *et al.* (1999) found an inverse relationship between dietary ME content and abdominal fat pad size of broiler chicks raised from 0 to 7 weeks of age.

5-2-Effect of feed additives:-

Feed additives had highly significant ($P < 0.01$) effect on fasting body weight and carcass weight, while dressing, abdominal fat, liver, heart, gizzard and giblets percentages at 6 week of age were not significantly affected by the addition of probiotics. Broiler chicks fed on diets with supplementary Ronozyme or Super Bio-Action had significantly ($p < 0.05$) higher fasting body weight and carcass weight compared to the unsupplemented control diets. However, broiler chicks fed on diets with supplemental Biomin had an intermediate fasting body weight and carcass weight at 6 weeks of age. Dressing% was not significantly affected by the addition of probiotics which was within the normal range. Also, addition of probiotics into broiler chick diets had no detrimental effects on the accumulation of abdominal fat in the carcass. Similar trend was also observed in liver, heart, gizzard and giblets percentages. The effects of interaction (Diets x Probiotics) on carcass traits were not significant.

Also, the effects of probiotic on carcass traits were not dependant on the type of diet. The higher chicks carcass weight observed with the supplemented-diets was attributed to the improvement in fasting body weight.

It is well known that there was a strong relationship between fasting body weight and carcass weight (Krapoth, 1987; Saleh, 1992). The present results are in agreement with the findings reported by Aly (1993), Zanella *et al.* (1999), Panda *et al.* (2000) and Kamran *et al.* (2002), who indicated that probiotic had insignificant effects on dressing %. Also, similar insignificant effects of probiotic on carcass yield were reported by (Esteve-garcia *et al.* 1997; Biswas *et al.* 1999; Kumprechtova, 1999) abdominal fat % (Esteve-Garcia *et al.* 1997; Zanella *et al.* 1999; Loddi *et al.* 2000) or the weights of internal organs (Aly, 1993; Panda *et al.*, 2000).

6-Some digestive tract measurements :

6-1-Effect of nutrient level:

Means of weight, length and density of the digestive tract in broiler chicks as affected by diets and probiotics are presented in Table 7. Broiler chicks fed on the low-CP diets recorded the highest ($P<0.05$) relative weight and length of duodenum compared to those fed on the control or the low-ME diets. However, those fed low-ME diets achieved the lowest ($P<0.05$) relative length of small intestine and cecum and lowest relative weight and length of large intestine compared to those fed on the control or low-CP diets. The low-ME diets increased ($P<0.05$) the density of cecum. From these results it could be noticed that broiler chicks were capable to adapt themselves to the alterations in diet formulations by increasing or decreasing the relative weight and length or the density of different parts of the digestive tract. Moreover, changes in the digestive tract parameters were more pronounced by lowering the ME level rather than by decreasing the CP content of the diet. These results agree with the findings of Osman *et al.* (1996), who reported that the relative weight of digestive tract and cecum length were increased when Pekin ducklings were fed on a non-conventional diet, which had higher level of crude fiber. Also, Swatson *et al.* (2000) found that increasing the level of protein in the diet increased significantly the empty weight of ileum and ceca. Moreover, duodenal weight varied significantly with the change in dietary protein level. Abd El-Fattah (2002) indicated that the relative length of small intestine of broiler chicks was increased progressively with decreasing density of diet. He found also that the effect of lowering ME on the relative length of small intestine was more remarkable than the effect of lowering the crude protein content of the diet.

6-2-Effect of feed additives:-

Probiotics had significant effect ($P<0.05$) only on the relative length of small intestine and density of large intestine. Broiler chicks fed on the diets supplemented with Biomin or Super Bio-Action recorded lower ($P<0.05$) relative length of small intestine compared to those fed on the nonsupplemented control diets.

The density of large intestine was increased ($P<0.05$) by the addition of Ronozyme or Super Bio-Action into broiler chick diets. While the density of large intestine was unchanged by the supplementation of diets with Biomin. The relative weight and length of duodenum and cecum and the relative weight of small intestine were insignificantly decreased in response to feed supplements used.

Table (6): Means of carcass traits in broiler chicks as affected by nutrient level and feed supplements.

TREATMENTS	fasting body weight (g)	Carcass weight (g)	Dressing %	Abdomina fat%	Liver %	Gizzard %	Heart %	Giblet %
Nutrient level	**	**	NS	**	NS	*	NS	NS
Control	1309 b	929 b	75.79	1.70 b	2.24	2.11 a	0.55	4.92
Low-C.P	1255 b	889 b	77.03	2.94 a	2.29	2.13a	0.59	5.02
Low-M.E	1598 a	1176.6a	78.55	1.77 b	2.47	1.80 b	0.61	4.87
SE	31.93	26.57	1.06	0.216	0.13	0.10	0.029	0.17
Feed additives:	**	**	NS	NS	NS	NS	NS	NS
control	1282.2b	810.2b	76.04	2.20	2.52	2.05	0.60	5.20
Ronozyme	1471.1a	1048.8a	75.79	2.02	2.32	1.82	0.54	4.69
Biomin	1372ab	994.8ab	77.79	2.11	2.21	2.16	0.61	4.97
Super Bio-Action	1425.1a	1038.8a	77.58	2.20	2.28	2.02	0.60	4.89
SE	36.87	30.68	1.22	0.25	0.15	0.11	0.03	0.20
Interaction	NS	NS	NS	NS	NS	NS	NS	NS
Control-Without	1178.6	802.6	73.36	1.69	2.23	2.33	0.53	5.11
Control-Ronozy	1372.6	950.0	73.59	1.53	2.13	2.10	0.53	4.78
Control-Biomin	1322.6	964.0	77.77	1.41	2.13	2.13	0.56	4.82
Control-Superbi	1362.0	999.3	78.43	2.16	2.46	1.90	0.60	4.98
Low-C.P-Witho	1260.0	876.0	74.56	3.16	2.43	2.03	0.60	5.12
Low-C.P-Ronzy	1321.3	962.0	77.19	2.81	2.16	1.70	0.50	4.37
Low-C.P-Biomi	1184.0	832.0	75.44	2.99	2.30	2.46	0.63	5.37
Low-C.PSuperi	1258.0	886.0	75.42	2.76	2.26	2.33	0.63	2.21
Low-M.E-With	1408.0	1052.0	80.19	1.73	2.90	1.80	0.66	5.37
Low-M.E-Rono	1719.3	1234.6	76.59	1.72	2.66	1.66	0.60	4.91
Low-M.E-Biomin	1609.3	1188.6	78.53	1.93	2.20	1.90	0.63	4.71
Low-M.E-Supb	1655.3	1231.3	78.91	1.68	2.13	1.83	0.56	4.49
SE	63.86	53.14	2.12	0.43	0.26	0.20	0.05	0.35

For each main factor, means within the same column bearing different letter (s) differ significantly ($P \leq 0.05$)

Table (7): Means of empty weight%, length% and density (weight (g)/length (cm) of certain parts of digestive tract in broiler chicks as affected by nutrient level and feed supplements.

TREATMENT	Duodenum			Small intestine		
	Weight%	Length%	Density g/cm	Weight%	Length%	Density g/cm
Nutrient level	**	**	NS	NS	**	NS
Control	0.62b	2.14b	0.32	2.43	9.91a	0.24
Low-C.P	0.91a	2.45a	0.36	2.78	10.49a	0.26
Low-M.E	0.66b	2.00b	0.32	2.48	8.87b	0.26
SE	0.02	0.081	0.01	13.0	0.271	0.009
Feed additives	NS	NS	NS	NS	*	NS
Control	0.84	2.37	0.35	3.27	10.48a	0.25
Ronozyme	0.74	2.11	0.35	2.41	9.27ab	0.26
Biomin	0.73	2.22	0.32	2.41	9.90b	0.24
Super Bio-Action	0.69	2.08	0.31	2.53	9.38b	0.27
SE	0.05	0.93	0.01	15.0	0.314	0.01
Interaction	NS	NS	NS	NS	NS	NS
control-Without	0.83	2.50	0.35	2.58	11.45	0.22
control-Ronozy	0.63	2.12	0.33	2.47	9.45	0.27
control-Biomin	0.60	2.15	0.31	2.26	9.73	0.23
control-Superbi	0.55	1.77	0.30	2.43	9.04	0.27
Low-C.P-Without	0.85	2.45	0.34	2.79	10.48	0.26
Low-C.P-Ronzym	0.85	2.22	0.37	2.57	9.83	0.26
Low-C.P-Biomin	0.95	2.62	0.35	2.61	11.31	0.22
Low-C.PSuperbi	0.98	2.52	0.39	3.17	10.34	0.30
Low-M.E-Without	0.81	2.16	0.37	2.77	9.50	0.28
Low-M.E-Ronozy	0.69	2.00	0.34	2.20	8.56	0.26
Low-M.E-Biomin	0.63	1.88	0.32	2.36	8.64	0.27
Low-M.E-Superb	0.53	1.97	0.26	2.01	8.78	0.23
SE	0.08	0.16	0.02	26.0	0.54	0.01

For each main factor, means within the same column bearing different letter (s) differ significantly ($P \leq 0.05$)

Table 7 (continued)

TREATMENT	Cecum			Large intestine		
	Weight%	Length%	Density g/cm	Weight%	Length %	Density g/cm
Nutrient level	NS	**	*	*	**	NS
Control	0.46	2.63a	0.17 b	0.17a	0.59a	0.29
Low-C.P	0.45	2.53a	0.17 b	0.18a	0.65a	0.28
Low-M.E	0.42	2.17b	0.19 a	0.15b	0.49b	0.29
SE	0.02	0.081	0.006	0.01	0.03	0.01
Feed additives	NS	NS	NS	NS	NS	*
Control	0.48	2.63	0.18	0.16	0.62	0.26 b
Ronozyme	0.41	2.31	0.18	0.17	0.52	0.32 a
Biomin	0.43	2.42	0.17	0.16	0.56	0.26 b
Super-bio-action	0.28	2.43	0.17	0.16	0.58	0.31 a
SE	0.03	0.94	0.007	0.86	0.03	0.015
Interaction	NS	NS	NS	**	NS	*
Control-Without	0.53	2.86	0.18	0.17	0.64	0.27
Control-Ronozy	0.4	2.43	0.16	0.16	0.57	0.28
Control-Biomin	0.47	2.59	0.17	0.15	0.54	0.27
Control-Superbi	0.45	2.66	0.17	0.21	0.64	0.33
Low-C.P-Without	0.40	2.55	0.16	0.15	0.62	0.27
Low-C.P-Ronzym	0.40	2.35	0.17	0.15	0.52	0.28
Low-C.P-Biomin	0.46	2.63	0.17	0.21	0.70	0.30
Low-C.PSuperbi	0.48	2.62	0.18	0.19	0.63	0.31
Low-M.E-Without	0.51	2.47	0.20	0.16	0.59	0.27
Low-M.E-Ronozy	0.44	2.15	0.22	0.18	0.47	0.39
Low-M.E-Biomin	0.37	2.04	0.18	0.10	0.44	0.23
Low-M.E-Superb	0.35	2.01	0.17	0.15	0.48	0.29
SE	0.04	0.16	0.01	0.05	0.05	0.02

For each main factor, means within the same column bearing different letter (s) differ significantly ($P \leq 0.05$)

Also, the effect of interaction (Diets x probiotics) on the digestive tract measurements was insignificant. From these results, it could be noticed that the addition of such supplements into broiler chick diets had negligible effect on the measurements of digestive tract and their positive effects was not strong enough to induce changes in carcass yield. The present results are partially in agreement with those reported by Brenes *et al.* (1993) and Viveros *et al.* (1994) who found that enzyme addition decreased the digestive tract weight by 20 % when expressed as a percentage of live weight. Fuente *et al.* (1995) indicated that empty weight of the intestinal tract was reduced by 5 % by enzyme addition. They also showed that the empty weight of the intestinal tract was linearly related to digesta viscosity ($r = 0.64$) and this relationship was higher in broilers fed diets without enzyme ($r = 0.90$).

From these results, it could be concluded that the addition of probiotics into broiler chick diets improved fasting body weight and carcass weight with no adverse effects on carcass traits or quality.

REFERENCES

- Abd El-Fattah, M. A. (2002). Using some natural growth promoters in broiler diets. M.Sc. Thesis, Fac. Agric., Minia Univ.
- Adams, C. A. (1995). Effect of commercial enzyme products for wheat-based broiler diets. Technical information first issue December 4.
- Alam, M. J., M. A. R. Howlider, M. A. H. Pramanik and M. A. Haque. (2003). Effect of exogenous enzyme in diet on broiler performance. *International J. Poult. Sci.*, 2 (2): 168-173.
- Aly, A. A. S. (1993). Influence of enzyme preparation and growth promoters on broiler performance. Ph. D. Thesis, Fac. Agric. Cairo. Univ.
- Annisson, G., and M., Choct (1991). Anti nutritive activities of cereal non starch polysaccharides in broiler diets and strategies minimizing their effects. *World's Poult. Sci. J.* 47:232- 242.
- Arram, G. A. (1984). The biological and physiological values of different rations in relation to body composition. Ph. D. Thesis, Fac. Agric, Cairo Univ.
- Attia, A. I. (1999). Effect of dietary energy concentration with/without pronutrients on the performance of broiler chicks. *Egyptian Journal of Nutrition and feeds*. 2 (special issue): 665-676.
- Babu, M., V. Sundararasu and P. Kothandaraman (1986). Studies on energy and protein requirements for broiler chickens. *Nutr. Abstr. Rev.* 57: 697.
- Barbour, G.W.; M. T. Farran; M. G. Uwayjan and V. M. Ashkarian (1999). Effect of adding varying levels of oil in a corn- soybean meal diet on performance and carcass quality of broiler chickens. *Poult. Sci.*, (Suppl. 1) 78: 53 (Abstr. 235).
- Bedford, M. R. (1995). Mechanism of action and potential environmental benefits from the use of feed enzymes. *Anim. Feed. Sci. Technol.*, 53: 145-155.
- Bedford, M.R. (2001). Enzymes, antibiotic and intestinal micro-flora. *Feed Mic.* vol., 9 number 2.
- Biswas, T.; L. Mandal and S. K. Sarker (1999). Studies of enzymes supplementation and herbal preparation at different levels of energy on the performance of broilers. *J. Interacademic*, 3: 53-58.
- Bougon, M. and M. Menec (1986). A study similar to field conditions, of different feeds of broiler chickens. *Nutr. Abstr. Rev.* 57: 632.
- Brenes. A.; M. Smith; W. Guenter and R. R. Marquardt (1993). Effect of enzyme supplementation on the performance and digestive tract size of broiler chickens fed wheat and barley based diets. *Poult. Sci.*, 72: 1731-1739.
- Bunchasak, C.; K. Tanaka; S. Ohtani and C.M. Collado. (1996). Effect of Met+Cys supplementation to a low-protein diet on the growth performance and fat accumulation of broiler chicks at starter period. *Animal Science and Technology*, 67 (11): 956-966.
- Choudhary, M. K. and C. K. Aggarwal (1987). Effect of early protein restriction on body measurements and meat quality of broilers. *Indian Vet. J.*, 64 (5): 408-414.

- Daghir, N. J. (1983). Effect of lysine and methionine supplementation of low protein roaster diets fed after six weeks of age. *Poult. Sci.*, 62 :1572 – 1575.
- Daster, B. and A. Golian (2003). Pinpointing the lowest protein diet for young male broilers. (Nut. Gorgan, Iran Univ.).
- Diamabra, O. H. and M. G. McCartney (1985). The effects of low protein finisher diets on broiler males' performance and abdominal fat. *Poult. Sci.*, 64: 2013-2015.
- Duncan, D. B. (1955). Multiple range and multiple F tests. *Biometrics* 11 :1-42
- Esteve-garcia, E.; J. Brufau; A. Pérez -Vendrell; A. Miquel and K. Duven (1997). Bioefficacy of enzyme preparations containing β -glucanase and xylanase activities in broiler diets based on barley or wheat, in combination with flavomycin. *Poult. Sci.*, 76: 1728-1737.
- Fancher, B. I. and L. S. Jensen (1989). Influence on performance of three to six-week old broilers of varying dietary protein contents with supplementation of essential amino acid requirements. *Poult. Sci.*, 68: 113-123.
- Friesen O. D.; W. Guenter; R. R. Marquadt and B. A. Rotter (1992). The effect of enzyme supplementation on the apparent metabolizable energy and nutrient digestibilities of wheat, barley, oats, and rye for the young broiler chick. *Poult. Sci.*, 71: 1710-1721.
- Fuente, J. M.; P. Pérez de Ayala, A. Flores and M. J. Villamind (1998). Effect of storage time and dietary enzyme on the metabolisable energy and digest viscosity of barley-based diets. *Poult. Sci.*, 77 : 90-97.
- Garcia Neto, M., G. M Pesti and R.I. Bakalli (2000). Influence of dietary protein level on the broiler chicken's response to methionine and betanie supplements. *Poult. Sci.*, 79: 1478 – 1484.
- Haddadin, M. S. Y.; S. M. Abd El-Rahim; N. H. Odelailah and R. K. Robinson (1997). A proposed protocol for checking the suitability of *Lactobacillus acidophilus* cultures for use during feeding trials with chickens. *Tropical. Sci.*, 37: 16-20.
- Hai, D. T and J. Blaha. (1998). The effect of low-protein diets with supplementation of essential amino acids on broiler chicken performance. *Agric. Tropic. Subtropic.*, 31: 109- 116.
- Hentges, D. J. (1983). Role of the intestinal micro flora in host defense against infection. In *Human Intestinal Micro-flora in Health and Disease*, edited by Hentges, D. J., pp. 311-331, Academic Press, New York.
- Huang, M. K.; Y. J. Choi; R. Houde; J.-W. Lee; B. Lee and X. Zhao (2004). Effects of *Lactobacilli* and an acidophilic Fungus on the production performance and immune responses in broiler chickens. *Poult. Sci.*, 83:788–795.
- Hussein H. H. and M. A. El Ashry. (1991). Some studies on the beneficial effects of *Lactobacillus* concentrate supplementation on broiler performance . *Egypt. J. Anim. Prod.*, 28 (1): 85-93.
- Kamran, M, T. N. Pasha, A. Mahmud and Z. Ali (2002). Effect of commercial enzyme (Natugrain) supplementation on the nutritive value and inclusion rate of guar meal in broiler rations. *International J. Poult. Sci.*, 1 (6): 167-173.

- Krapoth, H. J. (1987). Untersuchungen zur Schlacht-körperbewertung und Fleischbeschaffenheit bei Broiler. Ph.D. Thesis Kiel University, Germany.
- Kumprechtova, D. (1999). Effect of continuous administration of selected probiotic preparations on chicken broiler performance and nitrogen output in droppings. Mendel University of Agriculture and Forestry Brno, 127p. Dissertation thesis. Text in Czech.
- Loddi M. M., E. Gonzales, T. S. Takita and A. A. Mendes (2000). Effect of the use of probiotic and antibiotic on performance yield and carcass quality of broilers. *Revista Brasileira de Zootecnia* 29 (4) : 1124 - 1131.
- Mahapatara, C. M., N. K. Pandey and S. S. Verma (1984). Effect of diet, strain and sex on the carcass yield and meat quality of broilers. *Ind. J. Poultry Sci.*, 19 (4) : 236-240.
- Moran, E. T. (1979). Carcass quality changes with the broiler chicken after dietary protein restriction during the growing phase and finishing period compensatory growth. *Poultry Sci.*, 58, 1257-1270.
- NRC; National Research Council (1994). Nutrient requirements of poultry. 9th revised edition (Washington, National Academy Press.).
- Noy, Y. and D. Sklan (1994). Digestion and absorption in the young chicks. *Poultry Sci.*, 73: 366-373.
- Osman, A. M. A.; M. A. H. Soliman and S. A. Abd El-Latif (1996). Evaluation of a non-conventional diet for growing Pekin ducklings. *Egypt. Poultry Sci.*, 16 (1): 365-378.
- Panda, A. K.; M. R. Reddy; S. V. Rao; M. V. L. N. Raju and N. K. Praharaj (2000). Growth, carcass characteristics, immune competence and response to *Escherichia coli* of broilers fed diets with various levels of probiotic. *Archiv für Geflügelkunde*, 64 (4): 152 - 156.
- Sailer, K. (1985). Nutzung kompensatorischer Wachstumseffekte bei der Produktion schwerer Mastbroiler. Ph. D. Thesis, Hohenheim University, Germany.
- Saleh, K. (1992). Comparative studies on the performance of some broiler hybrids under certain feeding systems. Ph. D. Thesis, Fac. Agric., Minia Univ..
- SAS. (1994). SAS / STAT User's Guide: Statistics. Ver. 6.04, fourth edition, SAS Institute Inc., NC, USA.
- Sklan, D. (2002). Development of the digestive tract of poultry. *World's Poultry Sci. J.*, 57 (4): 415-428.
- Sohair, A. I. A. (1984). Studies on the effect of feeding broiler on rations varying in calcium sources and level or containing different values of metabolizable energy and crude protein. Ph. D. Thesis. Fac. Agric. Cairo Univ.
- Stilborn, H. L., Amy L. Izat and W. Waldoup. (1990). Lysine and crude protein requirements of male broilers from 3 to 6 weeks of age. *Poultry Sci* 69: 131 (Abstr.).
- Swatson, H. K., R. M. Gous and P. A. Iji (2000). Biological performance and gastrointestinal development of broiler chicks fed diets varying in energy : protein ratio. *South African J. Anim. Sci.*, 30 (Supplement1).
- Umesb, P. C. P. (1999). Probiotics benefits. *Poultry International*, 38 (12) : 40 - 44.

- Viveros, A., A. Brenes; A. M. Bizarro and M. Castano (1994). Effect of enzyme supplementation of a diet based on barley, and autoclave treatment, on apparent digestibility, growth performance and gut morphology of broilers. Anim. Feed Sci. Technol. 48: 237 - 251.
- Watkins, B. A. and F. H. Kratzer (1984). Drinking water treatment with a commercial preparation of a concentrated Lactobacillus culture for broiler chickens. Poult. Sci., 63:1671-1673.
- Yeo. J. and K. I. Kim (1997). Effect of feeding diets containing an antibiotic, or yucca extract on growth and intestinal urease activity in broiler chicks. Poult. Sci., 76 : 381-385.
- Zanella. I, N. K. Sakomura, F. G. Silversides, A. Figueirido and M. Pack (1999). Effect of enzyme supplementation of broiler diets based on corn and soybeans. Poult. Sci., 78 : 561-568.

تقييم الأداء الإنتاجي لبداري اللحم المغذاة علي علائق نباتية المضاف إليها بعض الإضافات الغذائية

أحمد محمد أحمد عثمان¹ ، مصطفى علي حسن سليمان¹، حسام حسين محمد حسنين² و

أحمد أبو بكر عبد المنعم عبد الوارث²

¹ قسم الإنتاج الحيواني - كلية الزراعة - جامعة المنيا

² قسم الإنتاج الحيواني والدواجن- كلية الزراعة - جامعة جنوب الوادي

تم تقسيم عدد ٣٦٠ كتكوت تسمين "هبرد" غير مجنس عمر يوم عشوانيا إلى ١٢ معاملة بكل منها ٣ مكررات واحتوت كل منها علي ١٠ كتاكيت (٣ علائق X ٤ مستويات اضافات اعلاف (منشطات بيولوجية)). تم تكوين عليقتين مقارنة (واحدة بادنة والاخرى نامية) من مصادر نباتية لتقابل الاحتياجات الغذائية لبداري اللحم. اما العليقتين البادئ وايضا العليقتين النامي الباقية فقد كونتا ليحتويا على بروتين خام اقل بنسبة ٢% ولكن طاقة ممثلة كافية (عليقة منخفضة البروتين) وعليقة تحتوى على مقدار اقل فى الطاقة الممثلة لكل كجم عليقة بمقدار ٢٠٠ كيلو كالورى ولكن مستوى البروتين الخام فى الحدود المثلى (عليقة منخفضة الطاقة). تمت إضافة أو عدم إضافة ٠,١% بيومين (منشط بكتيري) أو ٠,٣% رونوزيم (منشط إنزيمي) أو ٠,١% سوبربيو-اكشن (منشط ميكروبي وإنزيمي معا). غذيت البدارى على علائق التجربة حتى الاسبوع السادس من العمر. وكانت القياسات للأداء الإنتاجي للكتاكيت معدل الزيادة فى الوزن وعدل التحويل الغذائى، كما تم ايضا تقدير قياسات الذبيحة والقناة الهضمية، هذا بالإضافة الى العائد الاقتصادى. وقد اوضحت النتائج الآتى:

- حققت الطيور المغذاة علي العليقة المنخفضة فى الطاقة أعلى وزن جسم عند عمر ٦ أسابيع وكذلك أعلى معدل زيادة وزن جسم خلال فترة النمو من ٣-٦ أسابيع وطول فترة التجربة من عمر صفر - ٦ أسابيع.
- أدت إضافة المنشطات البيولوجية إلى العلائق إلى تحسن ($P \leq 0,05$) فى معدل الزيادة فى وزن الجسم خلال فترة البادئ (٠ - ٣ أسابيع) وطوال فترة التجربة (٠ - ٦ أسابيع).
- أدت تغذية بدارى اللحم على العلائق منخفضة الطاقة إلى زيادة الغذاء المستهلك خلال فترة النامي (٤ - ٦ أسابيع) وطوال فترة التجربة (٠ - ٦ أسابيع).
- حققت الطيور المغذاة على العليقة منخفضة الطاقة أو البروتين الخام اقل كفاءة تحويل غذائى بالمقارنة بالكنترول خلال فترة البادئ (٠ - ٣ أسابيع) ولكن هذه الفروق المعنوية اختفت تماما خلال فترة النمو وطوال مدة التجربة.
- أدت إضافة المنشطات البيولوجية إلى تحسن ($P \leq 0,05$) فى كفاءة التحويل الغذائى خلال فترة البادئ والنامي وطوال فترة التجربة.
- لم يكن هناك أى تأثير معنوي للعلائق أو المنشطات البيولوجية على حيوية بدارى اللحم طوال فترة التجربة.
- أدت التغذية علي العليقة المنخفضة فى الطاقة الممثلة إلى انخفاض ($P \leq 0,05$) فى تكلفة الغذاء/ كجم زيادة فى وزن الطائر مقارنة بالتغذية علي العليقة المنخفضة فى البروتين. بينما تحقق أعلى سعر بيع للطائر

- والعائد الاقتصادي بالتغذية على العلائق منخفضة الطاقة. كما تحسن ايضا العائد الاقتصادي باستخدام الإضافات الغذائية.
- لم يكن للعليقة تأثير جوهري على نسبة التصافي لبدارى اللحم بينما حققت الطيور المغذاة علي العليقه المنخفضة البروتين النسبة الأعلى ($P \leq 0.05$) في الدهن البطني عند عمر التسويق.
 - أدت إضافة المنشطات البيولوجية إلى تحسن جوهري ($P \leq 0.05$) في وزن الذبيحة بينما لم يكن هناك تأثير معنوي علي نسبة دهن البطن والكبد والقلب والقونصة (الحوانج).
 - سجلت الطيور المغذاة علي العليقة المنخفضة الطاقة الممتلئة اقل ($P \leq 0.05$) طول في نسبتي الأمعاء الدقيقة والأعورين وكذلك اقل وزن وطول نسبي للأمعاء الغليظة .
 - كان لإضافة المنشطات البيولوجية تأثير طفيف علي بعض قياسات القناة الهضمية.