

CHEMICAL AND SENSORY EVALUATION OF CHICKPEA AND SOYBEAN AS AFFECTED BY SOAKING, GERMINATION AND COOKING

El-Bagoury, A.A.*; Badea, A. Bessar** and Hanan, Ahmed. ***

* Home Econ. Dep., Fac. Specific Education, Tanta Univ.

** Food Sci. and Techno. Dept., Fac. Agric., Kafr El-Sheikh, Tanta Univ., Egypt.

***Food Tech. Inst., Agriculture Research Center, Giza, Egypt

ABSTRACT

Chickpea (cooker chickpea) and soybean were soaked either in distilled water or in 0.5% sodium bicarbonate solution for 12 hr, germinated for 72 hr and boiling cooked. The best cooked products, were those soaked in distilled water or in sodium bicarbonate solution before cooking. On the other side, germination up to 2 days was superior to three days. Generally, germination process reduced cooking times for the two legumes than soaking. While soaking processes elevated hydration coefficients for the two legumes, germination process reduced them. Also hydration coefficients of cooked unsoaked legumes were higher than those soaked or germinated before cooking. Cooking media of dry legumes contained higher amounts from total solids, ash and protein than soaking only or cooking after soaking or germination. The contents of the soaked distilled water from total solids, ash and protein were less than those of soaked in 0.5 % NaHCO₃ solution for both legumes. In contrast, the cooking media of the two legumes those were previously soaked in distilled water contained a higher amounts of the three ones. Soaking and germination processes raised water – soluble solids, wherein cooking process reduced them in all cases: raw, soaked and germinated chickpea and soybean. Soaking, germination and cooking increased crude protein and ether extract of chickpea and soybean. Soaking and cooking decreased ash contents, while they increased by germination. On the other hand, crude fiber was decreased by soaking, germination and increased by cooking. Finally, carbohydrates were decreased as a result of all processes for both legumes. Soaking and germination processes elevated total nitrogen (TN), non protein nitrogen (NPN), protein nitrogen (PN) and free amino nitrogen (FAN). On the other side, cooking process elevated TN, PN and reduced NPN, TFAN of all samples. Both soaking in distilled water or in 0.5 % sodium bicarbonate solution resulted in similar losses (1 – 21 %) for major minerals except sodium that increased by soaking in 0.5% sodium bicarbonate solution. Germination process lowered the contents of Zn, Mn, Fe, Na, K and P, meanwhile raised the contents of Ca, Cu and Mg. On the other side, cooking of dry legumes resulted in losses in all minerals with different ratios. Also, cooking after soaking or germination resulted in losses of some minerals (Zn, Mn, Fe, Cu, Na, K, and P) and increased others (Ca and Mg).

Keywords: chickpea, soybean, soaking, germination, cooking, sensory evaluation, cooking time, hydration coefficient, soluble solids, chemical composition, nitrogenous constituents and minerals.

INTRODUCTION

Legumes are important source of proteins and other nutrients and are commonly used as food and fodder. Particularly in developing countries, legumes represent the major component of daily dietary foodstuffs along with

bread (Shalaby 2000). In Egypt, it could be stated that legumes represent the major Egyptian diets whatever the standard of living (Abd-Allah *et al.*1988).

Chickpea (*Cicer arietinum*, L.) is an important source of protein in several developing countries. Among the world's grain legumes, chickpea is the second to dry beans (*Phaseolus vulgaris*, L.) in cultivated area and the third in production to dry beans and dry peas (*Pisum sativum*, L.)(Singh *et al.*1991).

Soybeans play a major role in agriculture, commerce, industry and nutrition and for centuries have been an important source of dietary protein for millions of people of the Orient (Sutardi and Buckle 1985).

Soaking and germination of legumes are an ancient and popular practice in many parts of the world (Shalaby 2000). Preparation time of dry legumes has been shown to be markedly reduced by procedures that influence cell integrity, texture, water imbibition and heat transfer. These procedures employ the use of a soak treatment capable of softening the seed coat of dry legumes. Several investigators have found that addition of sodium bicarbonate to the soak solution can markedly reduce cooking time of dry legumes by allowing better water imbibition through softening of the seed coat (Kilgore and Sistrunk 1981). A quick cooking method for legumes includes dehulling, soaking, germination and precooking. These processes reduced the cooking time and improved the nutritional value of beans (El-Sahan and Youssef 1992). It was reported that germination reduced cooking times and improved the sensorial attributes of cooked legumes (Trugo *et al.*2000)

The aim of the present investigation was studying effects of soaking in distilled water, 0.5% sodium bicarbonate solution (12 hr) or germination up to 72 hr following by boiling cooking on the sensory qualities, hydration coefficients, cooking times and the proximate composition of chickpea (cooker chickpea) and soybean. The investigation was extended to study solubility of solids, some nitrogenous constituents and minerals content of both legumes.

MATERIALS AND METHODS

A- Materials

Two samples of dry legumes, chickpea (*Cicer arietinum*, L.) variety Giza 195 and soybean (*Glycin maxl-merr*) variety Giza 82 (crops of 2004) were obtained from Agriculture Research Center, Giza, Egypt. The dried seeds were cleaned, freed from broken seeds, dust and other foreign materials and stored in polythene bags at 5°C until used.

B- Methods

1- Technological methods

Soaking: The dried seeds were soaked either in distilled water or in 0.5% sodium bicarbonate solution (1:3 w/v) for 12 hr (over night) at room temperature ($25 \pm 2^\circ\text{C}$). The unimbided water was recovered and the soaked seeds were rinsed twice with distilled water.

Germination: The soaked, washed seeds were germinated on stainless steel screen baskets, which were covered with two layers of cheesecloth. The germination process was carried out at room temperature ($25 \pm 2^\circ\text{C}$) for 72 hr. During the germination period, the seeds were rinsed with distilled water every 12 hr.

Cooking: The soaked and soaked -germinated seeds were added to boiling distilled water (1:3 w/v) in 500 ml Pyrex glass beaker, covered with aluminium foil and heated to maintain boiling until cooking. The seeds were considered completely cooked when they attained a soft texture and were easily mashed between the fingers (Wanjekeche *et al.* 2003). Raw seed samples (control) were washed in distilled water and cooked in the same manner, using a seed to water ratio of 1:5 (w/v). The cooked seeds were drained, allowed to cool at room temperature and homogenized.

2- Hydration coefficients were calculated as percentage increases in the weight due to soaking, germination and cooking for the two legumes (Hulse *et al.* 1977).

3- Sensory evaluation of cooked seed samples was carried out according to the method described by Watts *et al.* (1989). When the samples were used for sensory evaluation, table salt (0.22% w/w) was added before cooking. The panel was composed of 20 judges, using a fully structured 9-point rating scale to evaluate color, texture, flavor and overall acceptability of the cooked seeds. Drinking water was available for rinsing. The obtained data were subjected to analysis of variance (ANOVA) using the general linear of the Statistical Analysis System (SAS 1988). Also, means comparisons were performed using Duncan's multiple Range Test (Steel and Torrie 1980).

4- Analytical methods The soaked, germinated and cooked seeds were dried in an electric oven at 50°C for 24 hrs, ground to pass through a 60-mesh sieve and stored in polythene bags at 4°C in a refrigerator until used. Also, raw dry seed samples were ground to pass through a 60-mesh sieve and stored as above. All samples were analyzed in triplicate and all results were calculated on dry weight basis.

Proximate chemical composition including moisture, crude protein (N x 6.25), ether extract, crude fiber and ash was determined according to A.O.A.C. (1990) procedures. Carbohydrate content was calculated by difference.

Non-protein nitrogen (NPN) was determined by extracting the samples with 10% trichloroacetic acid (TCA) according to the method of Paredes-Lopez and Harry (1989). Kjeldahl nitrogen was determined on a 10 ml aliquot. Protein nitrogen (PN) was calculated as follows: $\text{PN} = \text{TN} - \text{NPN}$ where TN is total nitrogen.

Total free amino nitrogen was estimated by means of the ninhydrin colorimetric method outlined by Iodic *et al.* (1966) as tyrosine equivalent and converted to nitrogen by multiplying by the factor 0.077 (nitrogen content of tyrosine).

Water-soluble solids were measured according to the procedure of Agosin *et al.* (1989). Two grams of dry sample were suspended in 18 ml of distilled water. The suspension was gently stirred for 30 min at 80°C . The residue

was recovered on Whatmann No 1 filter paper, thoroughly washed, dried and weighed. Soluble solids were calculated by difference.

Zinc, manganese, iron, calcium, copper, magnesium, sodium and Potassium were measured using Perkin-Elmer atomic absorption spectrophotometer, Model 3300 and following the procedure of Kirk and Sawyer (1991). Total phosphorus contents were determined colorimetrically using the method of Mohamed *et al.* (1986).

On the other side, the cooking media were collected, allowed to cool at room temperature and taken with soaking media to estimate total solids, ash and soluble proteins

RESULTS AND DISCUSSION

Germination process was carried out at room temperature ($25 \pm 2^\circ\text{C}$). Although in the first day, germination was not well for the two legumes (chickpea and soybean), germination process was improved in the second day. After three days, fermentative flavor and sticky touch was observed, especially soybean.

Sensory evaluation

The results of the sensory evaluation of cooked legumes are presented in Table 1. The statistical analysis revealed significant ($P \leq 0.01$) differences for all parameters for the two legumes unsoaked, soaked or germinated. The best products of the two legumes were those soaked in distilled water or in sodium bicarbonate solution before cooking, respectively. It was noted that texture recorded the 3rd order for the two legumes, those soaked previously in 0.5 % sodium bicarbonate solution. On the other hand, germination up to 2 days was superior to three days, and for this reason, germinated legumes up to three days were eliminated from the other analysis. Lee *et al.* (1983) reported that beans (soybean and black-eyed peas) soaked in 1 – 3 % NaHCO_3 had no equal quality to the control. El-Bagoury *et al.* (2000) stated that lentil soaked in distilled water before cooking was the most acceptable followed by soaking in 0.5 % sodium bicarbonate solution and finally the control (unsoaked).

Table 1. Sensory evaluation of cooked chickpea and soybean

The treatment	The constituent				
	Color	Texture	Flavor	Taste	Acceptability
Chickpea					
1) Raw (unsoaked)	6.95 a	7.60 a	6.43 a	6.50 a	6.87 a
2) Soaked in distilled water (12 hr)	7.13 a	7.41 a	6.90 a	7.19 a	7.15 a
3) Soaked in 0.5% NaHCO_3 (12 hr)	7.20 a	7.18 a	6.71 a	6.88 a	6.99 a
4) Germinated (2 days)	6.88 a	6.46 a	5.73 a	6.31 a	6.35 a
5) Germinated (3 days)	5.59 b	5.41 b	4.67 b	4.72 b	4.97 b
Soybean					
1) Raw (unsoaked)	6.06 b	6.56 a	5.79 c	5.16 c	5.89 c
2) Soaked in distilled water (12 hr)	6.86 a	6.42 a	6.50 a	6.75 a	6.63 a
3) Soaked in 0.5% NaHCO_3 (12 hr)	7.01 a	5.37 b	6.27 b	6.33 b	6.25 b
4) Germinated (2 days)	5.73 c	4.35 c	4.57 d	4.95 d	4.90 d
5) Germinated (3 days)	5.14 d	3.88 d	3.39 e	3.75 e	4.04 e

* Score of 9 points.

** Values in the same column with different letters are significantly different ($P \leq 0.05$) using Duncan's multiple range test

Cooking times and hydration coefficients

Table 2 shows effects of different processes on cooking times and hydration coefficients of chickpea and soybean. Generally, germination process reduced cooking times for the two legumes than both soaking in distilled water or in 0.5% sodium bicarbonate solution. Adequate cooking of unsoaked chickpea required 99 min at 100°C. Chickpea soaked in distilled water or in sodium bicarbonate solution required 52 and 38 min, respectively, while germinated chickpea required only 35 min. The corresponding values of soybean were 126, 87, 75 and 71 min, respectively. Moreover, germination process reduced hydration coefficients, while soaking processes elevated them for the two legumes. Also, hydration coefficients of cooked unsoaked legumes were higher than those soaked or germinated before cooking. Generally, it is clearly that, hydration of both legumes through soaking and some hydrolysis through germination had shortened cooking times. Generally, the obtained results agreed with those of Rockland *et al.* (1979), Wang *et al.* (1979), Silva *et al.* (1981), Lee *et al.* (1983), El- Bedawey *et al.* (1989), Singh *et al.* (1991) and El. Bagoury *et al.* (2000).

Table 2: Cooking time and hydration coefficient of soaked, germinated and cooked chickpea and soybean

The treatment	Cooking time (min)	Hydration coefficient %		
		Soaking	Germination	Cooking
Chickpea				
1) Raw (unsoaked)	99	-	-	141.8
2) Soaked in distilled water (12 hr)	52	132.9	-	7.8
3) Soaked in 0.5% NaHCO ₃ (12 hr)	38	131.7	-	8.9
4) Germinated (2 days)	35	132.9	- 4.5	13.4
Soybean				
1) Raw (unsoaked)	126	-	-	126.2
2) Soaked in distilled water (12 hr)	87	119.7	-	5.4
3) Soaked in 0.5% NaHCO ₃ (12 hr)	75	117.5	-	7.7
4) Germinated (2 days)	71	119.7	-4.9	11.1

Room temperature during soaking (12 hr) and germination period (2days) was 25 ± 2 °C.

$$\text{Hydration coefficient} = \frac{\text{Sample weight after treatment} - \text{sample weight before treatment}}{\text{Sample weight before treatment}} \times 100$$

Losses of total solids, ash and protein into soaking and cooking media

Total solids, ash and protein of the soaking and cooking media recovered from chickpea and soybean are illustrated in Table 3. The dry legumes lost as much as 9 – 10 % of total solids into cooking medium in comparison to soaking only or cooking after soaking or germination. Also, cooking media of dry legumes contained higher amounts of ash and protein. It was noted that, the contents of the soaked distilled water from total solids, ash and protein were less than those of soaked in 0.5 % NaHCO₃ solution for both legumes. In contrast, the cooking media of the two legumes those were previously soaked in distilled water contained higher amounts of the three ones. Finally, the sum losses from the three ones by soaking and germination into the cooking media were nearly to those of cooked dry legumes. The obtained results were confirmed with Wang *et al.* (1979).

Table 3: Total solids, ash and proteins of soaking and cooking media of chickpea and soybean(g/100g dry wt.).

The constituent The treatment	%Total solids		Ash %		Proteins%	
	Soaking	Cooking	Soaking	Cooking	Soaking	Cooking
Chickpea						
1) Raw (unsoaked)	-	9.306	-	1.270	-	1.534
2) Soaked in distilled water (12 hr)	1.314	2.851	0.427	0.585	0.164	0.644
3) Soaked in 0.5% NaHCO ₃ (12hr)	1.706	2.511	0.777	0.384	0.182	0.629
4) Germinated (2 days)	1.314	4.446	0.427	0.697	0.164	0.811
Soybean						
1) Raw (unsoaked)	-	10.262	-	2.120	-	1.395
2) Soaked in distilled water (12 hr)	4.013	4.381	0.885	0.960	0.437	0.552
3) Soaked in 0.5% NaHCO ₃ (12hr)	4.398	3.867	1.486	0.461	0.495	0.517
4) Germinated (2 days)	4.013	5.429	0.885	1.017	0.437	0.825

Water – soluble solids

Water – soluble solids of chickpea and soybean are 22.123% and 24.883 %, respectively (Table 4). Soaking processes in distilled water or in 0.5% NaHCO₃ raised these values by 45.513%, 55.594% and 32.874%, 37.765% for chickpea and soybean, respectively. Moreover, the germination process raised the solubility, in addition to soaking process, by 14.679% and 17.349%, respectively. On the other side, cooking process for the two legumes, reduced the water-soluble solids in all cases: raw, soaked and germinated legumes, wherein dry legumes (without processes except cooking) recorded the lest value in this regard.

Table 4: Water- soluble solids of raw, soaked, germinated and cooked chickpea and soybean (on dry wt basis)

The treatment	% Soluble solids	
	Chickpea	Soybean
1) Dry (unsoaked)		
(A) Control*	22.123	24.883
(B) Cooking	19.241	14.964
2) Soaked in distilled water (12 hr)		
(A) Control*	32.413	33.063
(B) Cooking	16.764	16.399
3) Soaked in 0.5% NaHCO₃ (12 hr)		
(A) Control*	34.422	34.280
(B) Cooking	17.938	18.892
4) Germinated (2 days)		
(A) Control*	37.171	38.799
(B) Cooking	20.908	21.941

* Control = without cooking

The increase in soluble solids with soaking may be due to hydration process, while it may be due hydrolyzing enzymes through germination process. On the other hand, the reduction in soluble solids by cooking may be due to draining into the cooking medium.

Proximate chemical composition

Results in Table 5 show that crude proteins of dried chickpea and soybean were 25.069 % and 39.044, respectively. As a result of soaking in distilled water or in 0.5 % sodium bicarbonate solution for 12 hr, crude protein was increased by 3.690 % and 0.574 % in chickpea and by 10.693 % and 8.708 % in soybean, respectively. Also, crude protein of germinated legumes increased and reached 26.806 % and 43.963% in chickpea and soybean, respectively. On the other hand, cooking caused additional increases in crude protein of dry, soaked and germinated legumes. Such increases may be reflecting the decreases in the other components, especially carbohydrates, as a result of hydrolysis and leaching out. The obtained results are in accordance with those reported by El- Shimi *et al.* (1984 and 1992); Abdella *et al.* (1986); Uzogara *et al.* (1991); El- Bagoury *et al.* (1999 and 2000); Uwaegbute *et al.* (2000) and WanJekeche *et al.* (2003). Some investigators have reported a decrease in the protein content of some legumes as a result of soaking, germination or cooking (Abd – Allah *et al.* 1988; Hamza and Youssef 1988; El- Bedawey *et al.* 1989; El – Sahan and Youssef 1992; Khalaf Allah 1995 and Zia – ur – Rehman *et al.* 2003).

It was found that (Table 5) soybean contained higher amount of ether extract (22.727 %) than chickpea (5.549%). Soaking processes in distilled water or in 0.5% NaHCO₃ augmented ether extract, since it reached 7.753 % and 8.011 % in chickpea; 23.882% and 24.557% in soybean, respectively. An additional increment was recorded in ether extract of the two legumes by germination. These results are in agreement with those found by Damir and Shekib (1989). On the contrast, Khalaf Allah (1995) did not find change in the lipid content of mung bean after soaking, but there was a slight decrease during germination. Concerning cooking, the cooked legumes, whatever unsoaked, soaked or germinated, contained higher amounts of ether extract.

These increases could be explained by solubility of solids and leaching into soaking, rinsing and cooking media. These results agree with the reports of El – Samra *et al.* (1986), Shehata *et al.* (1994) and El – Bagoury *et al.* (2000). As shown in the same Table, ash content of chickpea and soybean was 3.842 % and 6.482 %, respectively. Soaking in distilled water or in 0.5% NaHCO₃ resulted in decrease in ash content of the two legumes. However, germination process induced a slight increase in the ash content (3.312% and 5.894% for chickpea and soybean, respectively.) On the other side, cooking slightly decreased the ash content of the two legumes, regardless of the previous processes. The decreases in the ash content may be due to leaching some minerals into soaking and cooking media.

The obtained results are in the line of Abd-Allah *et al.* (1988); Uzogara *et al.* (1991); El – Sahan and Youssef (1992); Shehata *et al.* (1994); Khalaf Allah (1995); Salama and Ragab (1997); El – Bagoury *et al.* (1999 and 2000); Habiba (2002) and Wanjekeche *et al.* (2003). by 17.480% and 18.388 % in soybean, respectively. Also, germination process resulted in slight decreases. In contrast, cooking raised the fiber content and may be as a result of discarding other components to cooking medium. The obtained results are in agreement with Abdella *et al.* (1986); Shehata *et al.* (1994); Khalaf Allah (1995).

It was observed that, chickpea and soybean contained 60.124% and 26.684% of carbohydrate, respectively. Soaking in distilled water, in 0.5% NaHCO₃, germination and cooking processes reduced the quantities of carbohydrate. The decline in the carbohydrate contents may be related to dissolution into soaking, rinsing and cooking media. These results accorded with those reported by Abdella *et al.* (1986); El-Samra *et al.* (1986); Abd – Allah *et al.* (1988); Uzogara *et al.* (1991); Youssef and Abdel – Gawad (1992); Shehata *et al.* (1994); El – Bagoury *et al.* (1999 and 2000); Uwaegbute *et al.* (2000) and El – Adawy (2002).

Table 5: Proximate chemical composition of chickpea and soybean as a result of soaking, germination and cooking (on dry wt basis).

The constituent	Moisture %	Crude Protein (Nx6.25) %	Ether Extract %	Ash %	Crude Fiber%	Carbohydrate** %
The treatment						
Chickpea						
1) Dry (unsoaked)						
(A) Control*	8.067	25.069	5.549	3.842	5.416	60.124
(B) Cooking	65.581	25.238	7.905	2.984	5.663	58.210
2) Soaked in distilled water (12 hr)						
(A) Control*	60.724	25.994	7.753	2.995	5.232	58.026
(B) Cooking	67.521	26.244	8.428	2.749	5.487	57.092
3) Soaked in 0.5% NaHCO₃ (12 hr)						
(A) Control*	60.922	25.213	8.011	3.163	5.105	58.508
(B) Cooking	68.210	25.756	9.233	3.114	5.393	56.504
4) Germinated (2 days)						
(A) Control*	62.455	26.806	7.885	3.312	5.041	56.956
(B) Cooking	69.453	27.356	8.374	2.504	5.239	56.527
Soybean						
1) Dry (unsoaked)						
(A) Control*	7.144	39.044	22.727	6.482	5.063	26.684
(B) Cooking	63.612	46.350	27.623	5.782	5.205	15.040
2) Soaked in distilled water (12 hr)						
(A) Control*	61.704	43.219	23.882	5.755	4.178	22.966
(B) Cooking	67.046	45.331	28.113	4.621	4.389	17.546
3) Soaked in 0.5% NaHCO₃ (12 hr)						
(A) Control*	61.129	42.444	24.557	5.932	4.132	22.935
(B) Cooking	67.155	44.169	29.627	4.374	4.421	17.409
4) Germinated (2 days)						
(A) Control*	61.682	43.963	24.203	5.894	3.874	22.066
(B) Cooking	70.542	46.325	28.377	4.220	4.087	16.991

* Control = without cooking. ** Carbohydrate was calculated by difference

Some nitrogenous constituents

The changes in some nitrogenous constituents of chickpea and soybean as a result of soaking, germination and cooking are presented in Table 6. Total nitrogen (TN) of chickpea was 4.011 %, while protein nitrogen was 3.538% that equals 88.207 % of total nitrogen. Mean while, non-protein nitrogen (NPN) was 0.473 %. Soaking processes led to increments in TN, NPN and PN, whereas soaking in distilled water increased them higher than soaking in 0.5% sodium bicarbonate solution. Also, germination process increased these contents. On the other side, cooking process increased TN, PN and decreased NPN for all samples under the study. Also, free amino

nitrogen (TFAA) was increased by soaking and germination and decreased by cooking.

Table 6: Changes in some nitrogenous constituents of chickpea and soybean as a result of soaking, germination and cooking.

The treatment	The constituent	TN%	NPN%	PN%	TFAN
Chickpea					
1) Dry (unsoaked)					
	(A) Control*	4.011	0.473	3.538	4.807
	(B) Cooking	4.038	0.327	3.711	4.699
2) Soaked in distilled water (12 hr)					
	(A) Control*	4.159	0.600	3.559	15.445
	(B) Cooking	4.199	0.543	3.656	5.858
3) Soaked in 0.5% NaHCO₃ (12 hr)					
	(A) Control*	4.034	0.501	3.533	12.119
	(B) Cooking	4.121	0.472	3.649	0.352
4) Germinated (2 days)					
	(A) Control*	4.289	0.711	3.578	19.248
	(B) Cooking	4.377	0.489	3.888	1.532
Soybean					
1) Dry (unsoaked)					
	(A) Control*	6.247	0.260	5.987	4.612
	(B) Boiling cooking)	7.416	0.223	7.193	1.042
2) Soaked in distilled water (12 hr)					
	(A) Control*	6.915	0.482	6.433	9.323
	(B) Cooking	7.253	0.257	6.996	1.308
3) Soaked in 0.5% NaHCO₃ (12 hr)					
	(A) Control*	6.791	0.397	6.394	5.307
	(B) Cooking	7.067	0.194	6.873	1.381
4) Germinated (2 days)					
	(A) Control*	7.034	0.540	6.494	12.070
	(B) Cooking	7.412	0.396	7.016	1.015

TN = Total nitrogen

NPN = Non-protein nitrogen

PN = Protein nitrogen

TFAN = Total free amino nitrogen

TN, NPN and PN were calculated as gm/100 gm dry weight, Where TFAN was calculated as mg/gm dry weight

*Control = without cooking.

Protein nitrogen of soybean was 5.987 % that represents 95.838% of total nitrogen (6.247 %). The rest (4.162%) was non-protein nitrogen (0.260%). Soaking and germination increased TN, NPN and PN, whereas these constituents were higher when soybean was soaked in distilled water than in 0.5% NaHCO₃. Also, free amino nitrogen (4.612 mg/g) was increased by soaking and germination. Concerning cooking process, TN, PN increased, while NPN, TFAN decreased. It was observed that the changes in NPN were normally parallel with the changes in TFAN with soaking, germination and cooking. Increases in TN by soaking, germination and cooking may be due to high losses in other constituents of chickpea and soybean, especially

carbohydrates. The increments in NPN and TFAN by soaking and germination may be due to dissociation of some protein fraction by active proteases. On contrast, the decrements in NPN and TFAN by cooking might be due to leaching them into the cooking media. The obtained results are confirmed, by El – Mahdy and El-Sebaiy (1982, 1985); Akapapunam (1985); Abdel – Aal and Rahma (1986); Allam 1987 and Shehata *et al.* (1994).

Minerals content

Tables 7 and 8 review minerals content of chickpea and soybean as affected by soaking, germination and cooking processes. Both soaking in distilled water or in 0.5 % sodium bicarbonate solution resulted in similar losses (1 – 21 %) for major minerals except sodium, that increased by soaking in 0.5% sodium bicarbonate solution (157 – 485 %). Germination process lowered the contents of Zn, Mn, Fe, Na, K and P by 1 – 20 %, meanwhile raised the contents of Ca, Cu and Mg by 2 – 17 %. On the other side, cooking of dry legumes resulted in losses in all minerals with different ratios. Also, cooking after soaking in distilled water or in 0.5 % Sodium bicarbonate solution resulted in losses of some minerals (Zn, Mn, Fe, Cu, Na, K, and P) by 1 – 58 %. In the same time, Ca and Mg raised by cooking to 1 – 9 %. Also, cooking of germinated legumes came to the same result. Ca and Mg are probably bound or otherwise rendered less accessible for leaching into the cooking water than the other major minerals (Chung *et al.* 1981). Generally, the obtained different results are attributed to leaching of some minerals into soaking and cooking media by different ratios; the mineral itself and the different processes. The obtained results are in accordance with those obtained by Chung *et al.* (1981); Abd – Allah *et al.* (1988); Chitra *et al.* (1996) and Salama and Ragab (1997).

From the obtained results, It could be concluded that soaking and /or germination processes of dry legumes reduced cooking time especially germination, concentrated nutritional elements and analyzed complex compounds into simple components that make them more useful.

T7-8

REFERENCES

- A.O.A.C. (1990): Official Methods of Analysis. Association of Official Analytical Chemists 15 th, ed., Washington, DC, USA.
- Abd-Allah, M.A.; F.M. Abu-Salem and S.M.A. Ahmed (1988). Relation between specific technological treatments, nutritive value and the HCN content of some common Egyptian legumes. *Annals Agric. Sci., Fac. Agric., Ain Shams Univ., Cairo, Egypt* 33 (2): 1101-114.
- Abdel-Aal, M.H. and E.H. Rahma (1986). Changes in gross chemical composition with emphasis on lipid and protein fractions during germination of fenugreek seeds. *Food Chem.* 22:1-13.
- Abdella, H.A.; S.H. El-Samra; H.N. El-Banna; S.A. Lotfy; A.M. Bahay and A.I. Labib (1986). Effect of heat and processing on the nutritive value of broad beans and lentils. *Annals of Agric. Sci. Moshtohor* 24(2): 805 - 814.
- Agosin, E.; D.Diaz; R.Aravena and E.Yanez (1989). Chemical and nutritional characterization of lupine tempeh. *J.Food Sci.*54: 102-104,107.
- Akapapunam, M.A. (1985) Effects of blanching, soaking and cooking on the hydrogen cyanide yields, nitrogen, ash and minerals of lima beans (*Phaseolus lumatus*). *J. Food Sci.* 50:1191.
- Allam, M.H. (1987). Chemical composition and nutritional value of fenugreek seeds during germination. *Annals Agric. Sci., Fac. Agric., Ain Shams Univ., Cairo, Egypt* 32 (3): 1537-1550.
- Chitra, U.; U.Singh and P.V.Rao (1996). Phytic acid, in vitro protein digestibility, dietary fiber and minerals of pulses as influenced by processing methods. *Plant Foods Hum. Nutr.* 49(4): 307-316.
- Chung, S.Y.; C.V. Morr and J.J. Jen (1981): Effect of microwave and conventional cooking on the nutritive value of colossus peas (*Vigna unguiculata*) .*J. Food. Sci.* 46: 472 - 273.
- Damir, A.A. and L.A. Shekib (1989). Chemical and nutritional evaluation of germinated fenugreek, debittered lupine and fowl mekelle. *Alex. J. Agric. Res.* 34(3): 15-27.
- El-Adawy, T.A. (2002). Nutritional composition and antinutritional factors of chickpea (*Cicer arietinum, L.*) undergoing different cooking methods and germination. *Plant Foods Hum Nutr.* 57(1): 83 - 97.
- El-Bagoury, A.A.; H.E.Kassab and R.E. Zein (2000). Effect of soaking and cooking methods on organoleptic evaluation and some chemical constituents of mung bean and lentil. *J.Agric.Sci.Mansoura Univ.*25 (5): 2753-2765.
- El-Bagoury, A.A.; M.A. Salem and H.E. Kassab (1999). Effect of soaking and germination of some legumes on their chemical and nutritional values. *J. Agric. Sci., Mansoura Univ.* 24(10): 5731 - 5745
- El-Bedawey, A.A.; E.H. Rahma; T.A. EL-Adawy and M.A. Goma (1989). Improvement in nutritional quality of faba bean by soaking treatment. *Egypt. J. Food Sci.* 17(1-2): 137 – 151.

- El-Mahdy, A.R. and L.A. El-Sebaiy (1982). Effect of germination on the nitrogenous constituents, protein fractions, in vitro digestibility and antinutritional factors of fenugreek seeds (*Trigonella Foenum graecum*, L.). *Food Chemistry* 8: 253-262.
- El-Mahdy, A.R. and L. A. El-Sebaiy (1985). Proteolytic activity, amino acid composition and protein quality of germinating fenugreek seed (*Trigonella foenum graecum*, L.) *Food Chemistry* 18: 19-33.
- El-Sahan, M.A. and A.M.M. Youssef (1992). Influence of quick cooking process on microstructure, protein quality, ash and polyphenols of chickpea (*Cicer arietinum*). *Egypt .J. Food Sci.* 20(2): 459-470.
- El-Samra, S.H.; H.M. Abdella; N.H. El-Banna; S.A. Lotfy; A.M. Bahay and A.I. Labib (1986). Effect of heat and processing on the essential amino acids content, lysine bioavailability and trypsin inhibitor of soybean. *Annals of Agric. Sci., Moshtohor* 24(2): 795-803.
- El-Shimi, N.M.; A.A. Damir and M. Ragab (1984). Changes in some nutrients of fenugreek seeds during germination. *Food Chem.* 14:11-19.
- El-Shimi, N.M.; L.A. Sheikeb and S.M. Kotet (1992). Changes in nitrogenous constituents, phytate and nutritive value during the germination of blackeye peas (*Vigna unguiculate*) and kidney beans (*Phaseolous vulgaris*). 2nd *Alex. Conf. Fd. Sci. Tech.* 173-184.
- Habiba, R.A. (2002). Changes in anti-nutrients, protein solubility, digestibility, and HCl-extractability of ash and phosphorus in vegetable peas as affected by cooking methods. *Food Chemistry* 77:187-192.
- Hamza, M.A. and M.M. Youssef (1988). Effects of decortication, soaking and germination on amino acid composition, in-vitro digestibility and some antinutritional factors of faba bean (*Vicia Faba*, L.). *Alex. J. Agric. Res.* 33(1): 103-114.
- Hulse, J.H; K.O. Rachie and L.W. Billingsley (1977). Nutritional standards and methods of evaluation for food legume breeders. International Development Research Center. Ottawa, Canada.
- Iodice, A.A.; V. Leong and I.M. Weinstock (1966). Separation of cathepsins A and D of skeletal muscles. *Arc. Biochem. Biophys.* 117:477.
- Khalaf Allah, A.M. (1995). Changes in chemical constituents of mung beans (*Phaseolus aureus*) as affected by dehulling, soaking and germination. *Egypt. J. Appl. Sci.* 10(2): 56-67.
- Kilgore, S.M. and W.A. Sistrunk (1981). Effects of soaking and cooking upon selected B-vitamins and the quality of blackeyed peas. *J. Food Sci.*46: 909-911.
- Kirk, R.S. and R. Sawyer (1991). Pearson's Composition and Analysis of Foods: Ninth Ed. Longman Scientific and Technical.
- Lee, Y.C.; D.B. Shin and D.W. Shin (1983). Improved cooking methods for dry beans and their effects on quality of cooked products. *Korean J. Food Sci. Technol.* 15(3): 307 - 313 [C.f., *FSTA*, 17 (4) 3 j 99, 1985].
- Mohamed, A.; P.A.J. Perera and Y.S. Hafez (1986). New Chromophore for Phytic acid determination. *Cereal Chem.* 63 (6): 475-478.

- Paredes-Lopez, O. and G.I.Harry (1989). Changes in selected chemical and antinutritional components during tempeh preparation using fresh and hardened common beans. *J.Food Sci.*54 (4): 968-970.
- Rockland, L.B.; W.R. Wolf; D.M. Hahn and R. Young (1979). Estimation of zinc and copper in raw and cooked legumes. An interlaboratory study of atomic absorption and x-ray fluorescence spectroscopy. *J. Food Sci.* 44:1711-1719.
- Salama, A.M. and G.H. Ragab (1997). Comparison of conventional and microwave cooking of kidney beans and carrot in relation to chemical composition, nutritive value and sensory characteristics. 2nd Egyptian Conf. Home Econ. *Menoufia Univ. Fac. Home Econ.* : 214 - 226.
- SAS. (1988). Statistical Analysis System .In A. A. Ray (Ed.), *User's guide*. Cary, NC: SAS Institute, Inc.
- Shalaby, A.R. (2000). Changes in biogenic amines in mature and germinating legumes seeds and their behavior during cooking. *Nahrung* 44(1): 23 – 27.
- Shehata, M.I.; F.A. Salem; E. Abd-Elatif and S.M. Abou-El-Maati (1994). Effect of processing of legumes on chemical composition and nutritive value. *Egypt J. Agric. Res.* 72 (4): 1123-1135.
- Silva, C.A.B.; R.P. Bates and J.C. Deng (1981). Influence of soaking and cooking upon the softening and eating quality of black beans. (*Phaseolous vulgaris*). *J. Food Sci.* 46: 1716 - 2720.
- Singh, U.; N. Subrahmanyam and J. Kumar (1991). Cooking quality and nutritional attributes of some newly developed cultivars of chickpea (*Cicer arietnum*). *J. Sci. Food Agric.* 55: 37 – 46.
- Steel, R. G. and J. H. Torrie (1980). Principles and Procedures of statistics 2nd ed.; Mc Graw -Hill Book: New York.
- Sutardi, A. and K. A. Buckle (1985): Phytic acid changes in soybeans fermented by traditional inoculum and six strains of *Rhizopus oligosporus*. *J. Appl. Bact.* 58: 539- 543.
- Trugo, L.C.; Donangelo, C.M.; Trugo, N.M.F.and Bach Knudsen, K.E. (2000). Effect of heat treatment on nutritional quality of germinated legume seeds. *J.Agric.Food Chem.*48: 2082-2086.
- Uwaegbute, A.C.; C.U.Iroegbu and O.Eke (2000). Chemical and sensory evaluation of germinated cowpeas (*Vigna unguiculata*) and their products. *Food Chemistry* 68:141-146.
- Uzogara, S.G.; I.D. Morton and J. W. Daniel (1991). Thiamin, riboflavin and niacin retention in cooked cowpeas as affected by kanwa treatment. *J. Food Sci.* 56 (2): 592 - 593.
- Wang, H. L.; E. W. Swain; C. W. Hesseltine and H. D. Heath (1979). Hydration of whole soybean affects solids losses and cooking quality. *J. Food Sci.*44: 1510-1513.
- Wanjekeche, E.; V.Wakasa and J.G. Mureithi (2003). Effect of germination, alkaline and acid soaking and boiling on the nutritional value of mature and immature mucuna (*Mucuna pruriens*) beans. *Tropical and subtropical Agroecosy stems*: 183 -192

Watts, B.M.; Yamaki, G.L.; Jeffery, L.E. and Elias, L.G. (1989). Basic sensory methods for food evaluation, Ed.; The International Development Research Center Pub, Ottawa, Canada.

Youssef, M.K.E. and A.S. Abdel-Gawad (1992). Available carbohydrates and in-vitro starch digestibility of some dry legumes: Effect of soaking and different methods of processing. 2nd Alex. Conf. Fd. Sci. Tech.: 164-172.

Zia-ur-Rehman; M. Islam and W. H. Shah (2003). Effect of microwave and conventional cooking on insoluble dietary fiber components of vegetables. *Food Chemistry* 80:237-240.

تأثير معاملات النقع والانبات والطهي على التقييم الكيماوي والحسي للحمص وفول الصويا

عادل عبد الحميد الباجوري * - بديعة عبد الرحمن بيصار ** - حنان احمد ***
* قسم الاقتصاد المنزلي - كلية التربية النوعية بطنطا - جامعة طنطا
** قسم علوم وتكنولوجيا الأغذية - كلية الزراعة بكفر الشيخ - جامعة طنطا
*** معهد تكنولوجيا الأغذية - مركز البحوث الزراعية - وزارة الزراعة

تم نقع كل من الحمص (حمص الطبيخ) وفول الصويا في الماء المقطر أوفى محلول بيكربونات الصوديوم (٠,٥ %) لمدة ١٢ ساعة والانبات لمدة ٧٢ ساعة ثم إجراء الطهي بالغليان . وكانت افضل المنتجات المطهية هي تلك التي تم نقعها أولا في الماء المقطر تليها المنقوعة في محلول البيكربونات. وكان الانبات لمدة يومين افضل من الانبات لمدة ثلاثة أيام. وبصفة عامة خفضت عملية الانبات مدة الطهي عن النقع فقط. وكانت معاملات التثرب أعلى في حالة البقوليات المطهية والتي لم يتم نقعها أولا ثم المنقوعة بينما أدت عملية الانبات إلى خفض معامل التثرب. و احتوت أوساط الطهي للبقوليات الجافة على كميات اكبر من الجوامد الذائبة والرماد والبروتين بالمقارنة بالنقع فقط أو الطهي بعد النقع أو الانبات. وفي حين احتوى ماء النقع المقطر على كميات أقل من الجوامد الكلية والرماد والبروتين بالمقارنة بمحلول البيكربونات حدث النقيض في سائل طهي البقول التي تم نقعها أولا في الماء المقطر. وبينما أحدثت عمليتي النقع والانبات زيادة فالجوامد الذائبة في الماء فان عملية الطهي قد أدت إلى العكس. ولقد ارتفع محتوى الحمص وفول الصويا من البروتين الخام و المستخلص الأثيري كنتيجة لمعاملات النقع والانبات والطهي بينما انخفض الرماد كنتيجة للنقع والطهي ارتفع بالانبات. ومن ناحية أخرى نقصت الألياف الخام بالنقع والانبات وزادت بالطهي. وأخيرا فان الكربوهيدرات نقصت كنتيجة لكل المعاملات. ولقد رفعت معاملتي النقع والانبات النتروجين الكلي والنتروجين الغير بروتيني والنيوتروجين البروتيني والنتروجين الأميني الحر بينما أحدثت عملية الطهي زيادة في النتروجين الكلي والنتروجين البروتيني وخفض النتروجين الغير بروتيني والنتروجين الأميني الحر في كل العينات. وتسببت معاملة النقع في حدوث فقد بنسب من ١ إلى ٢٠ % في غالبية المعادن فيما عدا الصوديوم والذي زاد كنتيجة للنقع في محلول بيكربونات الصوديوم. وتسبب الانبات في خفض مستويات الزنك والمنجنيز والحديد والصوديوم والبوتاسيوم و زيادة محتويات الكالسيوم والنحاس والمغنيسيوم. وأدى الطهي للبقول الجافة إلى نقص في كل المعادن بينما تسببت معاملة الطهي بعد النقع أو الانبات في نقص بعض المعادن والى زيادة في البعض الآخر. وتبين من النتائج السابقة أن معاملة البقول الجافة بالنقع في الماء أو محلول مخفف من بيكربونات الصوديوم أو الانبات تؤدي إلى خفض زمن الطهي وتركيز العناصر الغذائية وتحليل العناصر المعقدة منها إلى صورة يمكن الاستفادة منها بدرجة اكبر.

Table 7. Effect of soaking, germination and cooking on some minerals of chickpea (mg/ 100g dry weight).

Treatment Element	Dry (unsoaked)		Soaked in distilled water (12hr)		Soaked in 0.5 %NaHCO ₃ (12hr)		Germinated (2 days)	
	Control	Cooking	Control	Cooking	Control	Cooking	Control	Cooking
Zn	3.154	2.824	2.749	2.402	2.667	2.334	2.358	2.187
Mn	1.634	1.518	1.455	1.211	1.412	1.186	1.290	1.132
Fe	17.626	16.093	17.314	16.042	17.270	15.814	17.055	15.353
Ca	145.381	139.632	125.280	137.133	115.563	118.545	128.298	154.433
Cu	1.263	1.160	1.024	0.895	1.010	0.887	1.163	0.954
Mg	136.671	123.440	121.681	124.273	118.340	120.060	125.889	142.991
Na	1185.245	1065.614	1118.010	898.894	5435.831	2243.720	977.441	823.868
K	3436.285	3071.834	3123.228	2170.988	2687.636	2491.669	2065.544	1979.765
P	491.000	456.800	466.800	445.800	457.100	449.400	455.701	432.902

Control =without cooking

Table 8. Effect of soaking, germination and cooking on some minerals of soybean (mg/ 100g dry weight).

Treatment Element	Dry (unsoaked)		Soaked in distilled water (12hr)		Soaked in 0.5 %NaHCO ₃ (12hr)		Germinated (2days)	
	Control	Cooking	Control	Cooking	Control	Cooking	Control	Cooking
Zn	2.970	2.601	2.622	2.338	2.498	2.253	2.368	2.122
Mn	2.774	2.511	2.458	2.067	2.365	2.034	2.191	1.934
Fe	22.212	20.527	20.920	18.706	20.593	17.795	19.511	17.475
Ca	174.141	149.905	144.585	154.843	139.155	145.141	148.715	177.801
Cu	1.685	1.646	1.453	1.356	1.592	1.560	1.695	1.471
Mg	144.305	138.427	131.021	141.872	129.862	134.179	135.755	152.801
Na	1139.694	979.432	983.238	869.228	2936.841	1000.939	897.622	758.592
K	3453.284	2815.847	3124.703	2650.740	3095.175	2241.262	2591.848	2181.895
P	698.500	657.200	685.401	663.902	671.700	639.002	671.100	601.978

Control =without cooking