

EFFECT OF SHEEP MANURE, PHOSPHORUS LEVELS AND CHEMICAL FERTILIZERS AS WELL AS BIOFERTILIZERS ON GROWTH, YIELD, NITRATE ACCUMULATION AND CHEMICAL COMPONENTS OF BROCCOLI (*Brassica oleracea* var *italica*) PLANTS

Farrag, Amal M.*; A. H. Hanafy-Ahmed ** and Hanaa F. Ahmed**

* Vegetable Department, Faculty of Agriculture, Cairo University.

**Plant Physiology Division, Plant Department, Faculty of Agriculture, Cairo University.

ABSTRACT

Two field experiments were carried out on broccoli (*Brassica oleracea* var *italica*) plants cv. Premium to study the effect of sheep manure at the rate 10 m³ (4.28 tons/fed.), 3 levels of phosphate fertilizer i.e. 0, 10 or 20 kg P₂O₅/fed. or the recommended rate of chemical fertilizers (40 N + 30 P₂O₅ + 50 K₂O kg/fed.) with three kinds of biofertilizers i.e. yeast, *Pseudomonas*, *Bacillus* (bacteria dissolving phosphate) inoculated with seedling or uninoculation on vegetative growth characters at head initiation and at side head initiation as well as days from transplanting until head initiation, yield and chemical components in leaves and heads. Applying sheep manure plus 20 kg P₂O₅/fed. followed by sheep manure plus 10 kg P₂O₅/fed. gave higher values of plant length, plant, stem, leaves fresh weights, number of leaves per plant, head, side heads weights per plant and total yield/fed. as well as the percentage of N, P, K, dry matter, total sugar, soluble phenols and total free amino acids concentration in leaves and heads comparing with adding recommended chemical fertilizers. Days from transplanting until head initiation not affected by fertilization. The result indicated that using sheep manure plus 10 kg P₂O₅ gave better values than using chemical fertilizers. On the other hand, ascorbic acid of heads was higher by adding sheep manure plus 10 kg P₂O₅ than values of sheep manure plus 20 kg P₂O₅ or chemical fertilizers. Nitrate content of heads were the highest with application chemical fertilizer. The three biofertilizers treatments increased vegetative growth parameters, total yield and its components as well as chemical components in the leaves and heads as comparing with uninoculation treatment. *Pseudomonas aeruginosa* resulted in the highest yield of heads in the first season also side head weight and total yield per fed. as well as total sugars and amino acids in leaves in both seasons. Yeast caused the higher values of heads yield per fed. and total free amino acids in heads, in the second season. On the other hand, nitrate content of heads were the highest with inoculated plants with yeast, while *Bacillus* caused the lowest values. *Bacillus* results in the highest total amino acids in heads in the first season and ascorbic acid content in heads in both season. The highest total yields were obtained of plants received sheep manure plus 20 kg P₂O₅/fed and inoculated with *pseudomonas* or *Bacillus* in the first and second seasons, respectively.

INTRODUCTION

Broccoli (*Brassica oleracea* L. var *italica* Plenck) is an important nutritional cole vegetable. It is high in vitamin A, ascorbic acid and is a good source of calcium, niacin and riboflavin (Decoteau, 2000).

Broccoli is a cool-season crucifer. It has about the same climatic requirements as cauliflower. It is well adapted to all areas when grown during the coolest months of year, although it is not as sensitive to hot weather. It is

harvested over a longer period of time than cauliflower since lateral broccoli shoot develop marketable head after the center head is harvested. The upper stem and clusters of unopened flower buds (heads) are the edible parts of the plant (Johnson, 2000).

Broccoli product in combination of organic with inorganic fertilizer gave as good yields as those obtained with conventional inorganic fertilizer N. Biological activity in the soil tended to increase with the increase organo-mineral fertilizer rates (Belec *et al.*, 2001).

For increasing P uptake, plants can directly modify the rhizosphere in order to gain access to unavailable soil P reserves through exuding chemical compounds into rhizosphere and through association of roots with mycorrhiza (Raghothama, 1999).

Organic and bio-fertilizers are very important sources for providing the plants with their nutritional requirements without having an undesirable impact on the environment. Many investigators compared the effect of different organic manures (Perin *et al.*, 2004 and Perez-Murcia *et al.*, 2006), biofertilizers (Manivannan and Singh, 2004 and Chaterjee *et al.*, 2005) and mineral fertilizer especially phosphorus, Kumar and Sharma (2001) reported that the maximum values for growth, yield and quality characteristics were obtained at the highest N, P and K levels (150, 90 and 60 kg/ha., respectively) as well as their interaction on broccoli growth, yield and contents of dry matter, N, P, K and chemical components in leaves and heads (Chaterjee *et al.*, 2005).

Nitrate concentrations in vegetables were positively correlated with N rates. (Karitonas, 1999 and Rydz, 2001). Nitrate may be formed from NO₃ after ingestion, causing methaemoglobinemia (Wright and Davison, 1964). Presence of NO₂ in blood might result also in the formation nitroamines, which are carcinogenic (Craddock, 1983). Recent attention has been given to less pollution practices in modern agriculture. One of the ways to reduce soil pollution is the use of biofertilizers which have been recommended by several investigators to substitute chemical fertilizers.

Therefore, the objectives of this investigation were to find out the optimal phosphorus level and most efficient biofertilizers and organic fertilizers treatments and their combination to increase broccoli growth and increase yield as well as improve head quality.

MATERIALS AND METHODS

This investigation was conducted at The Experimental Station of the Faculty of Agriculture, Cairo University, Giza during the two successive winter seasons of 2002/2003 and 2003/2004 to study the effect of different levels of phosphorus, sheep manure or the biofertilizers treatments in single or in combined applications on broccoli growth characters, yield and nutritive value of heads. The physical and chemical characteristics of the experimental soil are presented in Table (1) as average in both seasons.

The chemical properties of soil were determined using the methods described by Jackson (1962). Seeds of broccoli hybrid prominence (Takii co.,

Japan) were sown in the two seasons on mid October in seedling trays. Transplanting was carried out four weeks later.

Table (1): Physical and chemical characteristic of experimental soil

Clay %	Silt %	Fine sand %	Coarse sand %	CaCO ₃ %	pH	EC ds/m	Organic mater %	Total N %	P ₂ O ₅ ppm	K ₂ O ppm
21.2	40.5	32.6	5.3	1.6	7.8	1.01	2.11	0.09	30.1	105.8

The experimental included 16 treatments which were the combinations of sheep manure with 3 levels of phosphorus fertilizers, i.e. 4.28 Tons/fed. (10 m²/fed.) sheep manure + 0 kg P₂O₅ (P₁), sheep manure + 10 kg P₂O₅ (P₂), sheep manure + 20 Kg P₂O₅ as calcium superphosphate 15.5% P₂O₅ (P₃) or the recommended chemical fertilizers 40 kg N + 30 kg P₂O₅ + 50 kg K₂O/ fed.. All organic treatment received potassium sulphate until received 50 kg KO₂ / fed..

Broccoli seedling non inoculated or inoculated of with 3 biofertilizers treatments i.e. without biofertilizers (M₁), soil yeast *Candida tropicalis* (M₂), *Bacillus megatherium* (M₃), phosphate dissolving bacteria and *Pseudomonas aeruginosa* (M₄) at transplanting.

Sheep manure and calcium superphosphate were applied during soil preparation. On the other hand, N, P and K₂O applied as ammonium sulphate (20.5% N), calcium superphosphate 15.5% P₂O₅ and potassium sulphate (48% KO₂), respectively for chemical treatment. Application was in two equal parts at 3 and 6 weeks after transplanting.

Chemical analysis of sheep manure used is presented in Table (2) as average in both seasons. Various cultural practices were conducted as commonly followed in commercial cauliflower field.

Table (2): Chemical analysis of sheep manure used at experimental period.

Organic matter %	pH	EC ds/m	N %	P %	K %
65.7	8.01	2.3	1.9	0.4	1.2

Experimental design:

The experiment was conducted in three replicates in split plot design, where sheep manure plus levels of phosphorus presented as the main plots and biofertilizers treatments were presented as the sub plot.

The experimental plot was 14 m² and consisted of 5 rows, each 4.0 m long and 0.75 m wide. The distance between plants were 60 cm.

Data were recorded on the following characters:

Vegetative growth characters:

Ten plants taken from each plot were chosen randomly when top head initiated (55 days after planting) and when side head initiated (85 days after planting) for measuring the following characteristics:

Plant length, fresh weights of plant, stem and leaves as well as number of leaves per plant. Days from transplanting until head initiated were determined on 50% of plants.

Yield and its component: Average fresh weight of top head and side head per plant were determined from each plot. Yield of top head, yield of side head and total yield (top head + side head) were recorded in two rows and calculated as ton per feddan.

Chemical composition of leaves and heads:

Samples of leaves, at head initiation and samples of top heads at harvesting were taken and dry matter percentage was determined.

Determination of N, P and K were carried out on the ground dry materials of plants which were digested using sulfuric acid, salicylic acid and hydrogen peroxide according to Linder (1944). Nitrogen was determined using the micro-kejedahl apparatus of Parnos–Wagner as described by Van-Schouwenburg and Walinga (1978). Phosphorus was estimated colorometrically by using chlorostannous reduced molybdophosphoric blue color method according to Chapman and Parker (1961). Potassium was determined using the flame photometer. $\text{NO}_3 - \text{N}$ was determined in distilled water extracts of dried tissue by the procedure of Cataldo *et al.*, (1975) by using salicylic acid and then calculated as mg/100 gram fresh weight. Ethanol extracts of fresh materials were used for the determination of total sugars, total free amino acids and total soluble phenols. Total sugar were determined by using the phenol-sulphuric acid method (Dubois *et al.*, 1956). Total free amino acids were determined by using ninhydrin reagent according to (Moore and Stein, 1954). Total soluble phenols were estimated using the Folin-ciocalteau colorimetric method (Swain and Hillis, 1959). Vitamin C content of heads determined by the methods of A. O. A. C. (1980) .

Statistical analysis:

Yield and yield components as well as chemical composition of leaves and heads were analysis as split plot design, whereas the vegetative characters of plant samples were analyzed as split – split plot in time, plant samples were considered as sub-sub plots. Data were subjected to the proper statistical analysis of variance according to Snedecor and Cochran (1980) using the LSD at 5% level.

RESULTS AND DISCUSSION

Vegetative Growth:

Results presented in Tables (3,4and5) indicate that the most studied vegetative growth parameters, i.e., plant length, fresh weights of plant, stem and leaves as well as number of leaves per plant were significantly influenced by different levels of phosphorus and biofertilizers treatments at the two different samples (55 and 85 days after transplanting).

Regarding to the effect of phosphorus levels, significant differences were detected on vegetative parameters, the highest values were obtained by adding mixture of sheep manure + 20 kg P_2O_5 (P_3) followed by mixture of sheep manure + 10 kg P_2O_5 . (P_2)

On the other hand, applying chemical fertilizers (NPK) alone or sheep manure alone (P_1) gave the lowest values, in both seasons. While, the number of leaves per plant, were not significantly affected by the different levels of P_2O_5 fertilization in the second season.

Table (3): Effect of sheep manure, phosphorus levels, chemical fertilizers, biofertilizers treatment, plant age and their interaction on length and fresh weight of plant.

Concerning the effect of the three different types of biofertilizers, data presented in Tables (3,4 and 5) reveal that plant length was the tallest by applying *Pseudomonas aeruginosa* (M₄) followed by *Candida tropicalis* (M₂) and *Bacillus megatherium* (M₃) while without using biofertilizers plant length was the lowest, in both seasons.

Plant fresh weight was the heaviest by applying M₄, while there were no significant differences between M₂ and M₃ in the first season. On the other, hand M₂ resulted in the highest values, while there were no significant differences between M₂, M₃ and M₄ in the second season. Without using biofertilizers (M₁), plants had the lowest values in both seasons.

Stem fresh weight was the heaviest by using *Pseudomonas aeruginosa* (M₄) or without using biofertilizers, while *Candida tropicalis* (M₂) gave the lowest values in the first season. *Candida tropicalis* (M₂) followed by *Pseudomonas aeruginosa* (M₄) gave the heaviest values while without using biofertilizers (M₁) gave the lowest values in the second season.

Leaves fresh weight per plant were the highest by using M₄ and M₂ in the first and second seasons, respectively. While, leaves fresh weight were the lowest without using biofertilizers in both seasons.

Number of leaves per plant were the highest by applying *Pseudomonas aeruginosa* (M₄) or *Bacillus megatherium* (M₃) without significant differences between them, in the first season. *Candida tropicalis* (M₂) followed by *Pseudomonas aeruginosa* (M₄) gave the highest values in the second season. The lowest values were obtained without using biofertilizers in both seasons.

As shown in Tables (3, 4 and 5), there were significant increases in plant length, fresh weight of plant, stem and leaves as well as number of leaves per plant with the progress of plant age.

It is clear from Tables (3,4 and 5) that the interaction between phosphorus fertilization and biofertilizers treatments were significant in both seasons. The tallest plants were obtained when using sheep manure + 10 kg P₂O₅ (P₂) and *Pseudomonas aeruginosa* (M₄) in the first season while the interaction between P₂×M₂ (*Candida tropicalis*) resulted in the tallest one in the second season.

The heaviest fresh weight of plants were obtained as a result of interaction between P₃×M₄ (P₃ : sheep manure + 20 kg P₂O₅, M₄ : *Pseudomonas aeruginosa*) in both seasons.

Without using biofertilizers treatments (M₁), broccoli plants were the lower in the length and fresh weight at all levels of phosphorus fertilizer. P₁×M₁ gave the lowest values.

The highest fresh weight of stem, leaves and number of leaves per plant were obtained by the interaction between P₃×M₄ in the first season, but the interaction between P₂×M₂ gave the highest values in the second season. The lowest values of stem fresh weight were obtained with the interaction NPK×M₄ and NPK×M₂ in the first and second seasons, respectively. The lowest leaves fresh weight and number of leaves per plant were obtained from the interaction NPK ×M₁ and P₁×M₁ in the first and second seasons, respectively.

Table (4): Effect of sheep manure, phosphorus levels, chemical fertilizers, biofertilizers treatment, plant age and their interactions on stem and leaves fresh weights per plant

Fertilizers	Biofertilizers	Stem weight /plant(g)						Leaves weight/plant (g)					
		2002-2003			2003-2004			2002-2003			2003-2004		
		C ₁	C ₂	X	C ₁	C ₂	X	C ₁	C ₂	X	C ₁	C ₂	X
P ₁	M ₁	87.5	356.7	222.1	76.3	251.7	164.0	372.5	805.0	588.8	302.5	591.7	447.1
	M ₂	95.8	380.0	237.9	82.5	366.7	224.6	393.4	796.7	595.0	372.5	906.7	639.6
	M ₃	80.0	434.3	211.7	88.8	285.0	186.9	389.6	823.3	606.5	367.5	677.7	567.1
	M ₄	121.3	322.5	221.9	80.0	285.0	182.5	390.0	905.0	647.5	427.5	617.5	522.5
	X	96.1	350.6	223.4	81.9	297.1	189.5	386.4	832.5	609.4	367.5	720.6	544.1
P ₂	M ₁	91.2	407.5	249.4	110.0	316.7	213.3	351.3	725.0	538.1	330.0	630.0	480.0
	M ₂	91.2	323.3	207.3	160.4	350.0	255.2	441.3	843.3	642.3	479.2	946.7	712.9
	M ₃	97.5	311.7	204.6	101.2	333.3	217.3	411.3	873.3	642.3	391.3	826.7	609.0
	M ₄	95.4	425.0	260.2	88.8	266.7	177.7	424.6	888.3	656.5	480.0	805.0	642.5
	X	93.8	366.9	230.4	115.1	316.7	215.9	407.1	832.5	619.8	420.1	802.1	611.1
P ₃	M ₁	132.5	383.3	257.9	85.0	283.3	184.2	392.5	863.3	627.9	412.5	656.7	534.6
	M ₂	90.0	380.0	235.0	82.5	373.3	227.9	416.3	888.3	652.3	421.3	866.7	644.0
	M ₃	105.0	400.0	252.5	67.5	340.0	203.8	472.5	923.3	697.9	541.3	743.3	642.3
	M ₄	108.7	466.7	287.7	127.5	361.7	244.6	496.3	966.7	731.5	498.8	916.7	707.7
	X	109.1	407.5	258.3	90.6	339.6	215.1	444.4	910.4	677.4	468.5	795.8	632.1
NPK	M ₁	111.2	350.0	230.6	67.5	283.3	175.4	311.3	570.0	440.6	383.8	661.7	522.7
	M ₂	113.7	313.3	213.5	41.7	233.3	137.5	466.3	725.0	595.6	452.5	700.0	576.3
	M ₃	118.8	390.0	254.4	46.3	251.7	149.0	335.0	710.0	522.5	415.0	700.0	557.5
	M ₄	97.5	270.0	183.8	105.0	308.3	206.7	361.2	915.0	638.1	388.8	840.0	614.4
	X	110.3	330.8	220.6	65.1	269.2	167.1	368.4	730.0	549.2	410.0	725.4	567.7
The interaction between biofertilizers treatments and plant age on stem and leaves fresh weight per plant													
	M ₁	105.6	374.3	240.0	84.7	283.8	184.2	356.9	740.0	548.9	357.2	635.0	496.1
	M ₂	97.7	349.2	223.4	91.8	330.8	211.4	429.3	813.3	621.3	431.4	855.0	643.2
	M ₃	100.3	361.3	230.8	75.9	302.5	189.2	402.1	832.5	617.3	428.8	759.2	594.0
	M ₄	105.7	371.0	238.4	100.3	305.4	202.9	418.0	918.8	668.4	448.8	794.8	621.8
	X	102.3	364.0		88.2	305.6		401.6	826.4		416.6	671.0	
LSD 0.05	P:1585 M:1292 C:847				P:1371 M:1591 C:982				P:464 M:533 C:378				P:749 M:619 C:395
	P x M: 25.83				P x M: 31.8				P x M: 106.5				P x M: 123.9
	P x C : 16.95				P x C : 19.6				P x C : 75.6				P x C : 78.9
	M x C: 16.95				M x C: 19.6				M x C: 75.6				M x C: 78.9
	P x M x C: 33.89				P x M x C: 39.3				A x M x C: 151.2				P x M x C:157.9
P ₁ : Sheep manure+0 kg P ₂ O ₅ /fed						P ₂ : Sheep manure+10 kg P ₂ O ₅ /fed.							
P ₃ : Sheep manure + 20 kg P ₂ O ₅ /fed.						NPK: 40 kg. N+ 30 kg P ₂ O ₅ + 50 kg K ₂ O/fed.							
M ₁ : Without inoculation						M ₂ : <i>Candida tropiclis</i>							
M ₃ : <i>Bacillus megatherium</i>						M ₄ : <i>Pseudomonas aeruginosa</i>							
C ₁ : 55 days after transplanting						C ₂ : 85 days after transplanting							

In general, without applying biofertilizers treatments the vegetative growth values were lower comparing with using biofertilizers treatments. The interaction between levels of phosphorus fertilization and the age of the plant were significant in both seasons .The highest values of vegetative growth were obtained by using sheep manure + 20 kg P₂O₅ (P₃) at the second stage of growth C₂ (at side heads initiation) in both seasons.

Results presented in Tables (3,4 and 5) indicated that there were significant differences in all studied vegetative growth characters due to the effect of interaction of biofertilizers treatments and the age of the plant. The greatest values of these parameters were recorded in the plant treated with *Pseudomonas aeruginosa* (M₄) and *Candida tropicallis* (M₂) in the first and

second seasons respectively at the second stage of plant growth. Whereas , the lowest values were recorded in plants untreated with biofertilizers treatment at the first stage of growth in both seasons.

The effect of the interaction between levels of phosphorus fertilization, biofertilizers treatments and the age of the plant were significant. The greatest values of the most vegetative growth characters were obtained with using sheep manure + 20 kg P₂O₅ (P₃) accompanied with the application of *Pseudomonas* (M₄) at the second stage of the growth (P₃×M₄×C₂).

Table (5): Effect of sheep manure, phosphorus levels, chemical fertilizers, biofertilizers treatment, plant age and their interactions on number of leaves per plant.

Fertilizers	Biofertilizers	Leaves numbers/plant					
		2002-2003			2003-2004		
		C ₁	C ₂	X	C ₁	C ₂	X
P ₁	M ₁	16.43	16.66	16.55	13.93	14.33	14.13
	M ₂	16.70	17.30	17.02	16.10	20.33	18.22
	M ₃	18.30	17.33	17.82	15.27	17.00	16.13
	M ₄	17.33	19.33	18.33	15.83	16.00	15.92
	X	17.19	17.67	17.43	15.28	16.92	16.10
P ₂	M ₁	15.66	16.00	15.83	15.00	15.67	15.33
	M ₂	15.90	18.33	17.12	16.27	19.33	17.80
	M ₃	18.50	19.50	19.00	16.83	17.67	17.25
	M ₄	16.33	21.00	18.67	16.67	17.67	17.17
	X	16.60	18.71	17.65	16.19	17.58	16.89
P ₃	M ₁	15.90	17.60	16.75	15.00	16.67	15.83
	M ₂	17.43	18.33	17.88	15.77	18.67	17.22
	M ₃	17.93	22.00	19.97	15.83	17.00	16.42
	M ₄	18.90	23.0	20.95	15.83	18.67	17.25
	X	17.54	20.33	18.89	15.61	17.75	16.68
NPK	M ₁	14.67	15.00	14.83	14.53	17.00	15.77
	M ₂	15.43	15.50	15.47	15.83	17.00	16.42
	M ₃	15.60	18.30	16.97	15.17	17.00	16.08
	M ₄	15.30	21.00	18.17	15.00	18.30	16.67
	X	15.26	17.46	16.36	15.13	17.33	16.23
The interaction between biofertilizers treatments and plant age on number of leaves per plant.							
	M ₁	15.67	16.32	15.90	14.62	15.92	15.27
	M ₂	16.37	17.38	16.87	15.99	18.83	17.41
	M ₃	17.58	19.29	18.44	15.77	17.17	16.47
	M ₄	16.98	21.08	19.03	15.83	17.67	16.75
	X	16.65	18.52		15.55	17.40	
LSD 0.05	P: 0.64 M: 0.89 C: 0.51			P: ns M: 0.77 C: 0.67			
	P x M: 1.78			P x M: 1.53			
	P x C : 1.02			P x C : 1.33			
	M x C: 1.02			M x C: 1.33			
	P x M x C: 2.03			P x M x C: 2.66			
P ₁ : Sheep manure+0 kg P ₂ O ₅ /fed				P ₂ : Sheep manure + 10 kg P ₂ O ₅ /fed.			
P ₃ : Sheep manure + 20 kg P ₂ O ₅ /fed.				NPK: 40 kg. N+ 30 kg P ₂ O ₅ + 50 kg K ₂ O/fed.			
M ₁ :Without inoculation				M ₂ : <i>Candida tropiclis</i>			
M ₃ : <i>Bacillus megatherium</i>				M ₄ : <i>Pseudomonas aerugionsa</i>			
C ₁ : 55 days after transplanting				C ₂ : 85 days after transplanting			

Days from transplanting to head initiation (DTH):

Concerning the effect of phosphorus levels and biofertilizers treatments and their interactions, data presented in Table (6) revealed that there were no significant differences between phosphorus levels on days from transplanting to head initiation (DTH) in both seasons. On the other hand, there were significant differences among biofertilizers treatment for DTH. Applying biofertilizers resulted in lower days from transplanting to head initiation compared to without using biofertilizers. There were no significant differences between biofertilizers treatments in the first season, while *Bacillus* and *Pseudomonas* resulted in lower days compared to *Candida*, in the second season.

With respect to the interaction between phosphorus fertilization and biofertilizers treatments, data in Table (6) show significant differences. The lowest days from transplanting to head initiation were obtained from plants fertilized with P₂ and treated with M₃ (*Bacillus*) or M₂ (*Candida*) as well as from plant fertilized with P₃ and treated with M₃ (*Bacillus*) in the first season. The same trend was obtained by M₄ (*Pseudomonas*) or M₃ (*Bacillus*) treatments which gave the lowest values when combined with all different kinds of fertilizers in the second season.

In general, applying biofertilizers treatments with all kinds of fertilizers resulted in the lowest days from transplanting to head initiation compared to without biofertilizers treatments.

Yield and its components:

Data presented in Table (6) revealed significant differences in the average top head and side head weight per plant as well as top heads, side head yields/fed. and total yield/fed. due to the effect of phosphorus levels. The highest value of these characters were recorded by the plants fertilized with sheep manure + 20 kg P₂O₅ (P₃), meanwhile the lowest values were obtained when applying sheep manure (P₁) or (chemical fertilization) NPK in the first and second seasons, respectively.

Concerning the effect biofertilizers treatments, data presented in Table (6) indicated that the highest values of average head weight and yield of heads/fed. were obtained by the plant supplied with *Pseudomonas* (M₄) followed by *Bacillus* (M₃) in the first season. Meanwhile, (*Candida*) M₂ followed by (*Pseudomonas*) M₄ gave the highest values in the second season. On the other hand, average side head weight per plant and yield of side heads/fed. were the greatest when applying *Pseudomonas* (M₄) followed by *Candida* (M₂) in the first season. In the second season, *Pseudomonas* (M₄) followed *Bacillus* (M₃) resulted in the highest values. While, total yield/fed. was the greatest when applying (*Pseudomonas*) M₄ in both seasons. On the other hand, the lowest values of these characters were obtained by the plants untreated with any of biofertilizers treatments in both seasons.

With respect to the effect of interaction between levels of phosphorus fertilization and biofertilizers treatments on yield and its components, data in Table (6) indicated significant differences. The highest average head weight, yield of heads/fed. and total yield /fed. were obtained by the plants fertilized with sheep manure+20 kg P₂O₅(P₃) and treated with *Pseudomonas*(M₄) in the first season. While, in the second season the interaction between P₃ and M₃

(*Bacillus*) gave the highest values. Meanwhile, the highest values of side heads weight / plant and yield of side heads /fed. were noticed in plants fertilized with sheep manure+10 kgP₂O₅ (P₂) and treated with *Pseudomonas* (M₄) in the first season. On the other hand , P₃ ×M₃ gave the highest values in the second season. However, the lowest values of average side heads/plant, yield of side heads/fed. and total yield/fed. were obtained with interaction P₂ ×M₁ in the first season. Meanwhile, average head weight and yield of heads/fed., in both seasons , average side heads/plant, yield of side heads / fed. and total yield were the lowest with using the interaction P₁×M₁ in the second season.

The results indicated that combined of sheep manure, phosphorus fertilizer and biofertilizers treatments affected in the productivity of broccoli plants.

From these results, it can be concluded that most of the studied vegetative growth parameters as well as yield and its components were increased with increasing phosphorus levels. Similar results and suggestion were reported by many workers on broccoli plants.

The effect of N, P and K fertilizers were in the same trend of studies were obtained by Dufault (1988), Kumar and Sharma (2001), Brahma *et al.*, (2002), Sharma *et al.*, (2002) and El-Magd *et al.*, (2005) who reported that increasing any of N, P or K ratio improved vegetative growth expressed as plant height and number of leaves per plant as well as total yield and head quality.

The vegetative growth and yield were affected by phosphorus levels. Wang *et al.*, (1997) Cited that application of N+ P gave 110% higher yields than N alone.

In this respect, this enhancement may be explained that phosphorus is a part of molecular structural of several vitally important compounds notable nucleic acids (DNA, the two forms of RNA). In addition, phosphorus plays role in the enzyme system necessary for energy transformation in photosynthesis and respiration. It is also a constituent of cell nucleus and essential for division for the development of meristematic tissue (Edmont *et al.*,1981) .

Furthermore, from the results it can be suggest that, vegetative growth parameters and yield of plants received combined of sheep manure and mineral fertilizer were higher than those supplied with mineral fertilizers alone or organic manure. These results are in agreement with those obtained by Subhan (1989) on cabbage, Sharma (2000) on broccoli plants and Bhardwai *et al.*,(2000) on cauliflower and tomato. The yield of broccoli harvested one week earlier and higher with combination of organic and mineral fertilizer. This was related to improved P nutrition (Brown *et al.*, 1994). In manured plots available P₂O₅ content increased. Cabbage yield were highest when chemical fertilizer were applied with FYM (Kamaiyama *et al.*, 1995). For every 1 ton compost applied, fertilizer N application could be reduced by about 10 kg (Inoune *et al.*, 1996).

Regarding the effect of biofertilizers treatments, results indicated that broccoli plants treated with biofertilizers, i.e. *Candida tropicalis*, *Bacillus megatherium* (Phosphate dissolving) or *Pseudomonas* had higher vegetative growth parameters, yield and its components compared with non biofertilizers

treatment. *B. megatherium* or *P. aeruginosa* were more superior than *C. tropicalis* in most characters. These results were confirmed by Subba Rao (1993) who reported that biofertilizers stimulate plant growth by increasing the number and biological activity of desired micro-organisms in root environment. Some of these micro-organisms fix-atmospheric nitrogen or convert of bond phosphorus or bond potassium to easily stimulated form. Broccoli plant received combination of organic, mineral fertilizer and biofertilizers had higher vegetative parameter and yield comparing with using organic or chemical alone. Inoculation plants or soil with different micro-organic has become recently new methods to reduce the amount of inorganic complete fertilizers.

Table (6): Effect of sheep manure, phosphorus levels, chemical fertilizers, biofertilizers treatment, and their interactions on day to head initiation, head weight, side head weight, head yield, side head yield and total yield

Fertilizers	Biofertilizers	Day to head initiation		Head weight (gm.)		Side head weight (gm/plant)		Head yield (tons/fed.)		Side head yield (tons/fed.)		Total yield (Tons/fed.)	
		2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
P ₁		57.2	58.7	504.0	557.0	290.0	268.8	4.480	4.951	2.578	2.389	7.059	7.341
P ₂		56.2	58.8	523.7	576.0	311.1	271.6	4.655	5.119	2.766	2.415	7.421	7.534
P ₃		56.9	59.0	589.1	606.9	332.9	298.6	5.237	5.394	2.959	2.654	8.192	8.049
NPK		56.6	58.4	539.2	548.0	291.0	243.8	4.793	4.871	2.588	2.167	7.380	7.039
LSD 0.05		NS	NS	50.12	25.66	40.42	5.69	0.445	0.230	0.360	0.055	0.551	0.209
	M ₁	57.8	60.6	520.9	502.3	263.7	181.9	4.631	4.464	2.344	1.617	6.975	6.082
	M ₂	56.1	58.7	521.5	625.7	311.3	243.7	4.635	5.562	2.768	2.167	7.404	7.728
	M ₃	56.4	57.8	556.3	546.5	301.6	319.9	4.945	4.858	2.681	2.844	7.626	7.702
	M ₄	56.7	57.9	557.2	613.5	348.5	337.3	4.953	5.453	3.098	2.998	8.051	8.451
LSD 0.05		0.59	0.75	32.38	31.68	35.1	20.66	0.288	0.276	0.312	0.183	0.476	0.332
	M ₁	59.1	60.6	470.9	413.2	289.5	143.7	4.186	3.673	2.573	1.277	6.759	4.950
	M ₂	56.0	59.4	474.4	626.1	293.2	256.7	4.217	5.563	2.607	2.282	6.824	7.847
	M ₃	57.2	57.6	522.5	537.2	289.3	352.9	4.644	4.774	2.572	3.129	7.217	7.903
	M ₄	56.7	57.5	547.9	651.8	287.9	322.9	4.870	5.794	2.559	2.870	7.429	8.664
	M ₁	56.5	60.4	515.1	536.2	220	219.5	4.579	4.767	1.926	1.951	6.535	6.718
	M ₂	55.8	58.5	495.0	603.2	302.9	271.1	4.400	5.362	2.692	2.410	7.092	7.772
	M ₃	55.8	57.9	573.9	536.2	300.8	236.3	5.102	4.764	2.673	2.100	7.775	6.864
	M ₄	56.8	58.3	510.2	628.5	420.8	359.7	4.540	5.586	3.741	3.197	8.281	8.783
	M ₁	59.3	61.8	590.2	509.5	296.0	186.8	5.246	4.528	2.631	1.661	7.877	6.189
	M ₂	56.2	59.1	506.6	658.0	373.0	264.6	4.503	5.849	3.316	2.352	7.819	8.201
	M ₃	55.8	57.3	594.8	666.2	290.1	409.6	5.287	5.922	2.579	3.641	7.866	9.563
	M ₄	56.3	57.9	664.9	593.9	372.6	333.4	5.911	5.279	3.312	2.963	9.223	8.242
	M ₁	56.3	59.4	507.5	550.2	249.2	177.8	4.511	4.890	2.215	1.581	6.726	6.471
	M ₂	56.4	58.0	609.9	615.4	276.0	182.6	5.422	5.471	2.458	1.623	7.880	7.094
	M ₃	56.7	58.3	534.2	446.6	326.2	281.7	4.748	3.970	2.900	2.504	6.648	6.474
	M ₄	57.0	57.8	505.1	579.8	312.5	333.1	4.49	5.154	2.778	2.961	7.268	8.150
LSD 0.05		1.19	1.50	64.77	63.35	70.20	41.32	0.576	0.551	0.624	0.365	0.953	0.666
P ₁ : Sheep manure+0 kg P ₂ O ₅ /fed						P ₂ : Sheep manure+10kg P ₂ O ₅ /fed.							
P ₃ : Sheep manure+ 20 kg P ₂ O ₅ /fed.						NPK: 40 kg. N+ 30 kg P ₂ O ₅ + 50 kg K ₂ O/fed.							
M ₁ : Without inoculation						M ₂ : <i>Candida tropicalis</i>							
M ₃ : <i>Bacillus megatherium</i>						M ₄ : <i>Pseudomonas aeruginosa</i>							

Many investigators studied the effect of microorganism on the growth and yield. Hsieh and Hsu (1995) studied the effect of addition of microorganism mixture EM (containing nitrogen-fixing, phosphate-dissolving and photosynthetic microorganisms, vesicular – arbuscular mycorrhizas, *Bacillus*, *Lactobacillus*, yeast and *actinomycetes*). The increases in cabbage yield in the organic plots compared with chemical plots were greater, especially when the microorganism mixture were included in the manures. Cabbage performed best with chicken manure compost combined with microorganisms. Bahadur *et al.*, (2005) evaluated the effect of organic manures and biofertilizers on growth and yield of cabbage. Biofertilizers were *Azospirillum*, vesicular arbuscular mycorrhiza (VAM) and phosphate solubilizers microorganisms (PSM). Pressmud + VAM+ the recommended dose of NPK (120:60:80 kg/ha) recorded the highest values for all parameters of growth, yield and average head weight. Chatterjee *et al.* (2005) studied the effect of different organic amendments and biofertilizers (*Azotobacter*, phosphate solubilizer and potash mobilizer) on broccoli. Mustard oil cake+ biofertilizer resulted in highest yield while recommended NPK rate, 150:60:80 kg/ha gave the lowest value.

Concerning the day from transplanting until head initiation, Gauss and Taylor (1996) reported on broccoli that at 8 weeks the apices reached its maximum width and 83% of the apices were initiation first order floral stalks. At 9 weeks, when the plants had an averages of 17 leaves, all apices were in the same stage of flowering and 52% of the plants sampled had inflorescence heads or button greater than 5.0 mm in diameter. Fortunato and Damato (2000) found that broccoli plants sown at October, apex transition occurred 55 days and ended 30 days afterwards.

Inoculation plants with microbial mixture and supplemented with organic and chemical fertilizer led to significantly decreased days from transplanting until flowering comparing with applying chemical fertilization (Darwesh, 2002 on tomato and Alian, 2005 on artishok).

Regarding the relationship between the growth of broccoli plants and plant age, results indicated that the highest values of vegetative growth parameter were recorded when side heads initiated after 83 days of transplanting. These results agreement with results obtained by Rincon *et al.*, (1999) who reported that the highest growth rate occurred between 73 and 87 days after transplanting.

Chemical composition in leaves and heads:

Data in Table (7) indicated clearly significant differences between phosphorus levels on the percentage of (N, P and K). The maximum values of N, P and K in leaves and heads were obtained as a result of fertilization with the highest phosphorus level 20 kg P₂O₅ + sheep manure (P₃). While, application of sheep manure only (P₁) gave the lowest values of N and K in leaves as well as N, P and K in heads. However, the lowest values of P in leaves were obtained with applying chemical fertilization (NPK), in both seasons.

Moreover, it is clear from data in Table (7) that application of M₄ (*Pseudomonas*) resulted in the highest values of N, P and K in the first season in the leaves. While M₂ (*Candida*) and M₃ (*Bacillus*) resulted in the

highest values of P and K, respectively in the second season. As regard to heads, (*Pseudomonas*) M₄ increased significantly N% in the first season. Meanwhile, (*Candida*) M₂ resulted in the highest values of N in the second season. The percentage of P and K% in head increased significantly with applying (*Bacillus*) M₃ in both seasons. Without using biofertilizers treatments (M₁), N, P and K % in the leaves and heads were the lowest in both seasons.

Table (7): Effect of sheep manure, phosphorus levels, chemical fertilizers, biofertilizers treatment, and their interactions on N% , P% and K% in leaves and heads.

Fertilizers	Biofertilizers	Leaves						Heads					
		N%		P%		K%		N%		P%		K%	
		2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
P ₁		2.39	2.59	0.234	0.289	1.69	1.76	2.05	2.23	0.246	0.199	1.86	1.97
P ₂		2.73	2.74	0.260	0.303	1.87	2.11	2.46	2.44	0.286	0.338	2.43	2.15
P ₃		2.73	2.91	0.294	0.391	2.38	2.20	2.66	2.51	0.322	0.361	2.44	2.32
NPK		2.62	2.70	0.218	0.225	1.86	1.86	2.41	2.21	0.249	0.237	2.17	2.21
LSD 0.05		0.223	0.14	0.032	0.009	0.17	0.17	0.099	0.245	0.032	0.032	0.141	0.105
	M ₁	2.01	2.52	0.155	0.222	1.93	1.71	2.09	2.03	0.218	0.214	1.97	2.01
	M ₂	2.83	2.75	0.210	0.352	1.88	1.83	2.35	2.57	0.297	0.322	2.34	2.15
	M ₃	2.69	2.85	0.308	0.333	1.77	2.41	2.37	2.42	0.379	0.353	2.38	2.33
	M ₄	2.93	2.82	0.333	0.305	2.22	1.97	2.77	2.36	0.210	0.246	2.21	2.17
LSD 0.05		0.190	0.21	0.008	0.027	0.107	0.08	0.169	0.146	0.027	0.027	0.146	0.148
P ₁	M ₁	1.85	2.62	0.143	0.163	1.38	1.54	1.750	1.91	0.213	0.160	1.80	1.75
	M ₂	2.44	2.59	0.133	0.410	2.08	1.54	1.93	2.54	0.253	0.208	1.83	1.94
	M ₃	2.41	2.53	0.394	0.375	1.56	2.18	2.00	2.47	0.330	0.279	1.93	2.16
	M ₄	2.88	2.66	0.265	0.208	1.73	1.76	2.53	1.99	0.188	0.149	1.86	2.04
P ₂	M ₁	2.61	2.16	0.181	0.214	1.50	2.00	2.35	2.07	0.203	0.213	2.04	1.96
	M ₂	2.89	2.72	0.174	0.423	1.56	2.04	2.36	2.85	0.381	0.438	2.73	2.16
	M ₃	2.62	3.01	0.321	0.264	2.20	2.19	2.54	2.63	0.325	0.353	2.53	2.30
	M ₄	2.79	3.05	0.365	0.311	2.21	2.20	2.60	2.22	0.235	0.348	2.43	2.16
P ₃	M ₁	1.99	2.69	0.166	0.285	2.40	1.68	2.06	2.09	0.231	0.326	1.95	2.16
	M ₂	2.86	2.75	0.298	0.419	2.40	1.98	2.68	2.68	0.328	0.419	2.53	2.28
	M ₃	3.0	3.12	0.304	0.421	2.06	3.34	2.69	2.44	0.500	0.406	2.74	2.54
	M ₄	3.06	3.07	0.409	0.440	2.66	1.78	3.21	2.81	0.230	0.291	2.54	2.30
NPK	M ₁	1.61	2.59	0.130	0.226	2.43	1.64	2.23	2.05	0.224	0.156	2.07	2.16
	M ₂	3.15	2.94	0.236	0.156	1.46	1.74	2.43	2.22	0.226	0.221	2.28	2.23
	M ₃	2.76	2.75	0.211	0.258	1.26	1.93	2.25	2.14	0.361	0.375	2.33	2.30
	M ₄	2.94	2.51	0.293	0.260	2.28	2.13	2.72	2.41	0.188	0.194	2.00	2.16
LSD 0.05		0.380	0.413	0.017	0.053	0.185	0.17	0.337	0.292	0.053	0.053	0.292	0.297
P ₁ : Sheep manure+0 kg P ₂ O ₅						P ₂ : Sheep manure+10 kg P ₂ O ₅ /fed.							
P ₃ : Sheep manure + 20 kg P ₂ O ₅ /fed.						NPK: 40 kg. N+ 30 kg P ₂ O ₅ + 50 kg K ₂ O/fed.							
M ₁ : Without inoculation						M ₂ : <i>Candida tropiclis</i>							
M ₃ : <i>Bacillus megatherium</i>						M ₄ : <i>Pseudomonas aeruginosa</i>							

Concerning the interaction between phosphorus level and biofertilizers treatments, there were significant differences. The interaction of P₃ × M₃ or P₃ × M₄ resulted in the highest percentage of N, P and K in both leaves and head in both seasons.

Dry matter percentage, total sugars, total soluble phenols, total free amino acids concentration in the leaves and heads (Tables 8 and 9) were significantly influenced by phosphorus levels. Sheep manures + 20 kg P₂O₅ (P₃) resulted in the greatest values followed by sheep manure + 10 kg P₂O₅

(P₂). Meanwhile, the lowest values were detected in leaves and heads of the plants fertilized with sheep manure only (P₁) or NPK chemical fertilizers. These results were obtained in two seasons.

Regarding the effect of biofertilizers treatments, data in Tables (8 and 9) indicated that plants received *Bacillus* (M₃) or *Pseudomonas* (M₄) had the highest values of dry matter percentage, total sugars, total soluble phenols and total free amino acids concentration in the leaves in both seasons. Meanwhile, inoculated with (*Candida*) M₂ resulting in the greatest phenols and total free amino acids concentration of the head in the first and second seasons, respectively. On the other hand, plants untreated with biofertilizers gave the lowest values of chemical components in leaves and heads.

Table (8): Effect of sheep manure, phosphorus levels, chemical fertilizers, biofertilizers treatment, and their interactions on dry matter %, total sugars, total soluble phenols and total free amino acids concentration (mg/100g f . w) in leaves

Fertilize-rs	Biofertilizers	Dry matter %		Total sugars		Total soluble phenols		Total free amino acids	
		2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
P ₁		13.15	12.81	60.9	70.8	43.3	33.3	246.3	294.7
P ₂		13.21	12.76	86.3	103.8	48.6	43.2	335.9	369.0
P ₃		13.30	12.83	87.9	111.4	50.2	47.8	366.9	541.3
NPK		12.22	12.02	61.3	78.8	34.3	33.7	321.0	362.5
LSD 0.05		0.926	ns	3.66	3.35	2.86	4.92	11.83	8.97
	M ₁	12.72	12.00	43.7	57.6	33.0	28.4	280.3	359.7
	M ₂	12.85	12.71	78.1	98.5	36.5	39.6	317.8	384.2
	M ₃	13.41	12.86	77.5	101.9	57.9	43.0	296.9	398.1
	M ₄	12.88	12.85	96.3	106.7	48.9	46.9	375.0	425.6
LSD 0.05		0.601	0.521	4.23	5.22	4.17	3.83	7.14	4.410
P ₁	M ₁	12.69	12.60	34.2	38.1	30.9	24.9	181.1	274.2
	M ₂	12.94	12.67	61.2	75.8	27.4	36.2	251.1	273.9
	M ₃	13.85	12.72	72.8	83.7	64.2	30.6	206.1	307.8
	M ₄	13.10	13.25	75.7	85.4	50.8	41.3	346.7	322.8
P ₂	M ₁	13.44	12.18	46.8	77.9	33.9	24.1	319.5	336.7
	M ₂	13.25	13.10	106.9	142.2	40.7	40.2	366.1	398.9
	M ₃	12.81	12.62	90.90	107.8	69.4	27.8	313.9	352.7
	M ₄	13.35	13.14	100.5	87.3	50.3	80.5	343.9	387.8
P ₃	M ₁	12.97	12.18	53.2	69.7	40.8	27.5	305.6	501.2
	M ₂	13.46	12.62	69.6	93.9	29.2	38.8	306.1	527.3
	M ₃	13.94	13.0	92.6	135.1	64.6	86.9	362.6	543.9
	M ₄	12.83	13.51	136.5	147.1	66.1	37.9	493.4	592.8
NPK	M ₁	11.79	11.02	40.6	44.6	26.4	37.2	315.0	326.7
	M ₂	11.74	12.47	78.6	82.2	48.8	43.2	347.0	336.7
	M ₃	13.05	13.11	53.5	81.2	33.3	26.8	305.0	387.8
	M ₄	12.32	11.47	72.6	107.0	28.7	27.7	316.1	398.9
LSD 0.05		1.201	1.043	8.56	9.04	8.34	7.65	14.28	8.82
P ₁ : Sheep manure+ 0kg kg P ₂ O ₅ /fed		P ₂ : Sheep manure+10kg P ₂ O ₅ /fed.							
P ₃ : Sheep manure + 20 kg P ₂ O ₅ /fed.		NPK: 40 kg. N+ 30 kg P ₂ O ₅ + 50 kg K ₂ O/fed.							
M ₁ : Without inoculation		M ₂ : <i>Candida tropiclis</i>							
M ₃ : <i>Bacillus megatherium</i>		M ₄ : <i>Pseudomonas aerugionsa</i>							

Data presented in Tables (8 and 9) show that dry matter percentage, total sugars, total soluble phenols and total free amino acids concentration in leaves and heads were significantly affected due to the interaction between phosphorus levels and biofertilizers treatments. The highest values of these chemical composition in leaves and heads were detected when plants fertilized with sheep manure + 20 kg P₂O₅ (P₃) and treated with (*Bacillus*) M₃ or (*Pseudomonas*) M₄ in both seasons.

The highest values of total phenols were found in leaves of plants fertilized with sheep manure + 10 kg P₂O₅ (P₂) and treated with *Bacillus* (M₃) in the first season, while the interaction between 20 kg P₂O₅+sheep fertilizer (P₃) and *Candida* (M₂) resulted in the highest values of dry matter percentage and total soluble phenols concentration in the heads in the first season. On the other hand, the lowest values of most chemical components were obtained with interaction NPK × M₁ or P₁ × M₁ in leaves and heads in both seasons.

As regarding ascorbic acid (VC) concentration in broccoli heads, data in Table 9 revealed that, VC values increased significantly with increasing phosphorus level up to 10 P₂O₅ + sheep manure (P₂) in both seasons. Ascorbic acid in heads of plants received chemical fertilizers alone (NPK) had lower values compared to P₂ in both seasons. The lowest values recorded with application sheep manure only (P₁). Biofertilizers treatments resulted in significant increasing in ascorbic acid concentration in heads comparing with uninoculated ones. Plants were treated with *Bacillus* followed by *Pseudomonas* had the highest values of VC when compared with those inoculated with *Candida*. The interaction between phosphorus and biofertilizers treatments were significant. The highest values were obtained from plants received P₂ × M₃ followed by NPK × M₄ in the first season, while the highest values were recorded by the plants treated with NPK × M₃ and P₂ × M₃ in the second season. However, P₁ × M₁ treatment gave the lowest values in both seasons.

Concerning the effect of phosphorus level on nitrate concentration, data in Table (9) revealed that nitrate concentration in broccoli heads significantly decreased with increasing P level. P₃ (sheep manure + 20 kg P₂O₅) treatment had the lowest values. Chemical fertilizer alone (NPK) gave the highest values in both seasons.

Moreover, the results indicated that biofertilizers treatments increased significantly nitrate concentration in head compared to uninoculated plants. Plants inoculated with *candida* (M₂) had the highest values while, inoculation with *Bacillus* (M₃) decreased nitrate in heads in both seasons.

The interaction between phosphorus fertilization and biofertilizers treatments were significant in both seasons. Plants received P₃ × M₁ and P₃ × M₃ gave the lowest values in the first and second seasons, respectively. Moreover, the results indicated that nitrate concentration in heads decreased with increasing phosphorus addition by adding phosphate to the soil or as a result of using active dissolving phosphate bacteria (*Bacillus megatherium*.) such values were still in the acceptable ranges for human health (Wang and Li, 2004)

Table (9): Effect of sheep manure, phosphorus levels, chemical fertilizers, biofertilizers treatment, and their interactions on dry matter%, total sugars, total soluble phenols, total free amino acids, ascorbic acid and nitrate concentration (mg/100g f. w) in heads.

Fertilizers	Biofertilizers	Dry matter %		Total sugars		Total phenols		Total amino acids		Ascorbic acid		Nitrate	
		2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
P ₁		11.67	11.28	125.8	120.3	36.5	40.4	484.0	576.1	108.1	76.3	243.2	228.6
P ₂		12.45	11.62	166.0	124.8	44.9	42.5	517.8	613.6	146.2	147.5	231.2	220.3
P ₃		13.60	13.19	179.4	180.1	50.2	51.6	605.0	636.1	126.3	146.3	210.6	192.0
NPK		11.73	11.10	157.9	106.5	38.0	39.5	466.0	540.7	135.6	126.9	245.3	235.7
LSD 0.05		1.235	0.649	0.510	3.13	3.89	2.49	12.51	10.17	4.31	5.93	6.44	7.90
	M ₁	11.67	11.06	133.9	114.7	38.8	37.9	460.1	573.0	90.6	91.9	223.8	210.9
	M ₂	12.26	11.87	152.7	139.3	52.5	42.3	505.4	616.0	133.8	119.4	254.9	236.1
	M ₃	12.68	12.27	185.5	134.2	36.2	46.4	567.3	569.5	151.1	146.9	218	208.2
	M ₄	12.85	11.99	157.1	143.3	41.9	47.4	539.9	607.7	140.6	138.8	233.6	221.5
LSD 0.05		0.674	0.636	4.05	2.55	3.41	4.21	9.60	7.33	4.50	4.34	5.36	4.71
	M ₁	11.1	10.63	121.9	106.6	36.2	36.8	419.4	537.5	45.0	50.0	236.4	225.9
	M ₂	11.7	11.57	144.3	111.6	53.2	40.9	448.2	595.9	140.0	62.5	267.3	236.2
	M ₃	11.66	11.61	157.1	107.5	31.6	42.1	491.7	551.4	135.0	92.5	220.8	211.6
	M ₄	12.22	11.32	79.7	155.5	24.8	41.7	576.8	619.5	112.5	100.0	248.1	240.7
	M ₁	12.0	11.26	161.3	109.7	30.6	41.0	506.5	602.2	112.5	130	221.5	210.1
	M ₂	12.16	11.52	130.6	129.9	46.5	42.3	430.5	622.3	132.5	132.5	247.7	228.2
	M ₃	12.39	12.38	214.1	126.7	46.2	42.9	627.1	601.5	202.3	177.5	225.6	219.4
	M ₄	13.27	11.32	158.1	133.1	56.1	43.8	507.6	628.5	137.5	150.0	230.1	223.6
	M ₁	12.79	12.23	132.5	140.5	52.1	44.7	547.2	612.6	100.0	125.0	191.6	175.4
	M ₂	13.08	13.65	156.0	194.6	79.0	49.9	600.7	633.7	112.5	155.0	230.8	220.3
	M ₃	14.41	13.28	201.1	219.5	33.4	59.6	613.0	678.5	155.0	137.5	201.3	170.5
	M ₄	14.13	13.59	228.0	165.8	36.2	52.3	659.0	619.5	137.5	167.5	218.5	201.6
	M ₁	10.79	10.13	119.8	102.1	36.3	29.0	367.3	539.6	105.0	62.5	245.5	231.9
	M ₂	12.11	10.75	180.0	120.9	31.3	35.9	542.4	613.4	150.0	127.5	273.7	259.7
	M ₃	12.27	11.78	169.6	83.3	33.7	41.0	537.5	446.7	112.5	180.0	224.3	231.2
	M ₄	11.76	11.73	162.4	119.7	50.8	52.0	416.7	563.2	175.0	137.0	237.8	220.0
LSD 0.05		1.347	1.272	8.11	4.42	5.91	7.29	16.63	12.69	8.75	8.18	10.57	9.43
P ₁ : Sheep manure +0kg P ₂ O ₅ /fed						P ₂ :Sheep manure + 10 kg P ₂ O ₅ /fed.							
P ₃ : Sheep manure + 20 kg P ₂ O ₅ /fed.						NPK: 40 kg. N+ 30 kg P ₂ O ₅ + 50 kg K ₂ O/fed.							
M ₁ : without inoculation						M ₂ : <i>Candida tropiclis</i>							
M ₃ : <i>Bacillus megatherium</i>						M ₄ : <i>Pseudomonas aeruginosa</i> .							

Results indicated that the percentage of dry matter, N, P and K of leaves and heads as well as chemical component of leaves and heads, i.e. total sugar, soluble phenols, free amino acids, ascorbic acid and nitrate increased with increasing P level. Combination of phosphorus sheep manure + 10 kg P₂O₅ (P₂) was superior than chemical phosphorus fertilizer alone at 30 kg P₂O₅ / fed. Broccoli plants inoculated with biofertilizers had higher content of dry matter, N, P and K as well as chemical components of leaves and heads comparing with un inoculated plants. These results agree with those obtained by Prabhakar *et al.*, (1986) they reported that cabbage yield and P uptake increased with P up to 75 kg/ha. Lu *et al.*, (1997) mentioned that peak nutrient absorption occurred during the head formation period. The

descending order of dry weight proportions was leaves, head, stem and roots.

Bowen *et al.*, (1999) and Rincon *et al.*, (1999) found that the highest concentration of N, P and K were accumulation in leaves, head and stem respectively occurred between 73 and 87 day after transplanting. Nkoa *et al.*, (2001) reported that the greatest dry weight of shoot were obtained at inflorescence initiation but subsequently decreased until harvest. Brahma *et al.*, (2002) pointed out that the highest percentage of N, P and K in leaves were 3.9%, 0.44% and 2.75%, respectively and ascorbic acid content of the head were 128.05 mg/100g .

Vagen (2003) mentioned that rapid N uptake in broccoli approximately starts when the small heads just visible to the eye. Choudhary and Choudhary (2005) reported that ascorbic acid content of heads were not much influenced by the N and P treatments, but dry weight was influenced significantly by both levels of N and P.

Combination of chemical, organic or biofertilizers fertilizers supplied to broccoli plants resulted higher chemical components in leaves and heads comparing with application of organic or chemical fertilizers alone. These results were confirmed by Machendran and Kumar (1997) they reported that the highest ascorbic acid content of cabbage were produced by applied 75% of the recommended rate of NPK combined with vermicompost and digested organic supplement. Suchorska – Orowska, (1998), Morselli *et al.*, (1999) and Perin *et al.*, (2004) found that the dry matter, sugar content of heads and uptake of N, P and K of cabbage, cauliflower or broccoli were significantly increased by organomineral fertilizers applied compared to chemical fertilizer.

The enhancing effect of biofertilizers, i.e. yeast, *Pseudomonas* and *Bacillus* (phosphate solubilizing bacteria), on organic chemical component may be attributed to increase phosphatase activity, P availability or/and producing growth regulating hormones. The use of biofertilizers may have an additional benefits such as nitrogen fixation, mobilizing phosphate and micronutrients through the production of organic acids and lowering soil pH (Saber, 1993). In this respect, Hewedy (1999) stated that, inoculation of tomato plants with mixture of some biofertilizers (phosphorieren, rhizobacterien and microbien) increased the availability of some nutrients which could be reflected on plants uptake and its content from these nutrients. Nanjian *et al.*, (2001) mentioned that increases in the level of amino acids and phenols were observed by inoculation with VAM (Vesicular arbuscular mycorrhiza) in tomato plants. The levels of peroxidase, phenol oxidase and phenylalanine ammonia – lyase increased with treated plants with VAM. Leaves and roots of treated plants with VAM had higher amount of cytokinins. In addition, microorganisms can secrete growth promoting substances, e.g., gibberellins, cytokinins and auxins (Brown, 1972). Abou-Hussein *et al.*, (2002) found that applying compost with chicken manure and biofertilizer, increased the percentage of nutrients in potato leaves (N, P, K and Ca) and total carbohydrates. On the other hand, using the chicken manure with biofertilizer increased nitrate content in potato tubers. Hanafy Ahmed *et al.*, (2002) working on lettuce, suggested that the use of rhizobactrein, nitrobein,

microbein and biogein may enhance the use of nitrate transformation with the available carbon into plant growth compounds which increased plant growth by increasing the organic components (total sugars, free amino acids and soluble phenols). Moreover, Chaterjee *et al.*, (2005) evaluated the effects of different organic amendments and biofertilizer (Azotobacter, vesicular arbuscular mycorrhiza and potash mobilizer. Poultry manure + biofertilizer produced broccoli heads with the highest ascorbic acid and reducing sugar contents.

Nitrate accumulation in plants occurs as a result of nitrate accumulation in the soil due to the activity of soil nitrifying organisms. This could be mainly due to the intensive application of nitrogen fertilizers which results in imbalancing nutritional status of the plants and consequently high nitrate accumulation as well as soil pollution. Nitrate content in broccoli heads varied significantly with the N and P rates (Karitonas, 1999 and Rydz, 2001). Nitrate concentrations in vegetables were positively correlated with N rates. As a result of addition N fertilizer to soil was the major cause for vegetables increasing their nitrate contents. With the addition of P fertilizer, yields of green cabbage and rape were increased, while those of spinach had no significant changes. The nitrate concentration was significantly decreased in green cabbage by P treatment. Different organs had different amount of nitrate accumulation, and it was found that nitrite concentration were much higher in roots, stems and petioles than in blades at any N rate. The head of broccoli had the lowest nitrate content (Dolanska, 2002 and Wang and Li, 2004).

It is important here to mention that, in the present work, there was a pronounce negative relationship between nitrate accumulation and organic compounds (total sugars, soluble phenols, free amino acids and ascorbic acid) which induce due to increasing phosphorus supply, either alone or combined with any of the different biofertilizers. In this respect, Hanafy Ahmed (1996) and Hanafy Ahmed *et al.*, (2002) suggested that simple organic molecules such as sugars, free amino acids, total soluble phenole and organic acids may act as an osmoticum for the regulation of plant osmosis. These simple organic molecules or solutes can replace nitrate in the cell vacuoles. Blom – Zandstra *et al.*, (1988) suggested that the availability of sugars might affect the need for nitrate as an osmoticum. The same authors working on lettuce, pointing out that NO₃ accumulation was inversely related to the accumulation of sugars and organic acids. Furthermore, Muller and Touraine (1992) and Imsande and Touraine (1994) mentioned that increasing the concentration of certain amino acids in the phloem sap causes an inhibition of nitrates uptake.

Biofertilizers are prepared nowadays from different microorganisms having a define role in the transformations of nutrients from the soil the plant roots. The effect of plant inoculation with different types of soil microorganism has become recently new methods to reduce the amount of inorganic complete fertilizers.

Finally it could be concluded that total yield increased due to increasing of top head and side heads of plant, which was a reflection for optimum plant growth. The best yield and quality were obtained in the present

study with applying 10 m³ sheep manure plus 20 Kg P₂O₅/fed. and inoculated plants with *Bacillus megatherium* or *Pseudomonas aeruginosa*.

REFERENCES

- A. O. A. C. (1980). Official Methods of Analysis. Association of Official Analytical Chemists Washington D. C. 13th ed.
- Abou-Hussein, S. D.; El-Oksh, L. and El-Shorbagy, A. (2002). Effect of chicken manure, compost and bio fertilizers on vegetative growth, tuber characteristics and yield of potato crop. *Egypt. J. Hort.*, 29 (1): 135-149.
- Alian, F. S. (2005). Response of globe artichoke to some organic manure and mineral nitrogen levels and biostimulant treatments under sandy soil condition. Ph.D. Thesis, Fac. of Agric. ,Cairo Univ.
- Bahadur, A.; Singh, J. and Singh, K. P. (2005). Response of cabbage to organic manures and biofertilizers. *Indian Journal of Horticulture*, 61 (3): 278-279.
- Belec, C.; Villeneuve, S.; Coulombe, J. and Tremblay, N. (2001). Influence of nitrogen fertilization on yield, hollow stem incidence and sap nitrate concentration in broccoli. *Canadian Journal of Plant Science*, 81 (4): 765-772.
- Bhardwai, M. L.; Harender, R. and Koul , B. L. (2000). Yield response and economics of organic sources of nutrients as substitute to inorganic sources in tomato and cauliflower. *Indian Journal of Agricultural Sciences* , 70 (10): 653-656.
- Blom – Zandstra, M.; Lampe, J. E. M. and Ammerlan, F.H.M. (1988). C and N utilization of two lettuce genotypes during growth under non-varying light conditions and after changing the light intensity. *Physiol. Plant.*, 74 (1): 147-153.
- Bowen, P. A.; Zebarth, B. J. and Toivonen, P. N. A. (1999) . Dynamics of nitrogen and dry matter partitioning and accumulation in broccoli in relation to extractable soil inorganic nitrogen. *Canadian Journal of Plant Science* , 79 (2): 277-286.
- Brahma, S.; Phookan, D. B. and Gautam, B. P. (2002). Effect of nitrogen, phosphorus and potassium on growth of broccoli. *Journal of the Agricultural Science Society of North East India*, 15 (1): 104-106.
- Brown, J. E.; Dangler, J. M. and Gilliam.T.Z. (1994). Comparison of broiler litter and inorganic nitrogen, phosphorus and potassium for double – cropped sweet corn and broccoli. *Journal of Plant Nutrition* , 17(5): 859-867.
- Brown, M. E. (1972). Plant growth substances produced by microorganisms of soil rhizosphere. *J. Appl. Bacteriol.*, 35:443-451.
- Cataldo, D. A.; Harnon, M.; Schrader, L. E. and Youngs, V. L. (1975). Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Commune in Soil Science and Plant Analysis*, 6 (1): 71-80.

- Chapman, H. D. and Parker, F. (1961). Methods of analysis for soil, plant and water. *Journal of Plant Nutrition* , 22 (1): 121-128.
- Chatterjee, B.; Ghanti, P.; Thapa, U and Tripathy, P. (2005). Effect of organic nutrition in sprouting broccoli. *Vegetable Science* , 32 (1): 51-54.
- Choudhary, R. K. and Choudhary, D.W. (2005). Effect of different levels of nitrogen and phosphorus on growth, yield and quality of cabbage. *Haryana Journal of Horticultural Sciences*, 34 (1/2): 145-146.
- Craddock, V. M. (1983). Nitrosamine and human cancer. Proof of an association ? *Nature*, 306: 638-641.
- Darwesh, F. M. (2002). Effect of different sources and levels of fertilizers on growth, yield and quality of tomato. Ph.D. Thesis, Fac. of Agric., Cairo Univ., Egypt.
- Decoteau, D. R. (2000). *Vegetable Crops*. Prentice-Hall, Inc upper saddle River, New Jersey, 464 p.
- Dolanska, L. (2002). Study of nitrate contents in broccoli cultivated in condition of different nitrogen doses. *Collection of Scientific papers. Faculty of Agriculture in Ceske Budejovice Series for Crop Sciences*, 19 (2) : 85-89.
- Dubois, M.; Smith, F.; Gilles, K. A.; Hamilton J. K. and Rebers, P. A. (1956). Colorimetric method for determination of sugars and related substances. *Anal. Chem.*, 28 (3): 350-356.
- Dufault, R. J. (1988). Nitrogen and phosphorus requirements for green house broccoli production. *Hort. Science* , 23 (3):576-578.
- Edmont, J. B. ; Sun, T. L. and Inderws, F. A. S. (1981). *Fundamental of Horticulture* Published by Tate. MC. Grawtill Publishing Co. Limited, Indian.
- El-Magd, M. M. A.; Mohamed, H. A. and Fawzy, Z. F. (2005). Relationship between growth, yield of broccoli with increasing, N, P or K ratio in a mixture of NPK fertilizers. *Annals of Agricultural Science Moshtohor*, 43 (2): 791-805.
- Fortunato, M. I. and Damato, A. (2000). Sowing times and ontogenetic cycle in two early cultivars of cima diraba (*Brassica rapa*). *Acta Hort.*, 533: 291-297.
- Gauss, J. and Taylor, G. A. (1996). A morphological study on the time of reproductive differentiation of apical meristem of broccoli. *Amer. Soc. Hort. Sci.*, 94:105-108.
- Hanafy Ahmed, A. H. (1996). Physiological studies on tip burn and nitrate accumulation in lettuce plant. *J. Agric. Sci. Mansoura Univ.*, 21 (11): 3971-3994.
- Hanafy- Ahmed, A. H.; Mishriky, J. F. and Khalil, M. K. (2002). Reducing nitrate accumulation in lettuce plants by using different biofertilizers. *Annals Agric. Ain Shams Univ.*, Cairo, 47 (1): 27-41.
- Hewedy, A. M. (1999). Influence of single and multibacterial fertilizer on the growth and fruit yield of tomato. *Egypt. J. Appl. Sci.*, 14 (7): 508-525.
- Hsieh, C. and Hsu ,A. (1995). Effect of continuous use of organic manures on the growth and yield of vegetable soyabean and cabbage. *Hulletin of Taichung District Agricultural Improvement Station* , 46:1-10.

- Imsande, J. and Touraine, B. (1994). N demand and the regulating of nitrate uptake. *Plant Physiol.*, 105 : 3-7.
- Inoune, T.; Imagawa, M. and Nishiwaki, K. (1996). The effects of the continuous application of rice straw compost on the nutrient uptake of vegetable in mineral soil upland fields. *Research Bulletin of the Aichi ken Agricultural Research Center*, 28: 157-164.
- Jackson, M. L. (1962). *Soil Chemical Analysis*. Contable and Company Ltd., London, 498 pp.
- Johnson, I. T. (2000). Brassica vegetables and human health: glucosinolates in the food chain. *Acta Hort.*, 539:93-104.
- Kamaiyama, K.; Fujiwara, S. and Funahashi, H. (1995). Effect of successive application of cow manure compost on growth of crops and the chemical properties, of the soil. *Bulletin of the Agricultural Research Institute of Kanagama Prefecture*, 136: 31-42.
- Karitonas, R. (1999). Optimization of mineral nutrition for broccoli. *Sodininkyste-ir-Darzininkyste*, 18 (3): 122-128.
- Kumar, P. and Sharma, S. K. (2001). Performance of different sprouting broccoli cultivars under different combination of nitrogen, phosphorus and potassium. *Indian Journal of Agricultural – Sciences*, 71 (8): 558-560.
- Linder, R. C. (1994). Rapid analytical methods for some of the more common inorganic constituents of plant tissues. *Plant Physiol.*, 19 (1) : 76-79.
- Lu, H.; Xu, X. and Zhang J. (1997). Study of features of nutrient absorption and dry matter accumulation in broccoli. *Acta Agriculture Shanghai*, 13 (4): 47-50.
- Machendran, P. P. and Kumar, N. (1997). Effect of organic manures on cabbage. *South Indian Horticulture*, 45 (5/6): 240-243.
- Manivannan, M. I . and Singh, J. P. (2004). Effect of biofertilizers on the growth and yield of sprouting broccoli under Allahabad agro-climatic condition. *Bioved* , 15 (1/2): 33-36.
- Moore, S. and Stein, W. H. (1954). A modified ninhydrin reagent for the photometric determination of amino acids and related compounds. *J. Biol. Chem.*, 211: 907-913.
- Morselli, T. B.; Fernandes, H. S. and martins, S. R. (1999). Response of cabbage and cauliflower to application of vermicomposts in the liquid form. *Revista Cientifica Rural*, 4 (2): 24-28.
- Muller, B. and Touraine, B. (1992). Inhibition NO₃ uptake by various phloem-translocated amino acids in soybean seedlings. *J. Exp. Bot.*, 43: 617-623.
- Nanjian, R.; Mariappan, G. and Velusamy, S. (2001). Biological control of fusarium wilt of tomato by VA mycorrhizal fungus. *Glomus fasciculaum*. *Bull. Oil. Srop.*, 24 (3): 33-36.
- Nkoa, R.; Coulombe, J. and Desjardins, Y. (2001). Towards optimtion of growth via nutrient supply phasing: nitrogen supply phasing increases broccoli growth and yield. *Journal of Exp. Bot.*, 52 (357): 821-827.
- Perez-Murcia, M. D.; Moral, R. and Moreno, C. J. (2006). Use of composted sewage sludge in growth media for broccoli. *Bioresource-Technology*, 97 (1): 123-130.

- Perin, A.; Santos, R. H. and Urquiaga, S. (2004). Residual effect of green manures on broccoli yield cultivated in succession to corn. *Ciencia Rural*, 34 (6): 1739-1745.
- Prabhakar, B. S.; Srinivas, K. and Subramanian, T. R. (1986). Phosphorus fertilization in sequence cropping of vegetables. *Indian Journal of Agricultural Sciences*, 56 (10): 710-716.
- Raghothama, K. G. (1999). Phosphate acquisition. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 50: 665-693.
- Rincon, L.; Saez, J. and Perez, J. A. (1999). Growth and nutrient absorption of broccoli. *Investigation Agraria Produccion Proteccion Vegetables*, 14 (1/2): 225-236.
- Rydz, A. (2001). The effect of foliar nutrition with urea on yield quality of broccoli. *Vegetable Crops Research Bulletin*, 54 (1): 61-64.
- Saber, M. S. M. (1993). A multi-strain biofertilizer. *Proc. The sixth International Symposium on Nitrogen Fixation with Non-Legumes*, pp. 138-145. Ismailia, Egypt, 6-10 September.
- Sharma, K. C. (2000). Influence of integrated nutrient management on yield and economics in broccoli under cold temperature conditions. *Vegetable Science*, 27 (1): 62-63.
- Sharma, S. K.; Rajender, S. and Korla, B. N. (2002). Effect of nitrogen and phosphorus on growth and seed yield of sprouting broccoli. *Horticultural Journal*, 15 (2): 87-90.
- Snedecor, G. W. and Cochran, W. G. (1980). "Statistical Methods", 7th Ed. The Iowa State Univ. Press, Ames, Iowa.
- Subba Rao, S. (1993). *Biofertilizers in Agriculture* (3rd Ed., Oxford & BD Publishing Co Ltd., New Delhi, Bombay, Calcutta, 242 p.
- Subhan, A. (1989). The effect of maize stalk compost, rice straw compost and sheep manure on the yield of cabbage. *Buletin Penelitian Horikultura*, 17 (4): 80-91.
- Suchorska – Orowska, J. (1998). Evaluation of fine brown coal and brown coal ash as organomineral fertilizers for vegetables. *Rozprawy Akademia-Rolniczaw Szczecinie*, 184 :92-102.
- Swain. T. and Hillis, W. F. (1959). The quantitative analysis of phenolic constituent. *J. Sci.; Food Agric.*, 10: 63-69.
- Vagen, I. M. (2003). Nitrogen uptake in a broccoli crop. 1: Nitrogen dynamics on a relative time scale. *Acta Horticulturae*, 627: 195-202.
- Van-Schouwenburg, J. C. H. and Walinga, I. (1978). *Methods of analysis of plant material*. Agric. Univ., Wagejnen – Netherlands.
- Wang, G.; Zhang, C. and Chun, Z. (1997). Effect of nitrogen, phosphorus and potassium fertilizer on yield and physiology target of broccoli. *China Vegetables*, 36(1) : 14-17.
- Wang, Z. and Li, S. (2004). Effects of nitrogen and phosphorus fertilization on plant growth and nitrate accumulation in vegetables. *Journal of Plant Nutrition*, 27 (3) : 539-556.
- Wright, M. J. and Davison, K. L. (1964). Nitrate accumulation in crops and nitrate poisoning of animals. *Advances in Agronomy*, 16:197-247.

تأثير التسميد بمخلفات الغنم ومستويات الفوسفات والسماذ الكيماوي والحيوي
على النمو والمحصول وتراكم النترات والمحتوي الكيماوي للبروكولي.
امل محمد فراج*، أحمد حسين حنفي أحمد** و هناء فتوح أحمد**
* قسم الخضراة الزراعة جامعة القاهرة.
** فرع فسيولوجي النبات قسم النبات كلية الزراعة جامعة القاهرة.

اجريت تجربتين حقليتين على نباتات البروكلي صنف برميم لدراسة تأثير سماذ الغنم بمعدل ١٠ م^٢ للفدان (٤,٢٨ طن) وثلاث مستويات من السماذ الفوسفاتي (صفر، ١٠، ٢٠ كجم فو٢أه / فدان) والسماذ كيماوي الموصى به (٤٠ ن، ٣٠ فو٢أه، ٥٠ ب٢ أ كجم / فدان). الشتلات لقت بالمخصبات الحيوية وهي الخميرة والبايسيلس وبسيدموناس وكذلك عدم التلقيح وتأثرها على صفات النمو الخضري عند بداية تكوين الرؤوس والرؤوس الجانبية وايضا عدد الأيام حتى بداية ظهور الأقراص، كمية المحصول ومكوناته والمحتوي الكيماوي للأوراق والأقراص. أظهرت النتائج ان استعمال سماذ الغنم + ٢٠ كجم فو٢أه يليه سماذ الغنم + ١٠ كجم فو٢أه أعطى أكبر القيم لصفات طول النبات، وزن النبات، والساق والأوراق وعدد الأوراق للنبات مقارنة باستخدام السماذ الكيماوي الموصى به، كذلك متوسط وزن القرص ووزن الأقراص الجانبية للنبات والمحصول الكلي وأيضا النسبة المئوية النتروجين والفوسفور والبوتاسيوم والمادة الجافة وتركيز السكريات الكلية والفينولات والأحماض الامينية الحرة الكلية في الأوراق والأقراص. لم يتأثر عدد الأيام من الشتل حتى بداية ظهور القرص الرئيسي بأنواع أو مستويات التسميد. استعمال سماذ الغنم + ١٠ كجم فو٢أه / فدان أعطى نتائج أفضل من استعمال السماذ الكيماوي الموصى به لمعظم الصفات. محتوى الأقراص من حمض الاسكوربيك كان أعلى عند استعمال سماذ الغنم + ١٠ كجم فو٢أه / فدان. كان محتوى الأقراص من النترات أعلى باستعمال السماذ الكيماوي. استعمال المخصبات الحيوية تسببت في زيادة لصفات النمو الخضري والمحصول ومكوناته وكذلك المكونات الكيماوية للأوراق والرؤوس مقارنة بعدم استعمال المخصبات الحيوية. بسيدموناس تسببت في أكبر قيم محصول الأقراص للفدان في الموسم الأول، أيضا محصول الأقراص الجانبية والمحصول الكلي للفدان والسكريات الكلية والأحماض الامينية الكلية في الأوراق في كلا الموسمين. الخميرة أعطت أعلى القيم لمحصول الأقراص للفدان والأحماض الامينية الكلية في الأقراص في الموسم الثاني. كان أعلى تراكم للنترات في الأقراص عند تخصيب النباتات بالخميرة بينما استعمال الباسيلس تسببت في أقل مقدار. بكتريا الباسيلس تسببت في أعلى قيم الاحماض الامينية الحرة في الأقراص في الموسم الاول، وحمض الاسكوربيك في الأقراص من الموسمين. أعلى محصول كلى تم الحصول عليه بتسميد النباتات بسماذ الغنم + ٢٠ كجم فو٢أه وتلقيحها بالبسيدموناس في الموسم الاول والبايسيلس في الموسم الثاني.