

## **GIZA 2000, A NEW EGYPTIAN BARLEY VARIETY FOR NEWLY RECLAIMED LANDS AND RAINFED AREAS**

**Ahmed, I. A.; R. A. Abo - El-Enin; A. A. Ei-Hag; Kh.A. Mcustafa; M.A. El-Moselhy; A.M.O. El-Bawab; H.A.T. Mahfouz\*; K.A. Amer; M.M. Abd El-Hamed and R.A. Rizk\*\***

Barley Res. Dept., Field Crops Res. Inst., Agric. Res. Center, Giza, Egypt

\* Agric. Genet. Eng. Res. Inst., Agric. Res. Center, ARC, Giza, Egypt

\*\* Plant Path. Res. Inst., Agric. Res. Center, ARC, Giza, Egypt

### **ABSTRACT**

This study reports the development and characteristics of Giza 2000, a superior new barley variety for the newly reclaimed lands and rainfed areas at the North Coast region. It was produced from a cross between the local variety Giza 121 and the Line 366/13/1 which was made in 1978/1979 season. In developing this variety, the check cultivars Giza 123, Giza 124, Giza 125 and Giza 126 were compared with Giza 2000 and two promising lines as well as the regional check Rihane-03. For this purpose, 39 yield trials were conducted from 1998/1999 to 2001/2002 growing season under rainfed conditions as well as under newly reclaimed lands.

The average yield of the new cultivar Giza 2000 recorded 3.51 ardab/feddan and significantly outyielded Giza 126 under rainfed conditions with an average increase of 0.57 ardab/feddan, i.e. 19.39%. In the newly reclaimed lands, the average yield of the new cultivar Giza 2000 significantly exceeded the average of national check Giza 123 by 2.46 ardab/feddan (17.25%). The new variety combines the good characteristics of its parents which including high yielding ability, early maturity, moderate resistance to leaf rust and resistance to powdery mildew and net blotch. Molecular fingerprinting utilizing PCR with 38 random 10mer primers was done for the new variety in order to be used as an appraisal when commercializing this variety and preserve the breeder's rights. A number of 115 reproducible and repeatable bands were recovered during PCR. Usefulness of this environment-independent molecular approach was discussed.

Seed multiplication of Giza 2000 was started since 2000/2001 growing season to produce certified seeds which will be available to cover the barley production area in Egypt.

**Key words:** New barley cultivar, Giza 2000, yield, yield stability, fingerprint.

### **INTRODUCTION**

Barley is the main crop grown in a large scale in North Coast of Egypt. It is also the main crop in the newly reclaimed land and in soils with chemical problems (saline soils) or where the irrigation water is limited. The total area of barley in Egypt fluctuated year after year according to the amount and distribution of rainfall. The barley cultivated production area in the Nile Valley has decreased, especially at locations where soil and irrigation is suitable for growing other strategic crops, such as wheat. On the other hand, the barley production area has increased in the newly reclaimed lands under different irrigation systems. The harvested area reached 99,356 hectare in 2001/2002. Barley yields have increased gradually over the past 50 years from 2.53 in the sixties to 2.98 ton/ha in 2000/2001 season. The increase in barley productivity

was mainly due to the release of improved barley varieties, which started when the varieties Sahrawy, Bonus, Giza 117 and Giza 118 were released in 1955, 1956, 1958 and 1963, respectively. All these varieties were resistant to the main barley diseases and characterized by high yielding ability. Later on, crossing work was further intensified by the introduction of useful resources, consequently, Giza 119, Giza 121, CC 89 and CC 163 were developed in 1973, 1975, 1977 and 1977, respectively. It should be clearly stated that breeding work achieved forward step in the past 15 years by releasing the varieties Giza 123 (for soil salinity and new reclaimed area), Giza 124 (for heat stress), Giza 125, and Giza 126 (for drought stress condition). These varieties were characterized by their high yielding ability and exhibited good performance under different environmental stresses.

Most progress in cultivar identification, so far, has relied only on a phenotypic assay of genotype, i.e. morphological characteristics that require extensive observations of individuals (Wrigley *et al.*, 1987). Factors like the environment, multigenic and quantitative inheritance or partial and complete dominance virtually affect gene expression. However, DNA-based genetic markers have been extensively integrated into several plant systems and are expected to play a very important role in the future of molecular genetics and plant breeding. Over the last decade, the revolution of polymerase chain reaction (PCR) technology has been initiated as a novel genetic assay based on selective DNA amplification (Saiki *et al.*, 1988; Krawets, 1989; Innis *et al.*, 1990). This assay relies on the enzymatic amplification of small DNA fragments using single arbitrary oligonucleotide primers (usually 10 mers) for cultivar identification.

The present work represents the procedures followed in developing the new barley variety Giza 2000, which was characterized by its wide adaptability and its high yielding ability under different environmental conditions.

## **MATERIALS AND METHODS**

Bahteem 52, Giza 117, and Giza 118 (Beecher) and Giza 121 were local varieties adapted to Egyptian conditions, but their popularity has declined because of their susceptibility to leaf rust, net blotch, and low yielding ability. FAO 86 was introduced from USDA and showed a good agronomic performance and disease resistance. During 1969/1970 growing season, the cross between Giza 117 and Bahteem 52 was performed at Giza Agric. Res. Center. In the same season, another cross was made between Giza 118 and FAO 86. These two crosses were crossed together in 1970/1971 and produced the cross Giza 117/Bahteem 52//Giza 118/FAO 86. The purpose of this cross was to combine the good characteristics of the four parents. The cross Giza 117/Bahteem 52//Giza 118/FAO 86 was crossed with the local cultivar Giza 121 at Giza Agricultural Research Station, Egypt in 1978/1979 growing season. The segregating material was handled according to the pedigree method at Giza and Sakha Research Stations where it was exposed to severe natural infection of leaf rust and net blotch.

The cross Giza 117/Bahteem 52//Giza 118/FAO 86\*Giza 121 was compared along with promising material in early yield trials. As a result of

these micro trials, the cross was promoted for further evaluation in Advanced Yield Trials compared with the commercial varieties (Giza 123, Giza 124, Giza 125 and Giza 126) along with Rihane-03 (long term regional check cultivar) and two promising lines (Table 1). For this purpose, a series of 15 and 24 yield trials were conducted, under rainfed conditions and newly reclaimed lands, respectively, in a randomized complete block design with 3 replications during the period from 1998/1999 to 2001/2002. Agricultural practices for barley were applied as recommended in each region.

Statistical analysis of data obtained was made using the methods outlined by Gomez and Gomez (1984). The stability parameters suggested by Eberhart and Russell (1966) were estimated for grain yield of the tested genotypes over all years and locations under study. One stability parameter was estimated as the linear regression coefficient *b* of a genotype mean on the average of all genotypes in the particular environment. The other stability parameter was deviation from regression  $S^2d$  for each genotype.

**Table (1): Name/cross and pedigree of the tested genotypes.**

Genotype name	Pedigree
Rihane 'S'	Long term check variety introduced from ICARDA
Giza 123	Giza 117/FAO 86
Giza 124	Giza 117/Bahteem 52// Giza 118/FAO 86
Giza 125	Sister line to Giza 124
Giza 126	Baladi Bahteem/SD 729-Por 12762-BC
Line 1	(Giza 121 x Chaaran-01/ Deir Alla 106/3/Asse/Alhs//Apm)
Line 2	(MAF 102/Volla//WW319 x Giza 119)
Giza 2000	Giza 117/Bahteem 52// Giza 118/FAO 86*Giza 121

#### **DNA isolation**

Leaf sample from 7-days-old seedlings of Giza 2000 was collected from five plants to saturate polymorphism within the variety and instantly frozen in liquid nitrogen. DNA was extracted from 0.5 g of fresh tissue by the modified procedure of Gawel and Jarrett (1991). DNA concentration was measured by UV-spectrophotometer at a wavelength of 260 nm.

#### **PCR conditions**

Thirty eight arbitrary 10 mer primers (primers A04, B01, B02, B05, B06, B07, B08, B10, B11, B12, B13, B14, B15, B16, B18, C01, C05, C08, C10, C19, D07, D09, D10, D20, E07, E09, E19, O02, O05, O09, O10, O12, O18, O20, Z08, Z10, Z11 and Z13 from Operon Technologies Inc., Alameda, CA 94501) were used for PCR based on the protocol of Williams et al. (1990). The reaction conditions were optimized and mixtures (37.5 µl total volume) consisted of 10 mM Tris-HCL, pH 8.8 at 25°C, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, nucleotides dATP, dCTP, dGTP and TTP (0.2 mM each), 0.2 µM primer, 50 ng template DNA and 1.5 units of Taq DNA polymerase (Promega). Amplifications were carried out in a thermocycler (Berkin Elmer) programmed for 37 cycles of 45 sec at 94°C, 50 sec at 36°C, 1 min at 72°C and ended with 8 min at 72°C.

### **Gel electrophoresis**

Agarose (1.2%) gel electrophoresis was used in this study according to Bahieldin and Ahmed (1994). A 1 kb plus DNA ladder was used as a standard. The run was performed for one hour at 100V in Pharmacia submarine (20 cm X 20 cm). Bands were detected, scored for molecular weights and photographed by Gel Documentation System. PCR was repeated twice for each primer and products that were generated at least twice were considered reproducible, while those generated only once were considered artifacts and eliminated from the fingerprint. Band sizes less than 100 or over 5000 bp were also eliminated from the fingerprint that are usually unrepeatably.

## **RESULTS AND DISCUSSION**

### **Yield performance under rainfed conditions**

A series of 15 yield trials were conducted at Mersa Matrouh (El-Mathani and El-Negela), North Sinai (El-Goura and Raffah) and Sakha and El-Noubaria (irrigated only once at sowing time) under rainfed conditions during the period of 1999/2000 to 2001/2002 to evaluate eight barley cultivars and lines. Drought is the main environmental problem occurred in the north coastal areas. Table (2) shows the yield performances (ardab/feddan) of the tested genotypes under the 15 yield trials. Highly significant differences in grain yield were detected between genotypes as well as all interactions.

It is evident from Table (2) that grain yield significantly decreased from 4.26 ardab/feddan (at Raffah) to 1.69 ardab/feddan at El-Mathani (severe drought stress). The new cultivar Giza 2000 exceeded the national check cultivar Giza 126 in all yield trials. In nine yield trials out of the 15, the yield of the new cultivar (Giza 2000) exceeded significantly the national check cultivar (Giza 126). Table (2) also presents the average yield in ardab/feddan for the tested lines and the check combined over years and locations. The combined analysis of tested genotypes over years and locations showed that Giza 2000 significantly exceeded Giza 126 in grain yield, with an average increase of 0.57 ardab/feddan, i.e. 19.4%. Under severe drought stress occurred at both El-Mathani and El-Goura, the new cultivar exceeded significantly the national check cultivar Giza 126 by about 0.38 and 0.41 ardab/feddan, i.e. 23.8 and 23.6%, respectively.

### **Yield performance in newly reclaimed lands**

Table (3) shows the average grain yield in ardab/feddan for the tested genotypes in 24 experimental yield trials representing different agro-climatic zones in Egypt during the period starting from 1998/1999 to 2001/2002. The experimental sites included El-Noubaria (calcareous soils), Sakha (diseases problems), El-Gemmeiza (optimum conditions), Mallowy (heat stress and aphids problems) and El-Hamoul, and El-Serw (soil salinity). The 24 yield trials performed included four barley cultivars (Giza 123, Giza 124, Giza 125 and Giza 126), three promising lines and the long term regional check cultivar (Rihane-03), Table (1). Analysis of variance presented in Table (3) shows highly significant differences among years, locations, genotypes and all types of interactions.

**Table (2): Average yield in ardab/feddan for the tested genotypes evaluated under rainfed conditions during 1999/2000 through 2001/2002.**

Season and Variety			Location			
1999/2000	El-Mathani	El-Negela	El-Goura	Raffah	Sakha*	Mean
Rihane-03	1.48	2.15	1.79	4.75	3.42	2.72
Giza 123	1.82	2.31	1.78	3.37	3.69	2.59
Giza 124	2.19	2.07	1.72	4.55	2.36	2.58
Giza 125	2.14	2.66	2.02	4.25	2.82	2.76
Giza 126	1.82	2.75	1.88	4.08	3.83	2.87
Line 1	2.08	1.98	1.72	5.05	4.15	3.00
Line 2	1.85	1.66	2.00	3.89	3.27	2.54
Giza 2000	2.55	2.93	2.45	4.97	4.68	3.52
Mean	1.99	2.30	1.92	4.36	3.53	2.82
2000/2001	El-Mathani	El-Negela	El-Goura	Raffah	Noubaria*	Mean
Rihane-03	1.65	2.70	1.75	3.21	4.44	2.75
Giza 123	1.76	2.36	1.68	4.06	5.62	3.10
Giza 124	1.57	2.47	1.87	4.16	5.07	3.03
Giza 125	1.54	2.76	1.42	3.93	5.01	2.93
Giza 126	1.55	2.38	1.78	4.08	4.24	2.80
Line 1	1.53	2.73	1.51	3.38	3.38	2.51
Line 2	1.46	2.69	1.64	4.25	5.57	3.12
Giza 2000	1.62	2.94	2.00	4.22	5.92	3.34
Mean	1.59	2.63	1.70	3.91	4.91	2.95
2001/2002	El-Mathani	El-Negela	El-Goura	Raffah	Noubaria*	Mean
Rihane-03	1.42	2.61	1.61	3.04	5.69	2.87
Giza 123	1.25	2.27	1.95	4.51	5.74	3.14
Giza 124	1.61	2.29	1.43	4.81	5.42	3.11
Giza 125	1.30	2.02	1.87	4.82	4.61	2.92
Giza 126	1.44	2.20	1.56	4.58	5.96	3.15
Line 1	1.61	2.51	2.01	4.27	5.53	3.19
Line 2	1.43	2.39	1.66	4.28	5.15	2.98
Giza 2000	1.78	2.85	1.97	5.77	6.04	3.68
Mean	1.48	2.39	1.76	4.51	5.52	3.13
Average of genotypes over years						
Rihane-03	1.51	2.89	1.71	3.67	4.52	2.78
Giza 123	1.61	2.31	1.80	3.80	5.02	2.95
Giza 124	1.79	2.28	1.67	4.51	4.29	2.91
Giza 125	1.66	2.45	1.77	4.33	4.14	2.87
Giza 126	1.60	2.44	1.73	4.25	4.68	2.94
Line 1	1.74	2.41	1.75	4.23	4.35	2.90
Line 2	1.58	2.25	1.77	4.14	4.66	2.86
Giza 2000	1.98	2.91	2.14	4.99	5.55	3.51
Mean	1.69	2.44	1.79	4.26	4.65	
L.S.D. at						
for Year						
Y	0.13	0.13	0.13	0.13	0.13	0.13
Location	L	0.29	0.32	0.29	0.29	0.29
Location (Year)LY		0.34	0.39	0.34	0.34	0.34
Genotypes	G	0.21	0.29	0.21	0.21	0.21
YG		0.37	0.49	0.37	0.37	0.37
LG		0.54	0.67	0.54	0.54	0.54
LGY		0.83	1.09	0.83	0.83	0.83

\* Supplied with sowing irrigation and left under rainfed conditions.

**Table (3): Average yield in ardab/feddan for the tested genotypes evaluated under six different locations during 1998/1999 through 2001/2002.**

Genotype	Location						Mean
	El-Gemmeiza	Sakha	Mallawy	El-Noubaria	El-Hamoul*	El-Serw*	
<b>1998/1999</b>							
Rihane-03	16.26	18.81	20.82	17.27	8.78	14.36	15.72
Giza 123	17.97	19.69	19.99	14.14	6.75	14.89	15.57
Giza 124	17.53	16.45	23.46	17.23	10.36	13.76	16.47
Giza 125	14.43	20.16	20.22	19.90	7.42	16.51	16.44
Giza 126	16.51	18.73	21.42	20.56	9.57	16.51	17.22
Line 1	17.65	18.31	22.14	18.18	7.83	15.32	16.57
Line 2	20.07	19.75	19.99	13.49	7.80	13.82	15.82
Giza 2000	20.31	22.50	23.99	20.87	10.34	16.28	19.05
Mean	17.59	19.05	21.50	17.71	8.61	15.18	16.61
<b>1999/2000</b>							
Rihane-03	17.57	16.34	20.94	13.88	9.83	12.20	15.14
Giza 123	20.44	20.22	21.06	18.23	7.42	11.42	16.47
Giza 124	24.77	17.77	19.39	14.40	8.86	10.05	15.87
Giza 125	23.07	17.83	21.54	17.50	9.73	11.25	16.65
Giza 126	22.92	19.81	20.64	15.13	9.93	11.49	16.65
Line 1	22.41	17.23	18.31	20.27	6.94	9.93	15.85
Line 2	23.34	16.51	17.83	18.44	7.78	9.16	15.51
Giza 2000	24.99	19.75	19.75	18.72	7.78	12.32	17.22
Mean	22.44	18.18	19.93	17.07	8.42	10.98	16.17
<b>2000/2001</b>							
Rihane-03	22.21	16.63	14.96	14.85	4.19	10.17	13.84
Giza 123	20.91	13.88	13.28	15.35	2.87	8.88	12.53
Giza 124	19.21	14.12	16.39	17.78	5.15	10.29	13.82
Giza 125	24.24	16.75	18.91	17.19	5.39	10.53	15.50
Giza 126	19.84	15.20	15.80	17.83	3.05	8.02	13.29
Line 1	18.68	15.68	15.92	15.99	4.07	9.57	13.32
Line 2	22.93	13.76	15.80	15.56	3.95	8.14	13.36
Giza 2000	23.59	18.79	17.59	19.03	5.51	10.77	15.88
Mean	21.45	15.60	16.08	16.70	4.27	9.54	13.94
<b>2001/2002</b>							
Rihane-03	16.63	14.77	16.83	13.52	6.80	7.67	12.70
Giza 123	17.60	15.96	18.12	11.42	5.06	8.73	12.48
Giza 124	17.95	14.34	17.57	11.30	7.23	7.78	12.70
Giza 125	18.32	16.24	18.00	12.32	6.39	7.35	13.10
Giza 126	17.58	15.94	17.17	10.65	6.69	8.31	12.72
Line 1	17.43	15.19	16.72	11.33	6.55	7.78	12.50
Line 2	19.68	14.84	15.90	13.40	5.79	8.09	12.95
Giza 2000	20.43	18.27	18.35	15.43	7.31	5.95	14.74
Mean	18.20	15.69	17.08	12.42	6.44	8.08	12.99
<b>Average of genotypes over seasons</b>							
Rihane-03	18.17	16.14	18.39	14.88	7.43	11.10	14.35
Giza 123	19.23	17.44	17.61	14.79	5.53	10.98	14.26
Giza 124	19.87	15.67	19.20	15.18	7.90	10.47	14.71
Giza 125	20.02	17.75	19.67	18.73	6.98	11.41	15.43
Giza 126	19.21	17.42	18.76	16.04	7.31	11.08	14.97
Line 1	19.04	16.60	18.27	16.44	6.35	10.65	14.56
Line 2	21.51	16.22	17.38	15.22	6.33	9.80	14.41
Giza 2000	22.33	19.83	19.92	18.51	7.66	12.08	16.72
Mean	19.92	17.13	18.65	15.97	6.94	10.95	
L.S.D. at		5%	1%		5%	1%	
for Year	Y	0.49	0.64	YG	N.S	N.S	
Location	L	0.59	1.56	LG	1.68	N.S	
Location (Year)	LY	1.19	1.57	LGY	3.36	N.S	
Genotypes	G	0.69	0.90				

\* Saline soils.

The average yield (ardab/feddan) ranged from 6.94 (at El-Hamoul, saline soils) to 19.92 (at El-Gemmeiza, optimum conditions). It is clear that grain yield is seriously affected by soil salinity.

The new cultivar Giza 2000 exceeded the national check cultivar Giza 123 in all yield trials except at Sakha and Mallawy in the second season (1999/2000).

The yield of the new cultivar Giza 2000 ranged from 7.66 (at El-Hamoul) to 22.33 ardab/feddan (at El-Gemmeiza) with an average of 16.72 ardab/feddan. The respective values for Giza 123 were 5.53, 19.23 and 14.26 ardab/feddan. Out of the 24 yield trials, the yield of the new cultivar Giza 2000 significantly exceeded the national check cultivar Giza 123 only in eight yield trials. The combined analysis over years and locations showed that Giza 2000 significantly exceeded Giza 123 in grain yield, with an average increase of 2.46 ardab/feddan, i.e. 17.3%.

Table (3) also presents the average yield in ardab/feddan for the tested lines and the check combined over years and locations. The combined analysis of tested genotypes over years and locations shows that under severe drought stress occurred at both El-Mathani and El-Goura, the new cultivar significantly exceeded the national check cultivar Giza 126 by about 0.38 and 0.41 ardab/feddan i.e. 23.8 and 23.6%, respectively. It could be noticed that Giza 2000 gave better yield than the other genotypes under saline soils. It exceeded the national check Giza 123 by about 2.13 and 1.10 ardab/feddan, i.e. 27.81 and 9.11%, in respective order.

#### **Yield stability**

A knowledge of genotype x environment interactions led to successful evaluation of stable genotypes which could be used in future breeding programs. Stability parameters for grain yield of the tested genotypes were estimated by the method described by Eberhart and Russell (1966), who defined the stable genotype as the one which had a regression coefficient of 1.0 and no deviation from regression mean square. An ideal genotype would have both a high average performance across a wide range of environments plus stability.

Table (4) indicates superiority of the new cultivar Giza 2000 as compared with the national checks Giza 123 and Giza 126 under rainfed conditions, newly reclaimed lands as well as all combinations of yield trials. Furthermore, the wider range of performance (2.71-26.74 ardab/feddan) may promise better yield. Slopes of regression on environmental indices did not differ from unity ( $b=1$ ) for Giza 2000, on the mean time,  $S^2b$  value was significantly different from zero, which proved the stability of the new cultivar Giza 2000.

#### **Diseases reaction**

The results presented in Table (5) show the reaction of the tested genotypes to the major barley diseases, i.e. powdery mildew, leaf rust and net blotch. It is clear from the table that the new cultivar Giza 2000 exhibited moderate resistance to leaf rust and resistant to both powdery mildew and net blotch.

Based on performance tests and agronomic characteristics of the new line Giza 121\*Giza 117/Bahteem 52//Giza 118/FAO 86, it was recommended to be released in 2000/2001 as a new variety designated as Giza 2000. Giza 2000 could be recommended for both rainfed and newly reclaimed lands. The cultivar proved to be wide adaptable and can be used under a wide range of environments.

**Table (4): Means performances of the studied genotypes under rainfed conditions and newly reclaimed lands as well as stability parameters over all locations and years.**

Genotypes	Environment	Yield (ardab/fed)			Stability parameter <sup>1</sup>			
		Min.	Max.	Avg.	X	b	S <sup>2</sup> d	CD
Rihane-03	Rainfed <sup>2</sup>	1.22	6.09	2.78	9.90	0.92	1.18	0.96
	New Land <sup>3</sup>	3.94	23.77	14.35				
Giza 123	Rainfed	1.18	6.14	2.95	9.91	0.97	1.02	0.97
	New Land	2.71	22.53	14.26				
Giza 124	Rainfed	0.91	5.80	2.03	10.17	0.97	1.00	0.97
	New Land	4.83	25.73	14.71				
Giza 125	Rainfed	1.22	5.36	2.87	10.60	1.06	0.92	0.98
	New Land	5.05	25.94	15.43				
Giza 126	Rainfed	1.35	7.08	2.94	10.34	1.01	0.76	0.98
	New Land	2.88	24.52	14.97				
Line 1	Rainfed	1.34	6.44	2.90	10.07	0.98	0.65	0.98
	New Land	3.83	23.98	14.56				
Line 2	Rainfed	1.35	5.96	2.88	9.98	0.99	1.28	0.96
	Newly Land	3.71	24.97	14.41				
Giza 2000	Rainfed	1.52	6.48	3.51	11.64	1.09	0.38	0.99
	Newly Land	5.16	26.74	16.72				

<sup>1</sup> Calculated from a series of 39 yield trials conducted under rainfed areas and newly reclaimed lands during the period from 1998/1999 through 2001/2002.

<sup>2</sup> Calculated from a series of 15 yield trials conducted under rainfed areas during the period from 1999/2000 through 2001/2002.

<sup>3</sup> Calculated from a series of 24 yield trials conducted on newly reclaimed lands during the period from 1998/1999 through 2001/2002.

**Table (5): Reaction of the tested genotypes to the major barley diseases.**

Genotype	Powdery mildew (seedling stage)	Powdery mildew (adult stage)	Leaf rust	Net blotch
Rihane-03	MS	MS	S	S
Giza 123	MR	MR	MS	S
Giza 124	MR	MR	MS	MS
Giza 125	R	R	MS	MS
Giza 126	R	R	S	R
Line 1	MR	MR	MS	MR
Line 2	MS	MS	MS	MR
Giza 2000	R	R	MS	R



#### **Seed Distribution of Giza 2000**

Increasing foundation seeds of Giza 2000 started since 2000/2001 and will be increased in 2002/2003 as certified seeds to cover parts of the barley growing area in Egypt.

#### **Molecular fingerprint of Giza 2000**

The entire fingerprint of this cultivar using the 38 primers was shown in Figure (1) and illustrated in Table (6). Reproductive PCR products were shaded in the table. The number of bands generated during PCR ranged from 1 (D09) to 5 (E19 and Z10). The size of PCR band ranged from 114 to 4547 bp. PCR products less than 100 and over 5000 bp were excluded from the fingerprint because PCR reaction with these short primers should generate products within this range. The total number of generated bands across the 38 primer and the three replicates was 234, only 115 of them were reproducible. These bands with certain molecular weights for certain primers should be developed for each PCR reaction in order to detect fingerprints of a given genotype.

The unstable bands were suggested to result from the formation of artificial heteroduplexes between multiple amplified fragments (Wenger and Nielsen, 1991), or from non-specific amplification. He *et al.* (1992) described that these artifacts were minimized on the gradient gel, where the latter controls the consistency of PCR products by denaturing artificial heteroduplexes.

However, Yang and Quiros (1993) reported that RAPD technology provided a new alternative for cultivar identification in celery. The advantages of DNA-based pedigree assessment have been, recently, demonstrated in maize (Marsan *et al.*, 1992) and barley (Bahieldin and Ahmed 1994).

Fingerprinting of newly developed cultivars is important in which it can be used as an appraisal when commercializing these cultivars to preserve the breeder's rights. Comparison with the available germplasm to develop cultivar-specific RAPD (random amplified polymorphic DNA) markers is another important approach to trap the material mix mistakenly happened during seed storage. Besides, molecular markers for agronomically important characteristics can be detected. Cultivar identification in Egyptian barley was recently done by Bahieldin and Ahmed (1994) utilizing six cultivars. As a recommendation, all Egyptian barley germplasm ought to be characterized on the structural as well as functional level in order to detect and/or isolate valuable gene (s) for subsequent improvement of Egyptian barley for yield components, and virus, fungus and insect resistance following conventional breeding as well as genetic engineering.




Figure 1. Photographs of PCR fingerprint of the new barley cultivar Giza 2000 with 38 different random 10 mer primers in three replicates (R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub>). Numbers represent primer codes shown in Table (6). M refers to 1 kb plus DNA ladder.

Table 6. PCR fingerprint of the new barley cultivar Giza 2000 with 38 different random 10 mer primers in three replicates (R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub>).

Lane No	Primer name	Band no	R <sub>1</sub> (bp)	R <sub>2</sub> (bp)	R <sub>3</sub> (bp)	Lane No	Primer name	Band no	R <sub>1</sub> (bp)	R <sub>2</sub> (bp)	R <sub>3</sub> (bp)
1	A04	1	2429	2715	1565	11	B13				1743
		2	1769	1715	932						1336
		3	921	986	513						
		4	565					1	976	921	784
								2		721	
2	B01				2315	12	B14	1		2413	2402
		1		1270	1140			2		1461	1408
		2	983	1012	932					1193	
		3	781	623	570			3	761	803	
		4	447		346				677		
											513
3	B02	1	3206	3241		13	B15			2685	
					2781					1872	
		2	2362	2357						1424	
		3	1536	1514	1408						
		4		959	890			1	1065		1082
			683					2	899		869
			490					3	569	509	447
4	B05			3129		14	B16	1		2300	2402
		1	2173	1982	2681			2		1723	1695
					1743			3		1121	1111
		2	1563	1553	1408			4	870	836	
			1307								540
					977	15	B18	1		2300	
		3	755		667			2		1641	1791
			545					3	774	1121	1111
			349							883	633
						16	C01	1		1766	1565
5	B06			3020				2		1241	1171
		1	1864	2357	1791			3	885	1021	932
				1935							723
				1335						588	
		2	1055	1092	1000						388
					827						327
6	B07			1965		17	C05			1900	
					1650			1	1230	1521	
			1455					2		859	1234
		1		1255	954			3	774	672	1000
		2	863	1051	743	18	C08	1	1275	1195	
		3	660	782	555					946	1202
				582					761		
7	B08				3341			2		594	585
			2056						301		
		1	1379		1695	19	C10	1			1140
		2	768		1202			2	688	1052	1026
					600					534	743
8	B10		4116								
		1			3220				394		
					1743				287		
		2	983		1336	20	C19	1	1557	2342	
			878		1140			2	1101	1419	
9	B11	1	1965	1935	2074			3	940	1284	
		2	1738	1477	1650			4	787	897	
			1093							704	
					869	21	C07	1		2000	
		3	606		540			2	1350	1162	
10	B12	1		1918	1791			3	990	873	
									764	704	
		2	1036	921	784						
			621								
			508								

Table 6. Cont.

Lane No.	Primer name	Band no.	R <sub>1</sub> (bp)	R <sub>2</sub> (bp)	R <sub>3</sub> (bp)	Lane No.	Primer name	Band no.	R <sub>1</sub> (bp)	R <sub>2</sub> (bp)	R <sub>3</sub> (bp)
22	D09			1825 1407 1179 460		30	O09			2575	1301 1000 911
		1	526					1	1189	1105	
								2	833	827	
23	D10	1	1350		1301						
		2	970	1094	977	31	O10		1732		
		3	735	678	827				1366		
		4	424	344	469			1		1000	1140
			125					2	788		685
24	D20			4381 3738				3	376	545 239	322
					1945 1524	32	O12				1555
		1	1010	1337	954				1169 700	1350	
				817				1	417	469	578
		2	681		685			2	250	250	334
		3	468	537	454	33	O18		1339		
		4		344	310			1		1078	1171
				222				2	771 606	723	850
25	E07			4381 3621							454
			1902							162	
		1	1487	1407 1150	1650 1336	34	O20		2376	1492	1082
								1	1215	1025	932
		2	750		827			2		667	743
		3	655		685			3		561	650
			228					4	347	286	
26	E09			4547 3212		35	Z08				2756
		1	1487		1408				1420	2342	
				1221				1		1252	1267
		2	681		1000			2	1040		911
				545	783			3	867	827	827
		3	335		274			4	461		530 310
27	E19			4269 3645 1850		36	Z10	1	2442	2269	2191
								2	1534	1455	1408
		1	1050		1000			3		1000	1171
		2	858	850				4	981		977
		3	642	685	867				786		
		4	403	500	454			5	449	500	500
		5		298	274					239	
28	O02			4136 3532		37	Z11	1		1650	1524
								2	1159 809	805	763
				1945	2632			3	640		
			1672					4	489	530	469
		1	1403						400	359	
		2	1202		1171	38	Z13	1	951	921	805
				947	932			2		613	545
			642					3	390		372
				500							
29	O05		2953		2191			4		200 114	145
		1	1082	1841	1026						
				805							
		2	531	484	613						

## ACKNOWLEDGEMENT

This study was supported in part by the Nile Valley and Red Sea Regional Program (NVRSRP) on Resource Management, Cool-Season Food Legumes, Cereals, and Wild Oats, (Barley Program), Financed by the European Union.

## REFERENCES

- Bahieldin, A. and I.A. Ahmed (1994). Use of agarose-RAPD and DGGE-RAPD molecular markers for the identification of some barley cultivars. *Egypt. J. Genet. Cytol.*, 23: 81-94.
- Eberhart, S.A. and W.A. Russel (1966). Stability parameters for comparing varieties. *Crop Sci.*, 6: 36-40.
- Gawel, N.J. and R.L. Jarrett (1991). A modified CTAB DNA-extraction procedure for *Musa* and *Ipomoea*. *Plant Mol. Biol. Rep.*, 91: 262-266.
- Gomez, K.A. and A.A. Gomez (1984). *Statistical Procedures for Agricultural Research*. John Wiley & Sons Inc., USA.
- He, S., H. Ohm and S. Mackenzie (1992). Detection of DNA sequence polymorphisms among wheat varieties. *Theor. Appl. Genet.*, 84: 573-578.
- Innis, M.A.; D.H. Gelfand; J.J. Sninsky and T.J. White (1990). *PCR protocols*, 1st Ed. Academic Press, San Diego, p. 482.
- Krawets, S.A. (1989). The polymerase chain reaction: opportunities for agriculture. *Ag. Biotech. News Info.*, 1: 897-902.
- Marsan, P.A.; C. Livini; M.M. Messmer; E. Mechinger; P. Franceschini; G. Monfredini and M. Motto (1992). Comparison between cluster analyses from RFLP and pedigree data of inbred lines related to stiff stalk synthetic heterotic group. *Maize Genet. Coop. Newl.*, 66: 18-20.
- Saiki, R.K.; D.H. Gelfand; S. Stoffel; S.J. Scharf; R. Higuchi; G.T. Horn; K.B. Mullis and H.A. Erlich (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science*, 239: 487-491.
- Wenger, R.H. and P.J. Nielsen (1991). Reannealing of artificial heteroduplexes generated during PCR-mediated genetic isotyping. *Trend Genet.*, 6: 178.
- Williams, J.G.K.; A.R. Kubelik; K.J. Livak; J.A. Rafalski and S.V. Tingey (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucl. Acid Res.*, 18: 6531-6535.
- Wrigley, C.W.; I.L. Batey and J.H. Sherritt (1987). Complementing traditional methods of identifying cereal varieties with novel procedures. *Seed Sci. Technol.*, 15: 679-688.
- Yang, X. and C.F. Quiros (1993). Identification and classification of celery cultivars with RAPD markers. *Theor. Appl. Genet.*, 86: 205-212.

**جيزة 2000 صنف جديد من الشعير يلائم المناطق الجديدة والزراعات المطرية**  
إسماعيل عيد المنعم أحمد ، رشاد أحمد أبو العينين ، عبد ربه عبد العزيز الحاج ، ، خالد أحمد مصطفى ،  
محمود أمين المصيلحي ، أحمد محمد عرابي البواب ، هشام طه محفوظ\* ، خيرى عبد العزيز عامر ،  
محفوظ عبد الحميد ، رزق عبد الخالق رزق\*\*  
قسم بحوث الشعير - معهد بحوث المحاصيل الحقلية - مركز البحوث الزراعية -  
الجيزة - مصر  
\* معهد بحوث الهندسة الوراثية الزراعية - مركز البحوث الزراعية - الجيزة - مصر.  
\*\* معهد بحوث امراض النباتات - مركز البحوث الزراعية - الجيزة - مصر.

يهدف هذا البحث إلى التعرف بصنف الشعير الجديد (جيزة ٢٠٠٠) الذى يتميز  
بالإنتاجية العالية من المحصول ويوجد فى الاراضى الجديدة والاراضى المطرية. إنتاج هذا  
الصنف من التهجين بين الصنف المحلى جيزة 121 والسلالة المباشرة ١/١٣/٣٦٦. قيم هذا  
الصنف بالمقارنة مع الاصناف المحلية جيزة ١٢٣ ، جيزة ١٢٤ ، جيزة ١٢٥ ، جيزة  
١٢٦ وسلالتين مبشرتين من البرنامج المحلى بالاضافة إلى الصنف واسع الانتشار ربحان  
٣. أقيمت ٣٩ تجربة محصولية (١٥ تجربة محصولية فى الاراضى المطرية على مدار  
ثلاث سنوات ١٩٩٩/٢٠٠٠ وحتى ٢٠٠١/٢٠٠٢ و ٢٤ تجربة محصولية فى الاراضى  
الجديدة على مدار أربع سنوات ١٩٩٨/١٩٩٩ وحتى ٢٠٠١/٢٠٠٢). أظهرت النتائج  
تفوق الصنف الجديد جيزة 2000 معنوياً على صنف المقارنة جيزة ١٢٦ تحت ظروف  
الاراضى المطرية وأعطى محصولاً مقداره ٣,٥١ أردب/فدان بزيادة مقدارها ٠,٥٧  
أردب/فدان (١٩,٤%) عن صنف المقارنة. وفى الاراضى الجديدة تفوق الصنف الجديد  
جيزة ٢٠٠٠ على صنف المقارنة جيزة ١٢٣ بمقدار ٢,٤٦ أردب/فدان أى بنسبة  
١٧,٢٥%. ويتميز الصنف الجديد بالعديد من الصفات المرغوبة من قدرة إنتاجية عالية ،  
التكيف فى النضج ، مقاومته لأمراض الشعير الشائعة. تم تعريف البصمة الوراثية لصنف  
الشعير جيزة ٢٠٠٠ باستخدام جهاز PCR وذلك باستخدام ٣٨ بادئ عشوائى وتم الحصول  
على ١١٥ حزمة يمكن عن طريقها تعريف صنف الشعير الجديد جيزة ٢٠٠٠ وذلك حفاظاً  
على ملكية الصنف لقسم بحوث الشعير بمعهد بحوث المحاصيل الحقلية ولقد بدأ فى أكتاف  
التقاوى المعتمدة لهذا الصنف حتى تغطى إحتياجات المزارعين فى المناطق المذكورة  
سابقاً.