

## MICROBIAL CONTROL BY THE FUNGUS *Beauveria bassiana* AGAINST INSECT PESTS IN HONEY BEE STORAGE

Mansour, H.M.

Plant Protection Research Institute, Agricultural Research Center,  
Dokki, Giza, Egypt.

### ABSTRACT

The present work was carried out at three locations in Kafr El-Sheikh governorate during 2000 to study the efficiency of the fungus, *Beauveria bassiana* against the insect pests infesting honey bee combs or associated with them namely, *Galleria mellonella* L., *Achroia grisella* L., *Ephestia* sp., *Tribolium confusum* Du Val., *T. castaneum* (Herbst.), *Cassida vittata* de Villers, *Hypera brunneipennis* Boh., *Sitona lividipes*, *Gryllus burdigaiensis* and *Oxycarinus hyalinipennis* Costa under laboratory and store conditions.

The obtained results revealed that *B. bassiana* ( $1.6 \times 10^8$  conidia/ml) had the higher mortality percentage against larvae of *G. mellonella*, *A. grisella*, *Ephestia* sp. and adults of *H. brunneipennis*, *S. lividipes*, *G. burdigaiensis* and *O. hyalinipennis* at 20 days after treatments recording 96, 96, 100, 96, 86, 52 and 80%, respectively, under laboratory conditions. Also, *B. bassiana* at the same concentration caused the highest mortality for these insects recording 82, 72, 82, 92, 78, 26 and 70%, respectively, under store conditions. On the other hand, *B. bassiana* had the lowest mortality percentage against *T. confusum*, *T. castaneum* and *C. vittata* recording 10, 18 and 10%, respectively under lab conditions, while the mortality percentage under store conditions for these insects were 22, 38 and 14%, respectively. It could be stated that *B. bassiana* may be utilized as a biological agent for controlling of *G. mellonella*, *A. grisella*, *H. brunneipennis*, *S. lividipes*, *G. burdigaiensis* and *O. hyalinipennis* under store conditions.

### INTRODUCTION

Some insects take the honey bee combs as a shelter throughout the store period that extends from September to January. Among these insects are *Galleria mellonella*, *Achroia grisella* and *Ephestia* sp. (Anderson and Mignot, 1970; Burges, 1978; Atallah *et al.*, 1983; El-Hemaesy, 1983 and Mansour, 1991). These insects damage the wax combs resulting in much losses to beekeepers.

Several attempts have been made to protect the combs against these insects using control measures, from which is the biological control agents such as the bacterium *Bacillus thuringensis* and the fungus, *Beauveria bassiana* (El-Sufty, 1983; Lacey, 1985 and Anop-Ongsakul and Surakrai-Permkam, 1986. No attempts were made to investigate the possible use of *B. bassiana* in the control of these insects in stored combs.

The fungus *B. bassiana* is safe to man and animal and is not injurious to honeybee Steinhaus (1963) and Mansour (1991).

The present work was carried out to study the efficiency of *B. bassiana*, against some insects attacking stored combs, as well as the feasibility of applying the fungus to control such insects in the store.

## **MATERIALS AND METHODS**

### **1. Survey of insect pests occurring in honey bees storage:**

An experiment was carried out at three localities at Kafr El-Sheikh governorate; El-Reyad, El-Abbasia and Sakha during 2000/2001 winter seasons. Each locality was represented by 100 wax combs enclosed in ten honey bee boxes. The boxes were examined from October until January and occurring insect species were reported.

### **2. Determination of effective concentration:**

Laboratory tests were conducted to determine the infectious concentration that cause satisfactory insect mortality for the surveyed species. Larvae or adults of the considered insects were individually dipped in a suspension of the fungus conidia containing 0.1% Tween 80. The later was added for the purpose of achieving better distribution of the fungus in the suspension. Tested concentrations were  $2 \times 10^7$ ,  $4 \times 10^7$ ,  $8 \times 10^7$ ,  $1.6 \times 10^8$  and  $2.3 \times 10^8$  conidia/ml. Fifty insects (larvae or adults) of each insect species were used for each concentration. After treatment, 10 insects from each species were introduced into a Petri-dish and fed on pieces of sterilized blocks of old bee wax. A batch of 10 larvae or adults of the tested insects were treated with water containing 0.1% Tween 80 as a check.

The insects (larvae and adults) were examined 10 & 20 days after treatments. The dead individuals were taken out, counted and recorded.

### **3. Fungus efficiency in storage:**

This experiment was conducted to evaluate the virulence of the fungus, *B. bassiana* against larvae of *G. mellonella*, *A. grisella*, *Ephestia* sp. and adults of *S. lividipes*, *H. brunneipennis*, *T. confusum*, *T. costaneum*, *C. vittata*, *G. burdigalensis* and *O. hyalinipennis* occurring in wax combs under storage conditions in the apiary of Sakha Agricultural Research Station, Kafr El-Sheikh. Old wax combs were kept into an oven at 45°C for 96 hours to kill the alive insects and their stages. Then, the combs were confined in 50 breeding boxes, each has five combs. Every insect species was inoculated into 5 boxes.

To accomplish artificially infestation, fifty insects (larvae or adults) from each species were introduced into the boxes. Two hours later, each wax comb was sprayed with 5 ml of the fungus suspension containing  $1.6 \times 10^8$  conidia/ml and 0.1% Tween 80 using an atomizer. In addition, wax combs harboured in five breeding boxes were treated with 0.1% Tween 80 as a check.

All breeding boxes were kept in the store of the apiary (22-26°C and 75-80% RH). The wax combs were examined 10, 20 and 30 days after applications, and numbers of dead insects were recorded.

## RESULTS AND DISCUSSION

### 1. Survey of insect species occurring in honey bees storage:

Data obtained in Table (1) indicated that ten species of insect pests belonging to 7 families and 4 orders were surveyed from honey bee combs under storage conditions at Kafr El-Sheikh Governorate.

*Galleria mellonella*, *Achroia grisella* and *Ephestia* sp. were the most abundant and destructive insect pests. The three insect species, *Tribolium confusum*, *T. costaneum* and *Gryllus burdigaiensis* were found causing no considerable damage to combs. *Hypera brunneipennis*, *Sitona lividipes*, *Cassida vittata* and *Oxycarinus hyalinipennis* used the combs as a shelter, but no signs of damage were observed. Some authors (Ahmad *et al.*, 1983; Nunamaker *et al.*, 1986; Hussein, 1988 and Mansour, 1991) surveyed some insect species in honey bee combs, such as *G. mellonella*, *A. grisella*, *Ephestia* sp., *G. burdigaiensis* and *Merops* spp. They attack hives, brood, stored combs.

**Table (1): Surveyed insects from stored honey bee combs.**

Order/fam.	Insect species	Stage
<b>Lepidoptera:</b>		
Galleridae	The greater wax moth, <i>Galleria mellonella</i> L.	Larvae
	The lesser wax moth, <i>Achroia grisella</i> F.	Larvae
Phycitidae	The fig moth, <i>Ephestia</i> sp.	Larvae
<b>Coleoptera:</b>		
Curculionidae	The alfalfa cultures pest, <i>Sitona lividipes</i> Fab.	Adult
	The Egyptian alfalfa weevil, <i>Hypera brunneipennis</i> Boh.	Adult
Tenebrionidae	The confused flour beetle, <i>Tribolium confusum</i> Du Val.	Adult
	The red flour beetle, <i>Tribolium costaneum</i> (Herbst.)	Adult
Chrysomelidae	The tortoise beetle, <i>Cassida vittata</i> de Villers	Adult
<b>Orthoptera:</b>		
Gryllotalpidae	The field cricket, <i>Gryllus burdigaiensis</i>	Adult
<b>Hemiptera:</b>		
Lygaeidae	The cotton seed bug, <i>Oxycarinus hyalinipennis</i> Costa	Adult

### 2. Effective fungus concentration:

Data in Table (2) show mortality of insects occurring in honeybee stores treated with five concentrations of *B. bassiana* conidia. *G. mellonella* mortalities at 10 days after treatment were 58, 58, 86, 96 and 98% at the concentrations  $2 \times 10^7$ ,  $4 \times 10^7$ ,  $8 \times 10^7$ ,  $1.6 \times 10^8$  and  $2.3 \times 10^8$  conidia/ml, respectively. *A. grisella* larvae mortalities at 10 days after treatments were 12, 40, 42, 72 and 82%, while those at 20 DAT were 46, 66, 76, 96 and 100% at the concentrations  $2 \times 10^7$ ,  $4 \times 10^7$ ,  $8 \times 10^7$ ,  $1.6 \times 10^8$  and  $2.3 \times 10^8$  conidia/ml, respectively. *Ephestia* sp. larvae recorded mortality 36% at 10 DAT for the concentration  $2 \times 10^7$  conidia/ml. While recorded 74 and 100% mortalities at concentrations of  $1.6 \times 10^8$  and  $2.3 \times 10^8$  conidia/ml, respectively after 10 days from treatment. Also, the fungus at concentrations  $1.6 \times 10^8$  and  $2.3 \times 10^8$  conidia/ml caused highest *Ephestia* mortality % (100 % each) after 20 days from treatment.

***Mansour, H.M.***

2

7062

Jassim *et al.* (1988) evaluated the pathogenicity of the fungus *B. bassiana* against the fig moth, *Ephesia cautella* (Walk.) (Lepidoptera: Phycitidae). They found that the best effective concentrations were  $3 \times 10^5$  and  $4 \times 10^5$  spores/cm<sup>3</sup> as they gave 96.6% and 98.6% mortality, respectively.

Also, the results indicate that adults of *T. castaneum* and *C. vittata* were similar in their susceptibility to infection of *B. bassiana*. The highest concentration ( $2.3 \times 10^8$ ) caused 6 & 22 and 8 & 24 mortality for the first and second insects 10 and 20 days after treatment, respectively. The concentration  $2 \times 10^7$  was neither lethal to flour beetles nor to tortoise beetle. These results indicate that *B. bassiana* is not suitable as a microbial control agent for controlling these coleopterous insects. In similar studies, Abo-Aiana (1985 & 1991) and Mansour (1991) found that the fungus caused low mortality to *C. vittata* even when treated with high conidia concentrations of *B. bassiana*.

The data indicate that treating insects, *H. brunneipennis*, *S. lividipes*, *G. burdigaisis* and *O. hyalinipennis* by the fungus *B. bassiana* gave sufficient results when compared to those of *T. confosium*, *T. castaneum* and *C. vittata*. The fungus was more effective and mortalities increased as the concentration of the fungus got higher. The mortality at 10 and 20 DAT were (42 & 50)-(34 & 42)-(12 & 26) and (22 & 32%) in *H. brunneipennis*, *S. lividipes*, *G. burdigaisis* and *O. hyalinipennis*, respectively, when the fungus was applied at concentration  $2 \times 10^7$  conidia/ml. Also the highest concentration  $2.3 \times 10^8$  conidia/ml caused the highest mortality after 10 and 20 DAT (100 & 100), (80 & 88), (42 & 55) and (52 & 84%), for the same insects, respectively. Frydocva *et al.* (1989) tested two biopreparations of *Beauveria* (Boverol and Boverosil) based on the entomogenous fungus *Beauveria bassiana* for their effectiveness against stored product pests; *Sitophilus granarius*, *Dryzaepphilus surinamensis* L. and *T. confusum* Duval in the laboratory. The first 2 species were more susceptible to the preparations than the latter one.

From the economic point view, the concentration  $1.6 \times 10^8$  conidia/ml could be satisfied as a microbial control agent against the considered insects in the current study.

### **3.Fungus efficiency in storage:**

Data presented in Table (3) revealed that the lepidopterous insects occurring in stored honey bee combs were more sensitive when treated with the concentrations of  $1.6 \times 10^8$  conidia/ml. Mortalities were 26, 64 and 60 at 10 DAT for *G. mellonella*, *A. grisella* and *Ephesia* sp. respectively. Also, the fungus was highly virulent against those insects after 20 DAT, since larval mortality ranged between 72 and 82%. At 30 DAT the *G. mellonella* larvae recorded mortality of 86%. Also the *Ephesia* sp. larvae took nearly the same trend (84% mortality) while the mortality for *Achroia grisella* larvae were 92%. El-Sufty *et al.* (1986), Mansour (1991), Hung *et al.* (1992), Maniania (1992) and Ibrahim (1996) tested the susceptibility of several lepidopterous larvae, including *G. mellonella*, to the fungus *B. bassiana*. They found that the fourth and fifth larval instars of lepidopterous insects were susceptible to the fungus infection.

**Table (3): Efficiency of the fungus *B. bassiana* ( $1.6 \times 10^8$  conidia/ml.) for the lepidopterous larvae occurring in stored honey bee combs.**

Insect Species	Mortality at days after treatment					
	10		20		30	
	No	%	No	%	No	%
<i>Galleria mellonella</i>	13	26	41	82	43	86
<i>Achroia grisella</i>	32	64	36	72	46	92
<i>Ephestia</i> sp.	30	60	41	82	42	84

No dead larvae in the control.

Table (4) show the efficiency of the fungus *B. bassiana* ( $1.6 \times 10^8$  conidia/ml) for the coleopterous insects in honey bee stores.

The fungus concentration  $1.6 \times 10^8$  conidia/ml was more effective and caused higher mortality for *H. brunneipennis* and *Sitona lividipes*. The mortality at 10, 20 and 30 DAT recorded 82, 92 and 100% for *H. brunneipennis* and 64, 72 and 82 for *S. lividipes*, respectively. The fungus exhibited low mortality to *Tribolium confusum*, *T. castaneum* and *Cassida vittata* when applied at the same concentration. The mortalities were 26, 42 and 16 at 30 DAT for *T. confusum*, *T. castaneum* and *C. vittata*, respectively. Frydoka *et al.* (1989) reported that Boverol and Boverosil based on the entomogenous fungus *B. bassiana* were low effective against *T. confusum* Duval. (Coleoptera: Tenebrionidae).

**Table (4): Effect of the fungus *B. bassiana* ( $1.6 \times 10^8$  conidia/ml.) on some Coleoptrous (minor or visitor) insects in honey bee stores.**

Insect species	Mortality at days after treatment					
	10		20		30	
	No	%	No	%	No	%
<i>Hypera brunneipennies</i>	41	82	46	92	50	100
<i>Sitona lividipes</i>	32	64	36	72	41	82
<i>Tribolium confusum</i>	0.0	0.0	11	22	13	26
<i>T. costaneum</i>	2	4	19	38	21	42
<i>Cassida vittata</i>	0.0	0.0	7	14	8	16

No dead larvae in the control.

Table (5) shows the effect of the fungus *B. bassiana* concentration  $1.6 \times 10^8$  conidia/ml on visitor insects in honey bee stores.

The fungus concentration  $1.6 \times 10^8$  conidia caused low mortality (56 and 52%) for *G. burdigaiensis* when the fungus applied at 20 and 30 days after treatments, respectively. Higher mortalities, (70 and 84%) for *O. hyalinipennis* were recorded at 20 and 30 DAT, respectively. Ibrahim (1996) reported that the fungus *B. bassiana* isolated from *O. hyalinipennis* adults killed 83.3% of *G. mellonella* larvae within 16 days.

**Table (5): Effect of the fungus *B. bassiana* ( $1.6 \times 10^8$  conidia/ml.) on visitor insects in honey bee stores.**

Insect species	Mortality at days after treatment					
	10		20		30	
	No	%	No	%	No	%
<i>Gryllus burdigaiensis</i>	7	14	28	56	26	52
<i>Oxycarinus hyalinipennis</i>	17	34	35	70	42	84

No dead larvae in the control.

## REFERENCES

- Abo-Aiana, R.A.D. (1985). Studies on the cabbage butterfly *Pieris rapae* (L.). M.Sc. Thesis, Fac. Agric., Tanta Univ.
- Abo-Aiana, R.A.D. (1991). Studies on pests of sugar-beet in Kafr El-Sheikh. Ph.D. Thesis, Fac. Agric., Tanta Univ.
- Ahmed, R.; N. Muzaffar and Q. Ali (1983). Biological control of the wax moths *Achroia grisella* (F.) and *Galleria mellonella* (L.) (Lep., Galleridae) by augmentation of the parasite *Apanteles galleriae* Wilk. (Hym., Braconidae) in Pakistan. *Apiacta*. 18(1): 15-20.
- Anderson, M.A. and E.C. Mignot (1970). The number of larval instars of the greater wax moth *Galleria mellonella* L. (Lepidoptera: Pyralidae), with characters for identification of instars. *Rev. of Appl. Entomol.*, 60(4).
- Annop-Ongsakul and Surakrai-Permkam (1986). Mycoses test of white muscardine fungus [*Beauveria bassiana* (Bals.) Vuill.] on some insect pests in the laboratory. *Songklankarin J. of Science and Technology*, 8(1): 1-6.
- Atallah, M.A.; A.A. Abdel-Naby and A.A. Mohamed (1983). Duration of different developmental stages of the greater wax moth, *Galleria mellonella* L. under field conditions in middle Egypt. *Proc. 5<sup>th</sup> Arab Pesticide Conf.*, Tanta Univ., Sept., 1, p. 104.
- Burges, H.D. (1978). Control of wax moth; Physical, chemical and biological methods. *Bee World*, 59(4): 129-138.
- El-Hemaesy, A.H. (1983). On the nutrition of the lesser wax moth, *Achroia grisella* Fab. (Lepidoptera: Pyralidae). *Proc. 5<sup>th</sup> Arab pesticide Conf.* Tanta Univ., Sept. 1: 113-118.
- El-Sufty, R. (1983). Enhanced culture of *Beauveria bassiana* by incorporation of insect material into the growth medium. *J. Agric. Res.*, Tanta Univ., 9(3): 871-879.
- El-Sufty, R.; R. Saleh; S.M.I. Metwally and R. Abo-Aiana (1986). Effectiveness of *Beauveria bassiana* (Bals.) on the immature stages of *Pieris rapae* (L.) (Lepi: Pieridae). *Proc. 1<sup>st</sup>. Hort. Sci. Conf.* Tanta Univ., 1: 300-308.
- Frydocva, B.; P.H. Verner and J. Bartos (1989). The comparative effectiveness of two biopreparations based on the entomogenous fungus *Beauveria bassiana*, especially their effects on storage pests. *Sbornik Vysoke Skoly Zemedelske, Praze Fakulta Agronomicka Rada, A, Rostlinna Vyroba*, 50: 247-264.

- Hung, S.Y. and D.G. Boucias (1992). Influence of *Beauveria bassiana* on the cellular defense of the beet armyworm, *Spodoptera exigua*. J. Invertebr Pathol., 60(2): 152-158.
- Hussein, M.H. (1988). Studies on some pests of honey bee colonies in Dhofar, Oman. Proc. 4 Int. Conf. Apic. Trop. Climates, Cairo, 1988: 370-376 (1989).
- Ibrahim, R.A.A. (1996). Studies on microbial natural enemies of certain insect pests. M.Sc. Thesis, Fac. Agric., Tanta Univ.
- Jassim, H.K.; L.M. Abdullah and I. Abd El-Ahad (1988). Determination of the exact concentration of *Beauveria bassiana* to control the larvae of the fig moth. Arab J. Invertebr. Pathol., 15: 328-340.
- Lacey, L.A. (1985). *Bacillus thuringiensis* serotype H. 14 in biological control of mosquitoes. Ed. H.C. Chapman., 6: 132-158.
- Maniania, N.K. (1992). Pathogenicity of entomogenous fungi (Hyphomycetes) to larvae of the stem borers, *Chilo partellus* swinhoe and *Busseola fusca* Fuller. Insect Science and its applications (ICIPE), 13(5): 691-696.
- Mansour, H.A.M. (1991). Studies on the natural enemies of honey bees. M.Sc. thesis, Fac. Agric., Tanta Univ.
- Nunamaker, R.A.; W.T. Wilson; M. Cal; L. Choote and G. Tween (1986). Beekeeping in Belize, central America, with notes on diseases and parasites. Bee-World, 67(4): 151-156.
- Steinhaus, E.A. (1963). Insect pathology. Vol. I. Academic Press. New York, 412 pp.

## تأثير الفطر *Beauveria bassiana* على الحشرات التي تتواجد في مخازن نحل العسل

حمدي احمد متولى منصور

معهد بحوث وقاية النبات مركز البحوث الزراعية - دقى - جيزه

أجريت هذه الدراسة فى ثلاث اماكن مختلفة داخل محافظة كفر الشيخ (الرياض - العباسية - سخا) وذلك خلال الفترة التى يتم فيها تخزين البراويز الزائدة عن حاجة طوائف النحل إليها من أكتوبر حتى يناير وذلك لإيجاد طريقة جديدة لمكافحة الآفات دون تلوث للبيئة.

حيث تم استخدام عدة تركيبات مختلفة من الفطر *B. bassiana* كالتالى  $2 \times 710$  ،  $4 \times 710$  ،  $8 \times 710$  ،  $1.6 \times 810$  ،  $3.2 \times 810$  كوتيديا/ملتر ماء رشاً على الحشرات التى تتواجد فى براويز الشمع حيث تم معاملتها فى اطاق بنرى تحت ظروف المعمل ، وتم دراسة أفضل التركيزات التى تعطى أعلى نسبة موت للحشرات وكان التركيز  $1.6 \times 810$  كوتيديا/ملتر هو أفضل تركيز حيث أعطى نسبة موت أعلى للحشرات بعد 20 يوماً من المعاملة ، وبعد ذلك تم استخدامه رشاً على البراويز بمعدل 5 مللى من معلقة القطر لكل برواز تحت ظروف المعمل على الحشرات الأتية دودة الشمع الكبيرة - دودة الشمع الصغيرة - ديدان جريش الذره - خنفساء الدقيق الصدائية - خنفساء الدقيق الحمراء المتشابهة خنفساء البنجر - سوسة أوراق البرسيم - سوسة جذور البرسيم - صرصور الغيط الأليف - بق بذره القطن وكانت النتائج كالتالى 96 ، 96 ، 100 ، 10 ، 18 ، 10 ، 96 ، 86 ، 52 ، 80% على التوالى. بينما أعطى نفس التركيز بعد 20 يوم من المعاملة تحت ظروف التخزين فى مخزن النحل نسب موت 82 ، 72 ، 82 ، 22 ، 38 ، 14 ، 92 ، 72 ، 56 ، 70% على التوالى. ومن نتائج الدراسة يتضح أن الفطر كان أكثر كفاءة على حشرات دودة الشمع الكبيرة دودة الشمع الصغيرة دودة جريش الذره وسوسة جذور البرسيم وخنفساء أوراق البرسيم وصرصور الغيط الأسود وبق بذره القطن وكان ذو تأثير ضعيف على خنفساء الدقيق الصدائية وخنفساء الدقيق المتشابهة وخنفساء بنجر السكر.

**Table (2): Efficiency of the fungus, *Beauveria bassiana* against insects occurring in honey bee stores under lab. conditions.**

Concentration com./ml.	Insect mortality at days after treatments (DAT)																			
	<i>G. mellonella</i> larvae		<i>A. grisella</i> larvae		<i>Ephestia</i> sp. larvae		<i>T. confosum</i> adult		<i>T. castaneum</i> adult		<i>C. vittata</i> adult		<i>H. brunneipennis</i> adult		<i>S. lividipes</i> adult		<i>G. burdigalensis</i> adult		<i>Oxycarmus hyalinipennis</i> adult	
	10	20	10	20	10	20	10	20	10	20	10	20	10	20	10	20	10	20	10	20
2 x 10 <sup>7</sup>	34	58	12	46	36	58	0.0	0.0	0.0	0.0	0.0	0.0	42	50	34	42	12	26	22	32
4 x 10 <sup>7</sup>	38	58	40	66	58	66	0.0	0.0	0.0	0.0	0.0	4	46	62	46	52	16	28	42	52
8 x 10 <sup>7</sup>	42	86	42	76	62	86	0.0	6	0.0	12	0.0	4	72	82	64	74	26	44	50	74
1.6 x 10 <sup>8</sup>	52	96	70	96	74	100	0.0	10	6	18	8	10	84	96	74	86	40	52	52	80
2.3 x 10 <sup>8</sup>	76	98	78	100	100	100	2	12	6	22	8	24	100	100	80	88	42	55	52	84

**No dead insects in the control**