

THE RELATION BETWEEN THE RENIN-  
ANGIOTENSIN SYSTEM AND FIBRINOLYTIC  
SYSTEM

M. M. Al-Zendah and A. Alfaqeh  
*Physiology & Biochemistry Departements - Faculty  
of Medicine  
Al-Fateh University for Medical Sciences*

ABSTRACT

*Background: Several lines of evidence point to an interrelation of the renin angiotensin system (RAS) with the endogenous fibrinolytic system.*

*Aim & Methods: this study was planned to examine the effect of salt depletion as a method of activation of the endogenous RAS on plasma fibrinolytic balance in 10 healthy human subjects in the presence and absence of Angiotensin converting enzyme inhibitor- ACEi (captopril). Activation of the RAS during low salt intake was documented by a significant increase in serum aldosterone concentration.*

*Results: the data suggest that activation of the RAS results in increased plasminogen activator inhibitors (PAI-I) antigen and that interruption of the RAS with the ACE inhibitor captopril significantly lowers PAI-I antigen without lowering tissue-type plasminogen activator (t-PA) antigen.*

*Conclusion: this data provides an evidence of a direct functional link between the RAS and the fibrinolytic system in humans and these findings may help to elucidate possible mechanisms by which ACE inhibition exerts vasculoprotective effects and reduces the risk of atherothrombotic events.*

## **INTRODUCTION**

Impaired fibrinolytic function, characterized by increased plasminogen activator inhibitor type 1 (PAI-1) levels and decreased tissue plasminogen activator (t-PA) activity, has been found in patients with hypertension and may account in part for the increased risk of atherosclerosis and its clinical complications in these patients.(1&2)

The balance between plasminogen activators and plasminogen activator inhibitors is a major determinant of net fibrinolytic activity(3&4).

Although the regulation of PAI-1 in vitro has been studied, (5,6,7) factors that control the production and secretion of PAI-1 in vivo are less well characterized.

The renin-angiotensin and fibrinolytic systems both play critical roles in cardiovascular homeostasis. Angiotensin II is involved mainly in blood pressure control and the handling of salt and water, thereby preventing the deleterious effects of hemorrhage. Angiotensin converting enzyme (ACE) inhibition has clearly been shown to improve prognosis in patients with congestive heart failure (8) and following myocardial infarction (9, 10).

This interrelationship may involve mechanisms other than changes in arterial blood pressure. In addition to various possible interactions, accumulating evidence suggests that the renin-angiotensin system is involved in the regulation of the fibrinolytic system (11).

The mechanisms through which activation of the RAS increases or ACE inhibition reduces the risk of ischemic cardiovascular events in selected populations are not known.

Angiotensin II has been suggested to mediate this interrelationship because this peptide was shown to stimulate plasminogen activator inhibitor-1 (PAI-1) in experimental settings (12). However, evidence

## **The renin-angiotensin system and fibrinolytic system**

---

from studies in man regarding effects of Angiotensin II on fibrinolytic function remains controversial.

To test this hypothesis, this study was planned to examine the effect of salt depletion on tissue-type plasminogen activator antigen and plasminogen activator inhibitor-1 antigen in normotensive subjects in the presence and absence of ACEi (captopril).

### **MATERIAL & METHODS**

Ten healthy male normotensive volunteers aged 30-45 years and with mean body mass index  $30 \pm 2.1$  Kg/m<sup>2</sup> were used in this study.

All subjects underwent a complete history and physical examination before investigation. Subjects with cardiovascular, renal, endocrine, or pulmonary disease were excluded. Written informed consent was obtained.

Each subject was provided with high (200mmol/d) salt, caffeine-free, and alcohol-free diet for 5 days. At 10 AM of the 5th day of diet, a catheter was placed in the antecubital vein and blood was drawn through the catheter. Then each subject was provided with a low (10 mmol/d) salt for another 5 days then another blood sampling is obtained. Then subjects were maintained on the 25mg BID dose of captopril for an additional 14 days (13). They were then provided a 10-mmol sodium diet for the last 5 days of captopril treatment. On the 5th day of the diet blood sampling were repeated at the end of the study.

All blood samples were obtained at 10 AM to avoid the diurnal variation of the measurements and placed on ice and immediately centrifuged. Blood for measurement of PAI-1 and tPA was collected in standard evacuated tubes containing 0.105 mol/L sodium citrate. The separated plasma or serum were frozen and stored at  $-70^{\circ}\text{C}$  until the time of assay.

Plasma samples were assayed for t-PA antigen and PAI-1 antigen using two site enzyme linked immunosorbent assay (kits purchased from

Biopool AB, Umea Sweden). Serum Aldosterone was measured by radio-immunoassay technique.

The PAI-1 and tPA mass ratio was determined by dividing plasma concentrations (ng/ml) by the molecular weights of the 2 proteins, with a value of 70.000 g/mol used for tPA and a value of 50.000 g/mol for PAI-1 (14)

**Statistical analysis:** All data were expressed as mean  $\pm$  SD (standard deviation) and comparison of the data in different groups was performed by applying the paired student-T test.  $P < 0.05$  was the criterion for statistical significance.

## **RESULTS**

Low salt intake was associated with significant increase in PAI-1 antigen level in relation to the level in high salt intake ( $p < 0.001$ ). Also captopril significantly decreased PAI-1 antigen level measured under low salt conditions ( $p < 0.001$ ). (Table 1 & Fig.1) tPA increase with low salt intake than in high salt intake but there is no effect of ACE inhibition on tPA antigen during salt depletion (Table 1 & Fig.2). So, PAI-1/tPA molar ratio was significantly lower with ACE inhibition than during low salt alone. Also low salt intake was associated with high PAI-1/tPA molar ratio than with high salt intake ( $p < 0.001$ ) (Table 1 & Fig.3). Low salt intake was associated with increased aldosterone compared with high salt intake ( $p < 0.001$ ), ACE inhibition decreased the aldosterone level ( $p < 0.001$ ) under low salt conditions, but the aldosterone level remained significantly higher than under high salt intake ( $p < 0.05$ ). (Table 1 & Fig.4) There was a highly significant positive correlation between PAI-1 antigen levels and serum aldosterone under low salt conditions ( $r = 0.8812$ ) ( $P < 0.001$ ). In contrast, there was no significant correlation between PAI-1 and aldosterone under high salt conditions ( $r = 0.3961$ ) ( $P > 0.05$ ).

---



---

### The renin-angiotensin system and fibrinolytic system

---




---

**Table (1): Effect of salt intake and ACE inhibition on fibrinolytic parameters and serum aldosterone level.**

<i>parameter</i>	<i>High salt</i>	<i>low salt</i>	<i>low salt + captopril</i>
PAI-1 antigen (ng/ml)	15.26 ± 1.32	23.69±2.08***	15.71 ± 1.16 ###
tPA antigen (ng/ml)	4.67 ± 0.54	5.79 ±0.66***	5.55 ± 0.55***
PAI-1/tPA	4.8 ± 0.78	6.37 ±0.86***	4.0 ± 0.39** #
Serumaldosterone conc.(ng/dl)	10.33 ± 1.05	23.81±2.66***	12.72 ± 2.73* ###

\* P< 0.05, \*\* P<0.01, \*\*\* P<0.001 vs. high salt ## P<0.01, ### P<0.001 vs. low salt




**Figure (1): Effect of salt intake and ACE inhibition on PAI-1 antigen(ng/ml). inhibition on tPA antigen (ng/ml).**

---

## The renin-angiotensin system and fibrinolytic system

---



**Figure (2): Effect of salt intake and ACE inhibition on tPA antigen (ng/ml).**





Figure (3): Effect of salt intake and ACE inhibition on PAI-1 /tPA mass

---

## The renin-angiotensin system and fibrinolytic system

---



**Figure (4): Effect of salt intake and ACE inhibition on aldosterone concentration (ng/dl).**

## **DISCUSSION**

This study examined the effect of activation of the endogenous RAS on plasma fibrinolytic balance in healthy human subjects. The data suggest that activation of the RAS by low salt intake results in increased PAI-I antigen and that interruption of the RAS with the ACE inhibitor captopril significantly lowers PAI-I antigen without lowering tPA antigen. In this study, activation of the RAS was documented by significant increases in serum aldosterone concentration.

The correlation between PAI-I antigen and serum aldosterone concentrations observed in this study further supports an interaction between the RAS and fibrinolytic system.

The sensitivity of the adrenal cortex to AngII varies with sodium intake such that conditions of salt depletion, Ang II is the major determinant of aldosterone (15). The highly statistically significant correlation between serum aldosterone and PAI-I antigen under low salt conditions and the lack of such a correlation under high salt conditions supports the hypothesis that Ang II regulates vascular PAI-I levels.

*Sechi et al.*, (16) stated that a strong and independent association exists between renin, aldosterone, plasma fibrinogen, and plasminogen activator inhibitor-1 (PAI-1) levels and this relationship might contribute to the development of hypertensive organ damage.

ACE inhibitors have been shown to reduce progression of atherosclerosis in several animal models (17, 18) and to reduce the vascular expression of PAI-I in normal and balloon-injured vessels (19).

The present study suggests a mechanism whereby ACE inhibitors could alter the incidence of ischemic cardiovascular events in the setting of an activated RAS. A lack of effect of ACE inhibition on tPA antigen in the present study may simply reflect the effects of ACE inhibition on the kallikrein-kinin system as well as the RAS. ACE inhibitors not only decrease the production of Ang II but also decrease

the degradation of bradykinin. Bradykinin has been shown to be a potent stimulus for tPA secretion *in-vitro* and *in-vivo* (20).

So, this study is in agreement with previous study by **Brown *et al.***, (20) that provides evidence for a direct functional link between the RAS and fibrinolytic system in humans. It suggests a mechanism through which ACE inhibitors could favorably alter progression of vascular disease, particularly in the setting of clinical states associated with activation of the tissue RAS.

Moreover, **Ridker *et al.***, (21) found that intravenous angiotensin II dose-dependently increased plasma PAI-1 antigen levels in healthy volunteers, whereas t-PA concentrations were unaffected or tended to be reduced. Also, in a clinical trial by **Pfeffer *et al.***, (8) has demonstrated that the administration of ACE inhibitors to patients with left ventricular dysfunction reduces the incidence of recurrent myocardial infarction by approximately 25%.

The positive effect of ACE-I on the fibrinolytic system has been related to: 1) inhibition of angiotensin II, which stimulates PAI-1 expression; 2) inhibition of degradation of bradykinin, a potent stimulus for tPA production; and 3) improvement of insulin sensitivity (4).

However, Larsson *et al.*, (22) found that in healthy volunteers a short-term infusion of angiotensin II increased t-PA activity and antigen levels in plasma, suggesting that angiotensin II enhances fibrinolysis under these experimental conditions. There may be several explanations for these apparently conflicting results of Larsson. One might be the duration of angiotensin II infusion, or due to the hemodynamic effects caused by angiotensin II. Another possible consideration is that the clearance of t-PA may have been reduced during angiotensin II infusion.

Vaughan (23) reported that individuals with reduced fibrinolytic activity are at increased risk for ischemic cardiovascular events, and reduced fibrinolysis may underlie some of the pathological consequences of reduced nitric oxide availability. Within the vasculature, angiotensin II stimulates the release of PAI-1, thereby reducing fibrinolytic activity. Thus the plasminogen activator system is largely controlled by the renin- angiotensin system (RAS) .In

## **The rennin-angiotensin system and fibrinolytic system...**

---

accordance with this finding; treatment with angiotensin converting enzyme inhibitors is associated with substantial reductions in the incidence of ischemic cardiovascular events.

Taken together, these findings raise the possibility that Ang II may contribute to the development of a prothrombotic state at least in part by increasing plasma levels of PAI-1, thereby reducing the net activity of the endogenous fibrinolytic system. This potential relation between the renin-angiotensin system and fibrinolytic function may have important clinical and therapeutic consequences.

### **REFERENCES**

- 1- Fogari R, Zoppi A. (2006):** Antihypertensive drugs and fibrinolytic function. *Am J Hypertens.*; 19 (12):1293-9.
- 2- Dielis AW, Smid M, Spronk HM, Houben AJ, Hamulyák K, Kroon AA, Ten Cate H, de Leeuw PW.(2007):** Changes in fibrinolytic activity after angiotensin II receptor blockade in therapy-resistant hypertensive patients. *J Thromb Haemost.* 5(7):1509-15.
- 3- Vaughan D.E (1998):** Fibrinolytic balance,the renin-angiotensin system and atherosclerotic disease. *Eur Heart J.vol.19suppl.G.*
- 4- Asselbergs FW, Williams SM, Hebert PR, Coffey CS, Hillege HL, Navis G, Vaughan DE, van Gilst WH, Moore JH.(2006):** The gender-specific role of polymorphisms from the fibrinolytic, renin-angiotensin, and bradykinin systems in determining plasma t-PA and PAI-1 levels.*Thromb Haemost.* 96 (4):471-7.
- 5- Brown NJ, Vaughan DE, Fogo AB.(2002):** The renin-angiotensin-aldosterone system and fibrinolysis in progressive renal disease. *Semin Nephrol.*; 22(5):399-406. 
- 6- Sawdey MS, Luskutoff DJ, (1991):** Regulation of murine type-1 plasminogen activator inhibitor gene expression in vivo: Tissue species and induction by lipopolysaccharide, tumour necrosis factor –a transforming growth factor-B.*J Clin Invest*; 88:1346-135.
- 7- Schneider DJ, Sobel BE (1991):** Augmentation of synthesis of plasminogen activator inhibitor type 1 by insulin and insulin-like

- growth factor type-1: implications for vascular disease in hyperinsulinism states. *Med. Sci*; 88:9959-9963.
- 8- **Pfeffer MA, Braunwald E, Moye LA, Basta L, Brown EJ, Cuddy TE, (1992):** Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* ; 327:669-677.
  - 9- **The Acute infarction Ramipril Efficacy study investigators (1993):** Effect of ramipril on mortality and morbidity of survivors of acute myocardial infarction with clinical evidence of heart failure. *Lancet*; 342:821-828.
  - 10- **Vaughan DE, Lazos SA, Tong K (1995):** Angiotensin II regulates the expression of plasminogen activator inhibitor-1 in cultures endothelial cells. *J Clin Invest*; 95:995-1001.
  - 11- **Lottermoser K, Hertfelder HJ, Vetter H, Dusing R (2000):** Renin-angiotensin-aldosterone system and fibrinolysis. *Med Klin* 15; 95 (12):683-8.
  - 12- **Lottermoser K, Hertfelder HJ, Gohlke P, Vetter H, Düsing R.(2004):** Short-term effects of exogenous angiotensin II on plasma fibrinolytic balance in normal subjects. *Clin Exp Hypertens.*; 26(1):13-26.
  - 13- **Lottermoser K, Wostmann B, Weisser B, Hertfelder HJ, Schmitz, Vetter H, Dusing R (1999):** Effects of captopril on fibrinolytic function in healthy humans. *Eur J Med Res* 26; 4 (1):31-4.
  - 14- **Brown NJ, Agirbasli MA, Williams GH, Reid W, Vaughan DE.(1998):** Effect of activation and inhibition of the renin-angiotensin system on plasma PAI-1. *Hypertension*; 32:965-971.
  - 15- **Oelkers W, Brown JJ, Fraser R, Lever AF, Morton JJ, Robertson JI(1974):** Sensitization of the adrenal cortex to angiotensin II in sodium –deplete man. *Cir.Res*; 40:69-77.
  - 16- **Sechi LA, Novello M, Colussi G, Di Fabio A, Chiuch A, Nadalini E, Casanova-Borca A, Uzzau A, Catena C.(2008):** Relationship of plasma renin with a prothrombotic state in hypertension: relevance for organ damage. *Am J Hypertens.* Dec; 21(12):1347-53.

---

### The rennin-angiotensin system and fibrinolytic system...

---

- 17- **Hayek T, Keider S, Mei-Yi, Oiknine J, Breslow J. (1995):** Effect of angiotensin converting enzyme inhibitors on LDL lipid peroxidation and atherosclerosis progression in apo E deficient mice. *Circulation*, 92.
- 18- **Aberg G, Ferrer P. (1990):** Effects of captopril on atherosclerosis in cynomolgus monkeys. *J Cardiovasc. Pharmacol*; 15:S65-S72.
- 19- **Hamdan AD, Quist WC, Gagne JB, Feener EP. (1996):** Angiotensin-converting enzyme inhibition suppresses plasminogen activator inhibitor-1 expression in the neointima of balloon-injured rat aorta. *Circulation*; 93:1073-1078.
- 20- **Brown NJ, Nadeau J, Vaughan DE (1997):** Selective stimulation of tissue type plasminogen activator (t-PA) in vivo by infusion of bradykinin. *Thromb Haemost.*; 77:522-525.
- 21- **Ridker PM, Gaboury CL, Colin PR et al., (1993):** Stimulation of plasminogen activator inhibitor in vivo by infusion of angiotensin II, *Circulation*; 87:1969-1973.
- 22- **Larsson P.T, Schwieder JH, Wallen NH and Hjerdahl (1999):** Acute effects of angiotensin II on fibrinolysis in healthy volunteers. *Blood Coagulation and Fibrinolysis*, 10:19-24.
- 23- **Vaughan DE (2001):** Angiotensin, fibrinolysis and vascular homeostasis. *Am. J. Cardiol.* 19; 87(8A):18C-24C.