

PRELIMINARY TRIALS FOR CONTROL OF CHICKEN SALMONELLOSIS USING LYOPHILIZED IMMUNIZED EGG YOLK WITH EMPHASIISON

ROFAIL, S. K., GERMINE S. SAAD and S. B. EL-GALLAD

Veterinary Serum and Vaccine Research Institute , ARC, Dokki Giza

(Manuscript received 24 June 2014)

Abstract

This study has been designed to prepare the lyophilized immunized egg yolk using different stabilizers as skimmed milk, sucrose lacto albumin and sucrose peptone. These preparations were tested in chicken against Salmonellosis. Chicken that received immunized egg yolk and experimentally infected with *Salmonella typhimurium* showed survival rate of (90 %) compared with the group that taken nonimmunized egg yolk (20 %). No difference in the immunogenic power of using different stabilizers with these batches. The immunogenic power of these preparations when stored at -20 °C , 4 °C and at room temperature for 3, 6 and 9 months after injection to mice revealed that these preparations were potent when stored at -20°C, then at 4°C . So lyophilized immunized egg yolks effective in protecting chicken against Salmonellosis.

INTRODUCTION

Chicken Salmonellosis is an acute highly fatal disease. The disease is enzootic and responsible for great economic losses to poultry industry where the immune deficiency status play an important role in the progress of the bacteria (Quinn *et. al.*, 2002 and Pietro and Duncan, 2006.

Treatment of chicken Salmonellosis is often ineffective due to the presence of antibiotic drug resistant strain of *Salmonella* and failure to identify the drug susceptibility pattern of involved strain. Many other drugs and chemicals that can enhance various aspects of the immune response, had been tested to increase the rate of immune response against such infection (Rofail, 2007 and Rofail *et. al.*, 2011) .

Recently great attention has been directed to the use of natural therapeutic agents against many infection of which yolk of vaccinated chicken have been used for prophylaxis and treatment, as used for control of chicken Salmonellosis (Rofail and Saad 2013) . Little effort have been Made to investigate the role Of egg yolk in the control Salmonellosis. So this study aimed to prepare a lyophilized immunized egg yolk using different stabilizers, and to test the efficacy of these preparations when stored at different temperatures for some months in chicken and mice .

MATERIALS AND METHODS

Experimental Hens

A total number of 20 laying hens of about 5 Months old were used for egg yolk production and another 60 chicken, of 2 weeks old age used for experimental designs.

Experimental Mice

Two hundred and forty healthy mice each of 25 gm were kept in isolated cages under strict Hygenic condition during the experiment.

Vaccine

Imported inactivated bivalent vaccine of *Salmonella typhimurium* and *Salmonella enteritidis* were used in vaccination of chicken.

Salmonella typhimurium isolate

Local isolate of *Salmonella typhimurium* was used and identified by gram staining, colonial morphology biochemical and serological reactions. This strain was used for challenge test in chicken and mice as described by Forbes *et al.*, (1998).

Vaccination of hens :

Hens were divided into 2 groups (10 each), the first group was vaccinated by inactivated bivalent vaccine of salmonella with 1 ml I/M weekly for 3 weeks as described by Ikemori *et al.* (1997), the other group was kept as negative control.

Egg Yolk samples :

Egg Yolk were collected weekly from non vaccinated group and 2 weeks after last vaccination of chicken for two months. It was collected in clean dry screw capped bottles and preserved frozen till used.

Stabilizers :

The following stabilizers were used :

1- **Skimmed milk** : It was produced by Sketon company, Iierk, Ireland (Wyeth SMA) it was used at dilution (10 %) in distilled water and sterilized by autoclaving then added to the immunized egg yolk and non-immunized as 1 :1 ratio.

2- **Sucrose lactoalbumin** : It was prepared according to Stanly and Walter(1999) then added to the immunized egg yolk and non immunized as 1 :1 ratio.

3- **Sucrose peptone** : It was prepared according to Stanely and Walter(1999) by mixing 0.5 % sucrose and 1 % peptone then added to the immunized egg yolk and non immunized as 1:1 ratio..

Lyophilization of immunized egg yolk :

The egg yolk fluid was mixed with different stabilizers and they were lyophilized by freeze drying method similetenuously in accord ance with Angus *et al.* (1977).

Experimental designs :

Table (1) illustrate the experimental design

Table 1. experimental design for chicken :

Group	No.	Administred preparation	Route	Challenge
1	10	Lyophilized immunized egg yolk with skimmed milk	Each chicken receive 1 ml of each preparation per os daily for ten days	Each chicken in each group received 1 ml I/P of twenty four hours broth culture of salmonella typhimurium containing 1.5×10^8 CFU/ml such chicken observed after experimental infection and mortalities were recorded, smears from internal organs of freshly dead chicken were subjected to bacteriological examination for reisolation of the organism according to Forbes et al (1998)
2	10	Lyophilized immunized egg yolk with sucrose lactoalbumin		
3	10	Lyophilized immunized egg yolk with sucrose peptone		
4 a	10	Lyophilized nonimmunized egg yolk with skimmed milk		
b	10	Lyophilized nonimmunized egg yolk with sucrose lactoalbumin		
c	10	Lyophilized nonimmunized egg yolk with sucrose peptone		

Keeping quality of lyophilized immunized egg yolk :

Lyophilized immunized egg yolk with sucrose lactoalbumin for example was exposed to different temperature (-20°C , 4°C and room temperature) and stored for period of 3,6,9 months. Table (2) Illustrate the experimental design for keeping quality in mice.

Table 2. Experimental design in mice :

Group	NO.	Administred preparation	Route	Challenge
1		Lyophilized immunized egg yolk with sucrose lactoalbumin as example	Each mice receive 0.5 ml of each preparation per Os daily for ten days	Each mice in each group received I/P with 0.5 ml of twenty four hours broth culture of salmonella typhimurium containing 1.5×10^8 CFU / mland mice observed after experimental infection and mortalities were recorded, smears from internal organs of freshly dead chicken were subjected to bacteriological examination for reisolation of the organism according to Forbes et al (1998)
a	10	Stored at -20°C for 0 time		
b	10	Stored at -20° C for 3 months		
c	10	Stored at -20° C for 6 months		
d	10	Stored at -20° C for 9 months		
2		Lyophilized immunized egg yolk with sucrose lactoalbumin as example		
a	10	Stored at 4°C for 0 time		
b	10	Stored at 4° C for 3 months		
c	10	Stored at 4° C for 6 months		
d	10	Stored at 4° C for 9 months		
3		Lyophilized immunized egg yolk with sucrose lactoalbumin as example		
a	10	Stored at room temperature for 0 time		
b	10	Stored at room temperature for 3 months		
c	10	Stored at - room temperature for 6 months		
d	10	Stored at room temperature for 9 months		
4		Lyophilized non immunized egg yolk with sucrose lactoalbumin as example		
a	10	Stored at -20°C for 0 time		
b	10	Stored at -20° C for 3 months		
c	10	Stored at -20° C for 6 months		
d	10	Stored at -20° C for 9 months		
5		Lyophilized non immunized egg yolk with sucrose lactoalbumin as example		
a	10	Stored at 4°C for 0 time		
b	10	Stored at 4° C for 3 months		
c	10	Stored at 4° C for 6 months		
d	10	Stored at 4° C for 9 months		
6		Lyophilized nonimmunized egg yolk with sucrose lactoalbumin as example		
a	10	Stored at room temperature for 0 time		
b	10	Stored at room temperature for 3 months		
c	10	Stored at - room temperature for 6 months		
d	10	Stored at room temperature for 9 months		

RESULTS AND DISCUSSION

This work has been designed to determine the protective potentiality of the prepared lyophilized immunized egg yolk against chicken salmonellosis using different stabilizers (skimmed milk, sucrose lactoalbumin and sucrose peptone) and to assess their keeping quality at different temperatures (-20° C, 4°C and room temperature) ,

for 3, 6 and 9 Months. Data presented in table (3) showed that chicken received lyophilized immunized egg yolk (with different stabilizers) resist infection with *Salmonella typhimurium* with survival rate reached 90 % compared with 20 % of the nonimmunized group . There was no difference in the immunogenic power between different stabilizers used, the antimicrobial immune response produced by treated chicken with lyophilized immunized egg yolk was similar to those obtained by Kuroski *et al.*(1993) who found that egg yolk immunoglobulin protect mice against experimental infection by Rota virus. Also Hiraga *et al.* (1990) reported that egg yolk derived from immunized hen was virtually a significant efficient source of antibodies and can prevent Rota virus infection. Egg yolk antibodies resulted from vaccination of laying hens are thought to provide a suitable source of immunity to chicken against experimental infection by *Salmonella typhimurium* Rofail and Germine saad (2013) proved that oral administration of the egg yolk to Mice could be an important tool of treatment against Salmonellosis in mice.

The strong effectiveness of egg yolk antibodies to infectious pathogens also reported by (Yokoyama *et al.* ,1992) who added that addition of egg yolk to the taad formula or even its application as a separate therapeutic agent is helpful for Neonatal piglets.

Reisolation of bacteria from treated groups revealed absence of *Salmonella typhimurium* in internal organs, blood samples and even no shedding in the feces. Meanwhile, the group received non immunized egg yolk as well as the control groups *Salmonella typhimurium* was detected clearly in internal organs , blood samples and shedding of the organism in the feces was also recorded. These findings were similar to those described by (Rofail and Germine saad 2013).

The considerable variations affecting the immunized egg yolk potency occur during lyophilization process depends on the effect of cooling rate, residual water content, storage temperature, length of storage and thawing rate before lyophilization and the containing additive such as stabilizers include that protein or other organic compounds that may extend the shelf life or enhance the period of the immunized egg yolk (Stanely and Walter 1999). Results of studying the effect of different temperature (-20 ° C, 4° C and at room temperature) on immunogenic power of lyophilized immunized egg yolk that determined in laboratory mice and challenged with *Salmonella typhimurium* as shown in table (4) It was found that preparations kept at -20° C for 3 months were the best and recorded higher immunogenic power percent (90%). These findings are parallel to the (Nisar 1988) who found that the freezing temperature is the best for vaccine and the room temperature fail to keep the vaccine potent, also these results were almost like to those recorded by (Tina *et al.*, 2000),

who recommended the storage of egg yolk at -20°C for better stability. Also (Lily *et. al.*, 2004) cleared that the prepared attenuated Rift Valley fever vaccine was being suitable for use when stored at -20°C without loss in its titre.

From the above mentioned results it could be concluded that the lyophilized immunized egg yolk was safe and immunogenic as it gave up to 90 % protection against *Salmonella Typhimurium* infection in chicken, as well as the best preservation in freezing temperature (-20°C) for 3 months storage can be used safely and effectively.

Table 3. Efficiency of lyophilized immunized egg yolk in chicken experimentally infected with *salmonella typhimurium*.

Group	No	Admisted preparation	No. of the dead chicken /total No of chicken	Mortality rate	No of survived chicken/total No of chicken	Protection %
1	10	Lyophilized immunized egg yolk with skimmed milk	1/10	10 %	9/10	90 %
2	10	Lyophilized immunized egg yolk with sucrose, lactoalbumin	1/10	10 %	9/10	90 %
3	10	Lyophilized immunized egg yolk with sucrose peptone	1/10	10 %	9/10	90%
4 a	10	Lyophilized non-immunized egg yolk with skimmed milk	8/10	80 %	2/10	20%
b	10	Lyophilized non-immunized egg yolk with sucrose lacto albumin	8/10	80 %	2/10	20%
c	10	Lyophilization non-immunized egg yolk with sucrose peptone	8/10	80 %	2/10	20%

Table 4 . keeping quality of lyophilized immunized egg yolk determined by mice experimentally infected with salmonella typhimurium.

Group	NO.	administred preparation	No. of dead mice /total No of mice	Mortality rate	No of survived mice /total no of mice	Protectio n %
1		Lyophilized immunized egg yolk with sucrose lactoalbumin as example				
	a	10 Stored at -20°C for 0 time	1/10	10%	9/10	90 %
	b	10 Stored at -20° C for 3 months	1/10	10%	9/10	90 %
	c	10 Stored at -20° C for 6 months	3/10	30%	7/10	70 %
d	10 Stored at -20° C for 9 months	4/10	40%	6/10	60 %	
2		Lyophilized immunized egg yolk with sucrose lactoalbumin as example				
	a	10 Stored at 4°C for 0 time	3/10	30%	7/10	70 %
	b	10 Stored at 4° C for 3 months	3/10	30%	7/10	70 %
	c	10 Stored at 4° C for 6 months	4/10	40%	6/10	60 %
d	10 Stored at 4° C for 9 months	4/10	40%	6/10	60 %	
3		Lyophilized immunized egg yolk with sucrose lactoalbumin as example				
	a	10 Stored at room temperature for 0 time	4/10	40 %	6/10	60 %
	b	10 Stored at room temperature for 3 months	4/10	40%	6/10	60 %
	c	10 Stored at - room temperature for 6 months	5/10	50%	5/10	50 %
d	10 Stored at room temperature for 9 months	5/10	50%	5/10	50 %	
4		Lyophilized non immunized egg yolk with sucrose lactoalbumin as example				
	a	10 Stored at -20°C for 0 time	8/10	80 %	2/10	20 %
	b	10 Stored at -20° C for 3 months	8/10	80 %	2 /10	20 %
	c	10 Stored at -20° C for 6 months	9/10	90 %	1/10	10 %
d	10 Stored at -20° C for 9 months	9/10	90%	1/10	10 %	
5		Lyophilized non immunized egg yolk with sucrose lactoalbumin as example				
	a	10 Stored at 4°C for 0 time	8/10	80 %	2/10	20 %
	b	10 Stored at 4° C for 3 months	8/10	80 %	2/10	20 %
	c	10 Stored at 4° C for 6 months	9/10	90 %	1/10	10 %
d	10 Stored at 4° C for 9 months	9/10	90 %	1/10	10%	
6		Lyophilized nonimmunized egg yolk with sucrose lactoalbumin as example				
	a	10 Stored at room temperature for 0 time	8/10	80 %	2/10	20 %
	b	10 Stored at room temperature for 3 months	8/10	80 %	2/10	20 %
	c	10 Stored at - room temperature for 6 months	9/10	90 %	1/10	10 %
d	10 Stored at room temperature for 9 months	9/10	90 %	1/10	10 %	

ACKNOWLEDGMENT

The authors would give thanks to prof. Dr. Roukaya Mohamed Osman for her scientific advice and guidance to accomplish this work.

REFERENCES

1. Angus, R.D. lowe, E.L. and Pietz D.E. 1977 . Evaluation of five media for stabilization of Brucella abortus strain 19 described by lyophilization. Dev. Bio. Stand 36 :307-312.
2. Forbes, B.A.D Sohu, F and Weiss field, K.S. 1998 . Baily and Scotts diagnostic microbiology , Moslly st. Louis.
3. Hiraga, CY, Kodama , T and Chikoura, Y.T. 1990. Prevention of Human rota virus infection with chicken egg yolk immunoglobulins containing rota virus antibodies . J. Jap avoc. infect. dis. 64 :118-123
4. Ikemori, Y.M, Ohta, M, Uneda, K, Catla, S.M., Yokoy and Kodama. Y. 1997. Passive protection of Neonatal calves against bovine corona virus induced diarrhea by administration of egg yolk or colostrum antibody powder. Vet. microbiol 58 (2-4) 105-111.
5. Kuroki, M, M, Ikemori , H., Yokoyama R.C. Pernatta F.C. and kodama, Y 1993.
Passive protection against Bovine Rota virus induced diarrhea in Murine model by specific immunoglobulin from chicken egg yolk. Vet. Microbiol, 37 (1-2): 135-145.
6. Lily, S.S., Marcoss, T.N, Gihan, K.M. sly, K.A. and Ibrahim, A.M. 2004. Trials for using skimmed milk as stabilizer for attenuating Rift Valley fever virus vaccine. J. Agic. Res. 82 (4) :1905-1915.
7. Nisar, A. 1988. Post vaccination antibody profile of haemorrhagic septicemia in adult cattle. Thesis M.Sc college of Vet Science, La Hore.
8. Pietro, M and dun can, M. 2006. Salmonella infectious, clinical immunological and molecular aspects. 1st., Ed. Cambridge University press.
9. Quinn P.J. , Markery, B.K. caste, M.E., Donnelly. W.J. and leased f.c. 2002. Veterinary microbiology and microbial disease. 1st published black well, science, LTD UK.
10. Rofail, S.k. 2007. Studies on some additives for reducing some enteric bacteria infection in swiss Mice. J. Egypt Vet. Med. assoc. 67 (1) : 233-239.

11. Rofaiil, S.K. and Germine, S. Saad. 2013. Effectiveness of protective potentials of chicken egg yolk in controlling Salmonellosis using laboratory Mice as a model. *J. Agric. Res.* 91 (1) : 323 -334.
12. Rofaiil, S.K., Rogaa, A.S, ola A, E and mahmoud, m.H. 2011. Studies on some medicinal plant extract as immune stimulant in rats experimentally infected with *Salmonella typhimurium* *J. agric. Res.* 89 (1) : 336-343.
13. Stanely , A.P and Walter A.O. 1999. Vaccines, overview of vaccine manufacturing and quality assurance, 40-46 chapter 4. 3rd ed. philadelphia, London, toronto, Tokyo.
14. Tina , D. J, Adel, K, A and Ofarrelly, C. 2000. Comparative analysis of methods of purification of egg yolk immunoglobulin. *Food Agric. immunol.* 12 : 77-85.
15. Yokoyama, HRC., perolt, R, Diaz, S, Sorda, Yikenor. 1992. Panine protective effect of chicken egg yolk immunoglulinic eschericha coli infection in Neonatal piglets. *infect. immun.* 60(3) : 998-1007.

محاولات مبدئية للسيطرة علي سالمونيلا الدجاج باستخدام

صفار البيض المناعي المجفد و دراسة كفاءته

صفوت كمال روفائيل ، جرمين صبحي سعد ،

سيد بيومي خليل الجلال

معهد بحوث الأمصال و اللقاحات البيطرية- مركز البحوث الزراعية - الدقي - الجيزة

أجريت هذه الدراسة لتحضير صفار البيض المناعي المجفد و ذلك باستخدام عدة مواد مثبتة مثل اللبن منزوع الدسم، السكروز اللاكتوبومين ، السكروز الببتون و تم استخدامه للسيطرة علي مرض سالمونيلا في الدجاج و أثبتت النتائج أن هناك تأثيرا مناعيا واضحا في الدجاج التي تم عدواها بالسالمونيلا تيفيميوريم و سبق معالجته بالمنتج المحضر حيث كانت نسبة الدجاج الذي لم ينفق و لم يتأثر أن بالعدوي ٩٠ % بالمقارنة بالمجموعة الغير معالجة و اظهرت النتائج لا يوجد فروق في القوة المناعية للمنتج المحضر بالمواد المثبطة المختلفة . و عند دراسة كفاءة المنتج المحفوظ في درجات التجميد (-٢٠ م°) ، درجة حرارة الثلجة العادية ، (٤ م°) درجة الغرفة ، و تخزينه لمدة ٣ ، ٦ ، ٩ شهور تم قياس القوة المناعية لكل هذه العوامل علي حدة في الفئران المعملية بعد عدواها صناعيا بالسالمونيلا تيفيميوريم فاثبتت النتائج أن المنتج المحفوظ في درجة التجميد و المخزن لمدة ٣ شهور أحتفظ بقوته المناعية.

من خلال تلك الدراسة يمكن أن يوصي بإمكانية استخدام صفار البيض المناعي المجفد و المخزن لمدة ثلاث شهور في درجة التجميد لوقاية الدجاج من الإصابة بميكروب السالمونيلا.