



BioBacta



Journal of Bioscience and Applied Research

www.jbaar.org



Molecular pathological detection of *S. typhimurium*

by fluorescence in situ hybridization (FISH) in lambs tissue sections

Basim M. Jwad*

Bushra I. AL-Kaisei**

Department of Pathology and poultry diseases, Collages of Veterinary Medicine, University of Baghdad, Baghdad, Iraq

Address For Correspondence

*Instructor Dr. Basim M. Jwad

** Assistant Professor. Bushra I. AL-Kaisei

E-mail: bassimpathology1979@gmail.com ; bassimpatho@yahoo.com Mobile; Viber & WhatsApp: (009647903310288)

DOI: 10.21608/jbaar.2018.154699

Abstract

Despite many different methods used to diagnose *Salmonella typhimurium*, especially that depended on isolation and biochemical identification, but use technique, *Fluorescence in Situ Hybridization (FISH)*, as one method from molecular fashion probe, has been developed and applied for the direct diagnosis of *S. typhimurium* in bacterial cell smears of pure cultures, or in formalin-fixed sections, and in paraffin-embedded tissue. Therefore, this study was designed to determine the single bacterial cells in the infected intestine, liver, and gallbladder by used (Sal-3 prop), through orally administered for lambs via stomach tube, with a volume of 0.5 ml contains (1×10^8 cfu/ml *S.typhimurium*), after bacterial identification by cultures media mainly *Salmonella* chromogenic agar, and biochemical diagnosis by [Iraq-CDC/central public health laboratory (CPHL) in the Baghdad province]. So results were concluded that Sal3-prop of FISH technique, was a good instrument for fortuity *S. typhimurium* in the histopathological tissue sections.

Keywords: *Fluorescence in situ hybridization, lambs, FISH, S. typhimurium.*

Introduction:-

Salmonellosis is one of the most important zoonotic diseases, it has a direct occupational with public health and related to food-borne illness, *Salmonella enterica* subspecies *typhimurium* are one of enteropathogenic bacteria, non typhoid *Salmonellosis*, cause diseases that range from diarrhea to systemic infections in human and animals (1 ; 2). *S. typhimurium*, is a facultative intracellular pathogen, that multiplication in intestinal submucosa, and infected the mesenteric

lymph node, which spread into liver and gallbladder (3).

This pathogen has been widely prevalent, that infected animals may carry these bacteria without any clinical symptoms, so they isolated from farm animal products like meat, milk, excretion of carrier animals, also from contaminated equipment, floors at a slaughterhouse, and personnel working in an abattoir (4; 5). *S. typhimurium* had many diagnostic methods in the microbiological laboratory for recognition, by use cultivation on culture media,

Received: October 3, 2018. Accepted: December 20, 2018. Published: December 26, 2018

also with biochemical tests i.e. manually like *Analytical profile index* (API 20E/NE), or automatically test systems such as (*MicroScan-WalkAway*, *VITEK* and *BD-phoenix*), but all these procedures usually require pure cultures (6- 9) that helped pathologist to give perfect diagnosis report.

Fluorescence in situ hybridization techniques(*FISH*), was considered as one of the molecular pathological methods, help to study the single bacterial strains in complex ecological systems, through the use specifically probes targeting ribosomes in *S. typhimurium*(10).

This study aims to use the *FISH* technique via a specific probe (Sal3) generate from oligonucleotide to targeting *S.typhimurium* which infected lambs tissue organs after embedding in paraffin wax, that main in formalin-fixed section.

Material and methods

Animals used: Eight lambs, in 3-4 months old, (5.00 - 5.50 Kg) were obtained from (animal market at Al-Mahmudiya District) in Baghdad-Iraq, and it divided into:

- I- The Control group included two animals
- II- The infected group included six animals, infected by 0.5 ml contain (1x10⁸ cfu/ml *S. typhimurium*), given as a single dose, through orally administered lambs by stomach tube.

Then kept in the animal house, range in dimensions (6x8) meter, an air of the room was changed by air vacuum, under conditions were

maintained at 20± 2C°, animals were fed on animal hay, grass, and water. The experimental study was taken in the department of pathology, college of veterinary medicine; Baghdad University, for one month. Sacrifice was occurred via slaughter animals, at (week, 15 days and month) of an experiment, and pieces of tissue were taken from animal to cultivation on *Salmonella* chromogenic agar, whereas other pieces fixed in 10% formalin saline (11), remain at less for 72 hours, tissue processing, and embedded in paraffin, was occur according to (12), slides stained with (Hematoxylin and Eosin).

Dose preparation:- Infected dose and concentration, was prepared in (Unit of Zoonotic disease, Department of Microbiology/College of Veterinary Medicine /Baghdad University). According to " McFarland standards

Preparation of Salmonella chromogenic agar:- According to manufacturer instruction,(37.1g.) of dehydrated media was dissolved in (1000 ml D.W.) at 80°C. The mixture shacked and boiled for one minute to dissolve the medium completely then cooled, to about 50°C and poured into sterile Petri dishes.

Selection of probe:- The systematic name of the Sal3-probe was [L-S-Sal- 1713-a-A-18], and sequences of *Salmonella* oligonucleotide probe obtained from (13 and 14), and (Sal3)specificity, is tested by hybridization with whole bacterial cells, and the slides prepare with the prop was occur according to (15), that explain in (Table:1).

Probe	Type	Sequence (5→3)	No. of Salmonella strains detected	No. of Non-Salmonella strains detected	Specificity (%)	Sensitivity (%)	reference
Sal-3	DNA	5'-AATCACTTCACCTACGTG	96	0	100	95.5	Nordentoft, et al.(1997).

(Table:1) sequencing of *FISH* probes for detection of *S. typhimurium* in embedding tissue.

Results and Discussion:-


Bacterial isolation:- The results of isolated bacterial infection, from tissue organs, is illustrated in (figure:1) that shows the *S. typhimurium* growing

on the culture media (*Salmonella* chromogenic agar), that appear as a mauve - magenta colonies in color, during the vision. This result is installed by the diagnosis report of "Iraq-CDC/central public

health laboratory (CPHL) in the Baghdad province", which appears in (figure:2).

The magenta colonies appear due to chromogen, is a target for esterase activity present in *Salmonella*, which utilization of this chromogen results in mauve or magenta colonies in color. These

results agree with (16), which act on Isolation and Identification of several types of *Salmonella* spp. from human and animal sources in Baghdad city. Also agreement with (17), that used the molecular and conventional detection of *S. typhimurium*, Isolation from different food sources.



(Figure:1) show magenta-mauve colonies of *S. typhimurium* colonies on chromogen agar.



مختبر الصحة العامة المركزي
Central Public Health Lab



وزارة الصحة العراقية
Ministry of Health

جمهورية العراق
وزارة الصحة
دائرة الصحة العامة
العدد : د.ص.ع/٧/٤/١٧٩
التاريخ : ٢٠١٧ /٤/٢٠

برجى الإشارة الى رقم الإصدار عند الإجابة

إلى / وزارة التعليم العالي والبحث العلمي / جامعة بغداد / كلية الطب البيطري

م/ تشخيص عزلة

تحية طبية ...

أشارة الى كتابكم ذي العدد ٢٤٥ في ٢٨/٣/٢٠١٧

تم إجراء الفحوص التوكيدية على العزلة المرسله من قبل طالب الدكتوراه (باسم محمد جواد)
جامعة بغداد / كلية الطب البيطري لغرض اكمال متطلبات بحثه الموسوم (Molecular pathological

changes induced by *Salmonella typhimurium* in lambs) ، وكانت النتيجة كما مبين في ادناه .

ID#	Specimen	Diagnosis
1	Sheep	<i>Salmonella typhi-murium</i>

للتفضل بالإطلاع مع الاحترام



الدكتور
محمد جواد جواد
مختبر الصحة العامة المركزي
٢٠١٧ /٤/٢

نسخة عنه الي :

- شعبة تدريب وتطوير الملاكات / كتابكم ٢٤٥ في ٢٨/٣/٢٠١٧ / للتفضل بالإطلاع مع الاحترام
- مختبر الصحة العامة المركزي / عضو الرباط للتدريب / للمعلم لطفاً
- مختبر الصحة العامة المركزي / شعبة البكتريولوجي / وحدة البكتريا المعوية
- مختبر الصحة العامة المركزي / اصابة وحدة المعوية

(١ - ١)

٧/٩/


عنوان مختبر الصحة العامة المركزي (م. ص. ع. م.) - بغداد- ساحة الأندلس- حي النضال- محلة (١٠٣) مقابل م. الشيخ زايد للطوارئ
 البريد الإلكتروني :- cphl_baghdad@moh.gov.iq هاتف الدائرة :- ٧١٨٩٦٣٩ بـدالة ذات خمس خطوط

(Figure:2) show Baghdad-CPHL, certification for a serological test.


Result of (FISH) in paraffin embedding tissues:-

Results of microscopic detection about the presence of *S. typhimurium* by using DNA-probe(Sal3), applied by fluorescent in situ hybridization on paraffin embedding tissues of liver and intestine of lambs after oral infected 0.5 ml contain (1×10^8 cfu/ml *S. typhimurium*), were indicating the presence *S. typhimurium* inside the epithelial cell of the liver, that demonstrates in (figure:3) and (figure:4). Also within intestinal villi


in the lamina propria show the infected cells that exhibition in (figure: 5) and (figure: 6), an equally important, all examined tissue, which illustrated as green shiny color, from a signal for tissue injury through the presence of bacteria in tissue epithelial cells of intestine and gallbladder within the liver, during the period of the experiment (7, 15 and 30 days) with paraffin-embedded tissue section of lambs (liver, gallbladder, and GIT).




(Figure:3) Microscopic picture of lambs liver, which patronized with (Sal3), after 7 days from infected by, show infected cells appear as green in color → (40X)



(Figure:4) Microscopic picture of lambs liver, which patronized with (Sal3), after 7 days from infected by, shows green shiny appearance inside infected cells,  are seen indicating the presence of both components of the probe (400X).



(Figure:5) Microscopic picture of lambs intestine, which patronized with (Sal3), after 15 days from infected by, show green shiny appearance inside infected cells,  are seen indicating the presence of both components of the probe (400X).



(Figure:6) Microscopic picture of lambs intestine, which patronized with (Sal3), after 30 days from infection by, show green shiny appearance inside infected cells, \Rightarrow are seen indicating the presence of both components of the probe (400X).

On the other hand, The scores of green signals appear in (table:2), were indicating the presence of *Salmonella typhimurium* in the organ of the animals during the period of the experiment (7, 15, and 30 days) with paraffin-embedded tissue section of lambs (liver and GIT). Whereas reduced autofluorescence from the color of the tissue, that appears red in the background, easily for detection of single infected bacteria.

Scores during (7 days), was highest after infected by *Salmonella typhimurium*, so the scores with paraffin-embedded tissue were recorded in all infected animals (100%), so it has a sign that *S.*

typhimurium infection with tissue organ was more severe in the period of infection during the first seven days of infection. However, notes that the two groups were equalized by the severity of the infection and spread the bacteria in the body within the period of 15 and 30 days after liver infection, wherefrom (table:2) in the intestine, we note that the proportions of the score were compared among them whether the infection with presence in the liver or intestine, so at end of the 15 days the green signals were found in all score that main the spread within animal from each observation, was recorded (100%) that recorded after 30 days.

Table(2): The scores of the (Sal3) DNA probe

Time	A green signal score of a paraffin-embedded tissue section of intestine lambs infected by 10^8 <i>S. typhimurium</i>			A green signal score of a paraffin-embedded tissue section of liver lambs infected by 10^8 <i>S. typhimurium</i>		
	Score 1	Score 2	Score 3	Score 1	Score 2	Score 3
7 days	0	1 B c	1	1	1 A c	1
15 days	3	0 A b	0	3	0 A b	0
30 days	0	1 B a	2	3	0 A a	0

Small letters significant differences between a group at ($p \leq 0.01$)

Capital letters significant differences with a group at ($p \leq 0.01$).

The results that show a shiny green color appear in the infected epithelial cell, occur due to (Sal3) prop within the FISH technique, was used to identification of *S. typhimurium* through alignment of mRNA gene sequences of representative Salmonella sequences, which characterized by the percentage of specificity to Salmonella strains detected was (100%) and the sensitivity was (95.5%), and it did not hybridize with any non-Salmonella sequence.

Furthermore, probe reaction with two regions from sequence variability in bacteria, which have previously been shown to exhibit high degrees of conservation in the probe-target region, that making detection and spatial localization of single bacterial cells in tissue samples.

Also used the paraffin embedding tissue with these probes, because the fixative tissue in paraffin wax, help to opening up the bacterial wall for probe penetration, also it protecting the ribosomes from degradation by endogenous RNase activity, and enhance the level of discrimination of the signal from single bacteria in tissue sections, and simultaneous excitation in two separate bands of the DNA, therefore enabling it to reduce autofluorescence from tissue considerably by

turning the color of the tissue red and thereby allowing for the detection of single bacteria.

This finding was supported by results reported by (18) in a dog; (19 - 21) they explained that "the development of a [fluorescence-labeled specific oligonucleotide probe] makes the (FISH) technique a promising tool for the rapid identification of *S. enterica* in bacterial smears, as well as for the detection of *S. enterica* in histological tissue sections".

We have verified in the present study investigated that used DNA-probe(Sal3), applied by fluorescent in situ hybridization on paraffin embedding tissue of liver and intestine of lambs after oral infected 0.5 ml contain (1×10^8 cfu/ml *S. typhimurium*), give a stability of the ribosome target allowed "*mRNA-DNA-S.typhimurium*", for the detection of single cells in clinical infection tissue, therefore we suggest that the FISH technique can be used for the detection of *S. typhimurium*, inside infective tissue after fixation and embedding in paraffin wax.

Conflict of Interests Statement:- The authors declare that there is no conflict of interest regarding the publication of this article.

Financial Disclosure Statement:- This study was not funded by any funding agency and was

conducted at the expense of the researcher. As part of the research requirements for the degree of Doctor of Philosophy in the Department of Pathology.

Animal Rights Statement:- The experiments and procedures involving local lambs, were approved by a scientific committee of the department of pathology; in the College of Veterinary Colleges of Veterinary Medicine, University of Baghdad, Iraq, and were conducted according to the guidelines of the committee.

References:-

1. Atyabi, N.; Zahraei Salehi, T.; Ghazisaeedi, F. and Ashrafi, I. (2012). The molecular investigation of widespread *Salmonella* serovars, *S. typhimurium*, and *S. enteritidis*, involved in salmonellosis of cattle and sheep.
2. Roaa Adnan Hussein (2016). Comparison study for treatment of enteritis in lambs induced by *Salmonella typhimurium* as compared colicin E1 or enterocin A1 with enrofloxacin. Degree of doctor of philosophy in veterinary medicine; College of veterinary medicine. Baghdad University. Iraq.
3. Aseel Jasim Mohammed (2017). Study the role of grape seeds polyphenols as anti-bacterial, anti-inflammatory, anti-apoptotic and immunomodulatory agent in mice Infected with *Salmonella typhimurium*. A dissertation degree of doctor of philosophy veterinary medicine/microbiology; Collage of veterinary medicine, Baghdad University. Iraq.
4. Narvaez-Bravo, C.; Rodas-Gonzalez, A.; Fuenmayor, Y.; Flores-Rondon, C.; Carruyo, G.; Moreno, M.; Perozo-Mena, A. and Hoet, A. E. (2013). *Salmonella* on feces, hides and carcasses in beef slaughter facilities in Venezuela. Intern. J. Food. Microbiol., 166: 226–230.
5. Paszkiewicz, W. and Pyz-Lukasik, R. (2010). Bacterial contamination of carcass surfaces in relation to the order of slaughtering cattle. Med. Weter. 66: 51–53.
6. O’Hara, C.M.; Tenover, F.C. and Miller, J.M. (1993). Parallel comparison of accuracy of API/20E, Vitek GNI, MicroScan Walk/Away Rapid ID, and Becton Dickinson Cobas Micro ID-E/NF

- for identification of members of the family Enterobacteriaceae and common gram-negative, non-glucose-fermenting bacilli. J. Clin. Microbiol. 31, 3165–3169.
7. Wain, J. and Hosoglu, S., (2008). The laboratory diagnosis of enteric fever. J. Infect. Dev. Ctries. 2, 421–425.
8. Snyder, J.W.; Munier, G.K. and Johnson, C.L. (2008). Direct comparison of the BD-phoenix system with the MicroScan WalkAway system for identification and antimicrobial susceptibility testing of Enterobacteriaceae and nonfermentative gram-negative organisms. J. Clin. Microbiol. 46, 2327–2333.
9. O’Hara, C.M., (2005). Manual and automated instrumentation for identification of Enterobacteriaceae and other aerobic gram-negative bacilli. Clin. Microbiol. Rev. 18, 147–162.
10. Licht, T.R.; Krogfelt K.A.; Cohen, P.S.; Poulsen, L.K.; Urbance, J. and Molin, S. (1996). Role of lipopolysaccharide in colonization of the mouse intestine by *Salmonella typhimurium* studied by in situ hybridization. Infect. Immun. 64:3811–3817.
11. Carson F L, and Hladik Ch. Histotechnology.: A Self-Instructional Text. Third Ed., Chicago ASCP Press, 2009. P:400.
12. Luna, L.G.: "Manual of histologic staining methods of the armed force institute of pathology" 3rd Ed. McGraw-Hill, New York. USA. , 1968.
13. Nordentoft, S.; Christensen, H. and Wegener, H.C. (1997). Evaluation of a fluorescence labelled oligonucleotide probe targeting 23S rRNA for in situ detection of *Salmonella* serovars in paraffin-embedded tissue sections and their rapid identification in bacterial smears. J. Clin. Microbiol. 35, 2642–2648.
14. Alm, E.W.; Oerther, D.B.; Larsen, N.; Stahl, D.A. and Raskin, L. (1996). The oligonucleotide probe database. Appl. Environ. Microbiol. 62:3557–3559.
15. Steen Nordentot ; Henrik Christenen and Henrik Caspar Wegener. (1997). Evaluation of a fluorescence-labeled oligonucleotide probe

targeting 23S rRNA for in situ detection of Salmonella serovars in paraffin-embedded tissue sections and their rapid identification in bacterial smears. *Journal of Clinical Microbiology*. 35(10):2642–2648.

16. Alyaa Abdulhussein Kadhim. (2014). Isolation and identification of Salmonella spp. from human and animal sources in the Baghdad city. Degree of Master of Science in Biology/Microbiology. College of Science for Women, Baghdad University. Iraq.

17. Ali Karim Husuni Al-Bakri. (2015). Molecular and Conventional detection of Salmonella typhimurium in different food sources. Degree of master in genetic engineering and biotechnology GEB. Genetic engineering and biotechnology institute for postgraduate studies/ Baghdad University. Iraq.

18. Antonio Giuliano; Thelma Meiring; Andrew J. Grant and Penny J. Watson. (2015). Acute Hepatic Necrosis Caused by Salmonella enterica Serotype I 4,5,12:-:1,2 in a Dog. *Journal of Clinical Microbiology*. 53(11): 3674 –3676.

19. Hagen Frickmann B. ; Andrea Hänle ; Andreas Essig ; Denise Dekker ; Kennedy Boahene ; Samuel Acquah ; Nimako Sarpong ; Yaw Adu-Sarkodie ; Norbert G. Schwarz ; Jürgen Mayd ; Florian Marksg ; Ralf M. Hagen and Sven Poppert. (2013). Fluorescence in situ hybridization (FISH) for rapid identification of Salmonella spp. from agar and blood culture broth—An option for the tropics. *International Journal of Medical Microbiology* (303):277– 284.

20. Almeida C. ; Azevedo N.F. ; Fernandes R.M. ; Keevil C.W. and Vieira M.J. (2010). Fluorescence In Situ Hybridization method using a peptide nucleic acid probe for identification of Salmonella spp. in a broad spectrum of samples. *APPL. ENVIRON. MICROBIOL.* 76(13):4476–4485.

21. Bledar Bisha. (2009). Fluorescence in situ hybridization-based detection of Salmonella spp. and *Listeria monocytogenes* in complex food matrices. A dissertation for the degree Doctor of philosophy. Iowa State University; Ames, Iowa. USA.