

CHARACTERIZATION OF POULTRY PRODUCTION SYSTEMS IN THE RURAL SECTOR OF FAYOUM

M.A. El-Wardani, Y.A. Abdel-Aziz, Amal S. Omar, A.H. Abdelmaged and O.M. Zatter

Animal Production Research Institute, Agricultural Research Centre, Ministry of Agriculture, Dokki, Giza, Egypt

SUMMARY

One hundred and twenty-one rural poultry farmers in 12 villages in six districts of Fayoum governorate were studied through semi-structured interviews with questionnaires. The objectives of this study were to describe the existing village poultry production systems. Rural poultry farmers were identified as those who raise flocks either inside their houses or in attached enclosures, beside small farms who operated on semi-commercial basis under rural conditions. The chi-square was used to test all differences between systems except flock size data which allowed making ANOVA between systems, districts and villages between districts.

Systems identified were: 1) Backyard or family poultry production system; involves two sub-systems: traditional and landless systems, and 2) semi-commercial village poultry production system. Backyard system represented about 76% of the studied farms versus 24% for the semi-commercial system. Chickens represented the highest component of the flock composition (82%) followed by pigeons and ducks (8.2% and 8 %), while geese, rabbits and turkeys represented minor percentages of 1%, 0.7% and 0.1%, respectively. Local breeds are the dominating breeds in all systems. Average flock size was 70 ± 6 and 70 ± 5 birds in the traditional and landless systems, respectively, versus 1322 ± 259 birds in the semi-commercial system. Most of the farmers (58-67%) in traditional and landless systems utilize poultry products for family consumption. In the semi-commercial system, only 10% of the products are used for home consumption and the rest (90%) goes to the ordinary market channels. Poultry are usually housed in primitive coops (73-76%) in traditional and landless systems. While in the semi-commercial system, poultry were generally raised in a room inside the house (66 %) or in a small pen attached to the house (34 %). Family labour is usually used. Disease control and unavailability of feed ingredients are major problems facing poultry production in the rural sector.

Keywords: *Village poultry production systems, landless, backyard, semi-commercial*

INTRODUCTION

In Egypt, family poultry production is the dominant system and is a part of the rural life. It has been one of the support systems to subsistence farmers, providing supplementary food and income which are badly needed in rural areas. Most families

keep poultry in the backyard or on rooftop. The exact number of the rural poultry population, backyard family production, rooftop systems etc. is not known (Hosny, 2006). Chicken production in the rural sector is estimated at about 99.430 million broilers (17 % of the total national production) and 1.2 billion eggs (29% of the total national production). There is no published data available on number of ducks, geese, turkeys, rabbits and pigeons (MALR, 2005). However, the rural sector is almost the sole source of ducks, geese and pigeons. A rural flock may hold different species of poultry, but chickens are mainly kept for egg and meat production, whereas turkeys, ducks, geese, rabbits and pigeons are mainly kept for meat production. Rural flock size can range from 10-20 birds up to a few hundreds (Hosny, 2006). Rural poultry sector depends mainly on local and improved local breeds.

Family poultry production in general and village chickens in particular represent a significant part of the rural and national economies (Gunaratne *et al.*, 1993; Panda and Mohapatra, 1993; Guèye, 1998; Sonaiya *et al.*, 1999 Guèye, 2000). According to a household expenditure survey for Egypt, poultry products account for nearly one third of the expenditure on animal protein products and account for 31 percent of the total food bill (AAFC, 2004). Sonaiya (1990) suggested the need to develop systems approaches to rural poultry development, and Lee *et al.* (1993) indicated that only by systems analysis, the production system could be better understood and interventions for improvement of production can be determined.

There is little available information about the management, constraints, and the productivity of rural poultry flocks and technological improvements that could be affordable to the low-input systems. The aim of this study –therefore- was to use systems approach to describe the existing village poultry production systems in rural areas and obtain reliable data on these systems in Fayoum governorate.

MATERIALS AND METHODS

A field survey was conducted on 121 randomly selected farmers in twelve villages in six districts of Fayoum governorate in middle Egypt. The number of poultry farmers surveyed in each of these villages is shown in table1. Data were collected as part of research study on development of market-oriented poultry production systems at the smallholders in rural areas, funded by the Egyptian National Academy of Scientific Research and Technology. The data were collected during the period from March to August, 2007.

A preliminary survey was conducted at the beginning to identify the village based poultry production and pilot-examine the survey formats. Poultry farmers in rural areas were identified as those operate in a village and raise flocks either inside or attached to their houses. They adopted simple management practices of poultry raised under rural conditions. One extension officer in each village was trained and assigned to collect data under supervision of the research team through weekly visits to the poultry farmers. The collected data included information on flock size, flock composition, flock structure, type of poultry, breeding purpose, housing systems, marketing, labour and constrains to improvement.

Enumeration data of the field survey were analyzed by the chi-square test of hypothesis and the Marascuillo procedure was used to test the significance of the proportions among systems (XLSTAT 1.01 computer program, 2009). The data collected on flock size were statistically analyzed by the least squares technique

using the general linear model procedure of SAS program (SAS, 2005). The following linear model was used in the analysis:

$$Y_{ijk} = \mu + S_i + D_j + V_k(D_j) + e_{ijk}, \text{ where}$$

Y_{ijk} is the observed flock size,

μ is the general mean,

S_i is the effect due to production system, $i = 1, 2, 3$ (1=Traditional, 2=Landless and 3=Semi-commercial),

D_j is the effect of the j district ($j = 1, 2, 3, 3, 4, 5, 6$),

$V_k(D_j)$ is the effect of the k village within district j , ($k=1, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12$),

e_{ijk} is a random effect associated with the individual observation and assumed to be independent, random and normally distributed.

Table1. Number of poultry farmers surveyed at different villages

Districts	Villages	Farmers
El-fayoum	Zawia Elkerdasa	9
	Elazab	10
Snors	Elkaapy	10
	Sanhor	11
Tamia	Kaffer Mahfouz	11
	Pander Tamia	10
Epshway	Abo-Denkash	11
	Abo-kesaa	10
Elseddeeq	Kaser Elgebaly	9
	Batn harred	10
Atssa	Elgaafra	10
	Gerdo	10

RESULTS AND DISCUSSION

Based on criteria of capital investment degree (extensive, semi-intensive or intensive) and the economic purpose of the poultry owner (subsistence, semi-commercial or commercial), the poultry production systems in rural areas can be classified -in general- into two main systems, 1) Back-yard or family poultry production system that involves two sub-systems; traditional and landless village poultry production systems and 2) Semi-commercial or small-farm poultry production system.

A. Brief description of the systems

1. Back-yard poultry production system: Back-yard or family poultry production is the prevailing system in nearly all the rural sector. The economic purpose of the poultry farmer is mainly to meet family needs (home consumption). The system involves little semi-subsistence-oriented production. In terms of capital investment it is considered as extensive system. Backyard poultry production system represented about 76% of the studied farms. Flock composition is heavily skewed towards chickens. This system involves two sub-systems; traditional and landless village poultry production systems.

1.1. Traditional village poultry production system: A mixed system where livestock, including animals; poultry and crop production are integrated in the same farm. Poultry owners in this system have access to cultivated area for crop production with an average of 1.7 Fadden (1 Fadden = 4200 m²). Poultry owners kept a limited number of different species of poultry, mainly chickens followed in numbers by ducks, geese, pigeons, turkeys and rabbits. This system represented about 41% (50 farmers) of the total farmers surveyed. Around 49 % of the families worked in the farm permanently and earned their living from agriculture. The other 51 % of the households had permanent jobs out of the farm as employees and worked on their farms as secondary jobs.

1.2. Landless village poultry production system: This system is common in the vicinities of the relatively large towns in rural areas. Poultry farmers in this system do not have access to cultivated areas. Poultry farmers kept birds of varying ages and different species (mainly chickens, followed in numbers by ducks, geese, pigeons, turkeys and rabbits). Poultry owners represented about 35% (42farmers) of the total farmers surveyed. The main profession for the largest portion of them is as employees or workers (82 %), the remaining (18%) work as poultry farmers. Among the landless families in rural areas particularly women, poultry used to provide independent income for the family in most cases. The importance of poultry in income generation especially for the poor and landless is quite evident in the study area. In rural Egypt, poultry account for 72 % of the total livestock income; chicken alone account for 61% of the livestock income (Croppenstedt, 2006).

2. Semi-commercial poultry production system: The semi-commercial poultry production system is rather market-orientated; therefore this system could be looked at as a transitional stage towards the commercial poultry production system. Poultry farmers who are involved in this system have to some extent management and marketing skills. It seems that more access to the know-how and capital are important factors for the development of this production system. The flock size is larger than the in other rural systems with four species being raised in this system (mainly chickens, followed by ducks, pigeons and rabbits). Poultry owners represented about 24% (29 farmers) of the total farmers surveyed. The majority of the poultry farmers are employees (65 %); the remaining (35%) work as only poultry farmers.

Another point of view was presented by Bessei (1987) and Sonaiya (1990) who classified poultry production systems into: free-range system or traditional village system, backyard or family system, semi-intensive system and intensive system. According to Gueye (1998a), the free-range system or traditional village system and backyard or family system are the most commonly practiced in rural Africa. Gueye (1998a) added that more than 85% of the rural families in sub-Saharan Africa keep one or more species of poultry.

B. Main Features of the systems:

1. Family labour: All family members including women, children and men tend to be involved in rural poultry production. Women, assisted in some cases by children, play a key role in the family labour, especially in the traditional and landless poultry

production systems. Daily managerial practices depend mainly on the women in 65% and 70% of the surveyed poultry farmers in traditional and landless systems, respectively. However, in the semi-commercial system farms depend mainly on men in 57% of the surveyed poultry farms as shown in table 2.

Table 2. Family labour participation in poultry management in the different production systems (expressed as percentage of system totals)

Items	Back-yard		Semi-commercial	P-value
	Traditional	Landless		
Women only	65 ^a	70 ^a	28 ^b	0.0001
Men only	22 ^{ab}	10 ^a	57 ^b	0.0006
Family	13 ^a	20 ^a	15 ^a	0.67

Means with different letters within the same rows are significantly different.

In Egypt, poultry raising is a popular activity among rural women (Hosny, 2006). The same trend has been observed by Sonaiya, (2000) who stated that Nigeria family poultry is usually the responsibility of women. A major portion of backyard poultry production in the village is managed and implemented by women. In sub-Saharan Africa, more than 70% of the chicken owners are women, while traditionally pigeons are the responsibility of only children (Gueye, 1998b).

2. Types of poultry: Native breeds of poultry of different species are mainly kept followed in numbers by improved native breeds and exotic breeds as shown in table 3. The largest percentage of chicken of native breeds such as Fayoumi, Balady and Dandarawy (77% and 61%) are kept in traditional and landless systems, respectively. On the other hand the semi-commercial system included the largest percentage of improved native breeds such as Dokki4, Mandarah, Montazah, Matrouh, Bandara, El-Salam and Baheig (51 %) and exotic broiler strains (33%).

Table 3. Poultry species* and breeds raised in the different production systems (expressed as percentage of the system totals)

Poultry species		Back-yard		Semi-commercial	P-Value
		Traditional	Landless		
Chickens	Native breeds	77 ^a	61 ^a	16 ^b	0.0001
	Improved Native breeds	23 ^a	39 ^a	51 ^c	0.0001
	Exotic Broiler strains	---	---	33	---
Ducks	Native breeds	44 ^a	48 ^a	---	0.47
	Exotic breeds	56 ^{ab}	52 ^a	100 ^b	0.0001
Rabbits	Native breeds	60 ^a	100 ^b	---	0.0001
	Exotic breeds	40 ^a	---	100 ^b	0.0001
Geese	Native	100	100	100	---
Turkey	breeds	100	100	100	---
Pigeon	Native	100	100	100	---

Means with different letters within the same columns are significantly different.

*Including geese, turkeys and pigeons were native breeds only.

This emphasizes the importance of the native breeds for rural poultry production. This could be due to its tolerance to harsh weather conditions, lower feeding requirements and also to consumer preferences of the taste of eggs and meat of native breeds. No specific breed could be pinpointed in the rural poultry sector as it depends mainly on the local non-specified crosses between endogenous native breeds such as Fayoumi, Balady and Dandarawy or improved native breeds (Hoseny, 2006).

Duck farmers in traditional, landless and semi-commercial systems have the same attitude towards keeping exotic breeds in 56% and 52% and 100% of cases, respectively. Exotic duck breeds such as Muskovy and Pekin as shown in table 3. Similar trend has been observed in favour of the native rabbits breeds such as Balady White and Balady Red versus exotic rabbit breeds such as New-Zealand and Chinchilla. In the surveyed farms, only native breeds of geese, turkeys and pigeons were found. This reflects the preference of family poultry producers in rural sector to keep native breeds of these types of poultry.

3. Flock composition: Rural poultry farmers raised different species of birds, mainly chickens, followed in number by ducks, geese, pigeons and little numbers of turkeys and rabbits as shown in table 4. Chicken represented the highest population of the flock composition (82.0%) followed by pigeons (8.2%) and ducks (8.0%), while geese, rabbits and turkeys represented minor percentages of about 1.0%, 0.7% and 0.1% , respectively of the total numbers of poultry on the farms.

Table 4. Proportions of the farms raising different species of poultry under the studied production systems (expressed as percentage of system totals)

Poultry species	Traditional	Landless	Semi-commercial	P-Value
Chicken	58 ^a	68 ^a	85 ^b	0.0001
Ducks	18 ^a	18 ^a	6 ^b	0.0001
Geese	9 ^a	6 ^b	---	0.0001
Turkey	1 ^a	0.5 ^b	---	0.0001
Pigeon	12 ^a	7 ^a	8 ^b	0.0001
Rabbit	2 ^a	0.5 ^a	1 ^b	0.0001

Means with different letters within the same rows are significantly different.

Chickens constituted 58% and 68% of the flock composition in traditional and landless poultry production systems, respectively versus 85% in the semi-commercial poultry production system. Family poultry flock composition is heavily skewed towards chickens in Africa as more than 85% of the rural families in sub-Saharan Africa keep one or more species of poultry (Gueye, 1998), and towards ducks in Asia and turkeys in Latin America (Brabnckaert and Gueye, 1999).

4. Flock size and structure: Flock size is more related to the objectives of the poultry farmer. The average flock size was 70 birds in both of backyard systems, and ranged between 13 and 199 birds and between 20 and 180 birds in the traditional and the landless poultry production systems, respectively. On the other hand, the average flock size of chickens in the semi-commercial system was 1322 birds, and ranged from 100 to 5000 birds (Table 5). This could be due to that poultry owners in the system are more market oriented and have access to market channels. The wide

variation noted in the flock size in many rural areas, depends on household objectives (home consumption, income generating or both) and investment.

Table 5. Least squares means \pm standard errors of flock size under different production systems (birds)

Items	Back-yard		Semi-commercial
	Traditional	Landless	
Poultry flock size	70 \pm 6 ^b	70 \pm 5 ^b	1322 \pm 259 ^a
Chicken flock size	41 \pm 5 ^c	47 \pm 5 ^b	1235 \pm 226 ^a
Duck flock size	16 \pm 2 ^b	14 \pm 1 ^c	575 \pm 312 ^a
Geese flock size	12 \pm 1 ^a	13 \pm 4 ^b	---
Turkey flock size	9 \pm 4 ^a	7 \pm 4 ^b	---
Pigeon flock size	19 \pm 4 ^a	13 \pm 3 ^b	---
Rabbit flock size	9 \pm 3 ^b	5 \pm 1 ^c	125 \pm 3 ^a

Means with different letters within the same columns are significantly different ($P < 0.05$).

*Including chickens, ducks, geese, turkey, pigeons and rabbits.

Flock size was reported in some studies in Egypt to range from 10-20 birds up to a few hundreds depending on the objectives of the farmers (Hoseny, 2006). Flock size ranged from 4-130 birds in Philippines (Lambio, 2005) while in South East Asia, flock size ranged from 10 to 50 birds (Aini, 1999). Household flock sizes range from 3 to 97 birds in Africa, 10 to 31 in South America and from 50 to 2,000 in Asia (Brabnckaert and Gueye, 1999). The wide variations in rural flock size could be attributed to production system and local factors (Kuit, 1986).

With respect of flock structure, data were available only on chickens since other species were found in scattered small numbers. The largest flock size was that of the semi-commercial being 1235 birds with 100% young chicks and the smallest was that belonging to the traditional system (41 birds, of which 44 % were young chicks). The flock size for the landless system was 47 birds, of which 61% were young chicks. The high percentage of chicks in the whole flock as compared to mature hens (40%, 32%), cocks (4%, 3%) and pullets (12%, 4%) for both of the traditional and landless systems, respectively could be due to the high mortality rate during incubation period. In general, the proportion of hens in the flock is an indication of egg and meat production of the farm (Mwalusanya, 1999, Abdou, 1992 and Wilson, 1987).

Table 6. Analysis of variance of flock size by system, district and village within district

Source of variation	DF	MS
System	2	13.18 ^{***}
District	5	0.18 ^{NS}
Villages within district	6	0.37 ^{**}
Error	107	0.08
Corrected total	120	----

*** $P < 0.0001$, ** $P < 0.01$ and NS=not significant.

Analysis of variance showed significant effects among systems ($P < 0.0001$), and between villages within districts ($P < 0.01$) as shown in table 6. However no

significant differences were detected among districts which indicate the similarity among them as far as poultry production is concerned.

5. Source of chicken: For the same reason mentioned in the discussion of flock structure, source was studied only for chicken. Table 7 shows that traders are the main source of chickens (mainly day-old chicks) in the different production systems. Traders are the source of 41%-61% of the chickens in the surveyed farms depending on the system. This reflects the importance of poultry traders in rural areas. Local hatcheries are the only source of chickens in 5-14% of the cases. Governmental sites in Fayoum such as Al-Azzab integrated project and poultry research stations are good source for chickens in the Semi-commercial and landless systems (23% and 48%, respectively).

Table 7. Source of chickens in the different production systems (expressed as percentage of the system totals)

Source of chickens	Production system			P-Value
	Traditional	Landless	Semi-commercial	
Owner's flock	21 ^a	31 ^a	---	0.57
Traders	61 ^c	41 ^a	43 ^a	.0001
Hatcheries	14 ^a	5 ^b	9 ^b	.0001
Governmental sites	4 ^{bc}	23 ^a	48 ^a	.0001

Means with different letters within the same rows are significantly different.

6. Utilization of poultry and their products: Poultry owner's objective is usually more related to the production system. Table 8 shows that the two first systems (traditional and landless) are mainly directed towards home consumption (67% and 58%, respectively), and the surplus is sold in the local markets to increase family income. In the traditional system, 23% of the farmers sold most of their poultry and poultry products directly to the consumers in village and urban markets, where 10% of them sold their products to the traders. A similar trend was observed in the landless system as indicated in table 8. The semi-commercial system is more market-oriented. Around 50 % and 40 % of the farmers in the semi-commercial system preferred to sell their products to regular markets and traders, respectively. The remaining 10 % keep poultry and poultry product for home consumption.

Table 8. The utilization of poultry and their products in the different production systems (expressed as percentage of the system totals)

Production Systems		Home consumption	Market	Traders
Back-yard	Traditional	67 ^a	23 ^a	10 ^a
	Landless	58 ^a	25 ^{ab}	17 ^a
Semi-commercial		10 ^b	50 ^b	40 ^b
P-Value		.0001	.010	.0001

Means with different letters within the same columns are significantly different.

7. Housing: Poultry in the traditional and landless systems are usually housed in primitive coops that are built from locally available material in rural areas such as mud bricks and palm wood, reed or plant stalks. In most cases, these houses; are

located either on the rooftops or attached to the house have no artificial light, and with small windows. The coop represents the largest percentage (73% and 76 %) for the traditional and landless systems, respectively, while in the semi-commercial system, poultry were generally kept in a suitable room inside the house (66 %) or in a small poultry house (34 %) as indicated in table 9.

Table 9. Types of housing under the different production systems (expressed as percentage of the system totals)

Production Systems		Coop	Room inside the house	Small poultry house
Back-yard	Traditional	73 ^a	18 ^a	9 ^a
	Landless	76 ^a	11 ^a	13 ^a
Semi-commercial		---	66 ^b	34 ^b
P-Value		0.39	.0001	.0001

Means with different letters within the same columns are significantly different.

8. Constraints to improvement: Feeding is a major problem in 20-25% of the surveyed farms followed by diseases and high mortality (20%-24%) in the different production systems; farms suffering from low production, lack of good incubation and housing facilities, and unavailability of appropriate poultry breeds represent together a considerable percentage of the surveyed farms in all systems (Table 10). Lack of equipment represented a minor problem (1-2%) to landless and traditional systems, respectively. However, in the semi-commercial system the lack of equipment is a problem for 12% of the farms as the farmers purchase feeders and drinkers which are considered expensive.

Table 10. Type of problems facing rural poultry farmers in the different production systems (expressed as percentage of system totals)

Items	Back-yard		Semi-commercial	P-Value
	Traditional	Landless		
Feeding	22 ^a	25 ^b	20 ^b	.0001
Diseases and mortality	24 ^{ab}	23 ^a	20 ^a	.001
Low production	21 ^a	21 ^b	7 ^c	.0001
Incubation facilities	12 ^a	13 ^b	12 ^a	.001
Breeds	10 ^a	8 ^a	12 ^a	.45
Housing	9 ^a	9 ^b	7 ^a	.0001
Equipment	2 ^a	1 ^a	12 ^b	.0001
Marketing	---	---	10	---

Means with different letters within the same columns are significantly different.

Traditional and landless systems have no problems in marketing and they are able to sell their products directly to the consumer or in the village market. Marketing problems were found in 10% in the semi-commercial system.

ACKNOWLEDGEMENT

This work is part of a study on the “development of market-oriented poultry production systems at the smallholders in rural areas”. The authors would like to

express their gratitude to the financial assistance provided to this study by the Egyptian National Academy of Scientific Research and Technology. Also, acknowledge the assistance received from Dr. M. A. Ibrahiem and Dr. U. El-Saied in the statistical analysis and Prof. S. Galal for reading of the manuscript and valuable advice.

REFERENCES

- AAFC, Agriculture and Agri-Food Canada, 2004. Market Information Africa and the Middle East, July 2004.
- Abdou, I., Bell, J.G., 1992. Dynamique de la volaille villageoise dans la region de Keita au Niger. In: Village poultry production in Africa, proceedings of an international workshop held in Rabat, Morocco, 7-11 May, 0 6-11.
- Aini, I., 1999. Village chicken production and health: South-east Asian perspectives. FourthAsia-Pacific Poultry Health Conference. <http://www.avpa.cia.au/confer/ideris.htm>.
- Bessei, W., 1987. Ttendencies of world poultry production . Proceeding of the third International dLG- Symposium on Poultry Production in Hot Climates , June 19-22, Hameln, Germany .
- Branckaert,R.D.S. and E.F. Gueye, 1999. FAO,s programme for support to family poultry production poultry as a tool in poverty eradication and gender equality- Proceedings of a workshop, March 22-26, 1999, Tune Landboskole, Denmark.
- Croppenstedt, A., 2006. Household Income Structure and Determinants in Rural Egypt, ESA Working Paper No. 06-02, FAO, Rome.
- Guèye, E.H.F., 1998. Village egg and fowl meat production in Africa. World's Poultry Science Journal 54: 73-86.
- Guèye, E.H.F., 2000. The role of family poultry in poverty alleviation food security and the promotion of gender equality in rural Africa. Outlook on Agriculture 29: 129-136.
- Gueye,E.F., 1998. Poultry plays an important role in Africa Village Life. World Poultry , vol.14,NO,14-17.
- Gunaratne, S.P., A.D.N. Chandrasiri, W.A.P. Mangalika Hemalatha, J.A. Roberts, 1993. Feed resource base for scavenging village chickens in Sri Lanka. Tropical Animal Health Production 25: 249-257.
- Hosny,F.A., 2006. the structure and importance of the commercial and village based poultry system in Egypt. FAO publications. <http://www.fao.org/docrep/008/af840e/af840e00.htm>.
- Kuit, H.G., A. Traore and R.T. Wilson, 1986. Livestock production in Central Mali: Ownership, management and productivity of poultry in traditional sector, Trop. Anim. Hlth Prod.18 222-231.
- Lambio, A.L., 2005. "The Future Prospects for Smallholder Native Chicken Producers in the Philippines". Elpidio Quirino Professorial Chair Lecture. 24 June. ADSC Lecture Hall, UPLB-CA.
- Lee, J.V.D., H.M.J. Udo and B.O. Brouwer, 1993. Design and validation of animal traction module for smallholder livestock systems simulation model. Agricultural Systems 43: 199- 227.
- MALR, 2005. Ministry of Agriculture and Land Reclamation - Bulletin of Economic Affairs Sector, Egypt (In Arabic).

- Mwalusanya, N.A., 1999. Productivity and nutritional status of local chickens under village management conditions, INFPD Newsletter 2 18-20.
- Panda, B., and S.C. Mohapatra, 1993. Poultry development strategies in India. *World's Poultry Science Journal* 49: 265-273.
- S.A.S., 2005. SAS User's Guide: Statistics. SAS Inst., Inc., Cary, NC.
- Sonaiya, E.B., 1990. The context and prospects for development of smallholder rural poultry production in Africa. In: CTA seminar proceedings, volume 1. *Smallholder Rural Poultry Production*, Thessaloniki, Greece, pp. 35-52.
- Sonaiya, E.B., R.D.S Branckaert and E.F. Guèye, 1999. Research and development option for family poultry. First INFPD/FAO Electronic Conference on Family Poultry.
- Sonaiya, E.B., 2000. Backyard poultry production for socio- economic Advancement of the Nigeria Family: Requirement for Research and Development. *Nigeria Poultry Science Journal* 1: 88- 107.
- Wilson, R.T., A. Traore, H.G. Kuit, and M. Slingerland, 1987. Chick mortality in scavenging village chickens in Sri Lanka, *Trop. Anim. Hlth Prod.* 19 229-236.
- XLSTAT 1.01 computer program, (2009).

توصيف نظم الإنتاج الداجني في القطاع الريفي في محافظة الفيوم

محمد عبد العزيز الورداني، ياسر أحمد عبدالعزيز، آمال صالح عمر، أحمد حسين عبد المجيد،
أسامه محمود محمد زعتر

معهد بحوث الإنتاج الحيواني، مركز البحوث الزراعية، وزارة الزراعة، الدقي، جيزة، مصر

أجريت الدراسة في محافظة الفيوم على 121 مربيا للدواجن في ١٢ قرية تابعة لعدد ٦ مراكز. وتم تعريف مربى الدواجن الريفي في هذه الدراسة بأنه المربي الذي يحتفظ بالقطيع الداجني داخل المنزل/ ملاصق للمنزل أو في مزرعة صغيرة في النظام شبه التجاري الريفي. تم تجميع البيانات عن طريق إستمارة استبيان مع مقابلات شخصية. وكان الهدف من هذه الدراسة هو تحديد أنظمة إنتاج الدواجن السائدة في القطاع الريفي وتوصيفها. أجري تحليل مربع كاي للبيانات لاختبار الفروق بين الأنظمة فيما عدا حجم القطيع حيث سمحت البيانات المتعلقة به بإجراء تحليل التباين بين الأنظمة و بين المراكز وبين القرى داخل المراكز.

ميزت الدراسة نظامين رئيسيين لإنتاج الدواجن في القطاع الريفي: (١) نظام التربية المنزلية الريفية (العائلي) ويندرج تحت هذا النظام نظامان فرعيان هما النظام التقليدي الريفي ونظام التربية بدون حيازة زراعية. (٢) نظام التربية شبه التجارية الريفية. شكل نظام التربية المنزلية الريفية حوالي 76% من المزارع بينما يشكل النظام شبه التجاري الريفي حوالي 24% ، ومثل الدجاج حوالي ٨٢ % من إجمالي أنواع الدواجن ومثل الحمام والبط ٨.٢%، ٨.٠% علي التوالي، بينما مثل الأوز والأرانب والرومي ١%، ٠.٧%، ٠.١% علي التوالي.

كان متوسط حجم القطيع في النظامين التقليدي و بدون حيازة زراعية ٧٠ طائر وفي النظام شبه التجاري ١٣٢٢ طائر، ووجد أن ٥٨ % إلى ٦٧ % من المربين في النظام التقليدي ونظام التربية بدون حيازة زراعية يستعملون الدواجن ومنتجاتها في الإستهلاك العائلي والباقي يباع في الأسواق المحلية وللتجار، بينما يستعمل نظام شبه تجاري حوالي ١٠ % فقط من منتجات الدواجن للإستهلاك العائلي والباقي ٩٠ % يذهب إلى القنوات التسويقية. تسكن الدواجن في النظامين التقليدي والتربية بدون حيازة زراعية في أعشاش بنسبة ٧٣ % إلى ٧٦ % بينما النظام التجاري تكون التربية في مساكن خاصة بنسبة ٦٦ % أو في ملحق للمسكن بنسبة ٣٤ % ويستعمل عادة العمالة العائلية. الأمراض والتغذية هي أكثر المشاكل التي تواجه مربى الدواجن في هذه الأنظمة بالقطاع الريفي.