

IMPROVEMENT OF THE RHEOLOGICAL AND PHYSICAL PROPERTIES OF NON-FAT YOGURT USING MICROBIAL TRANSGLUTAMINASE, EXOPOLYSACCHARIDES PRODUCING CULTURE AND COMMERCIAL STABILIZERS

Dina A.M. Amer⁽¹⁾, M.Y.A. El-Hawary⁽¹⁾ and M.A. El-Soda⁽²⁾

⁽¹⁾ Department of Food science and Technology, Faculty of Agriculture, Tanta University, Egypt

⁽²⁾ Department of Dairy science and Technology, Faculty of Agriculture, Alexandria University, Egypt

Received: Sep. 24 , 2017

Accepted: Oct. 18 , 2017

ABSTRACT: *The aim of this study was to improve the rheological, physical and organoleptic properties of non-fat yogurt using microbial Transglutaminase (MTGase), Exopolysaccharides producing culture (EPS) and two commercial stabilizers based on pectin and gelatin. Results showed that non-fat yogurt samples produced by textural impotents haven't significant differences with control full-fat yogurt sample in moisture content at fresh, while there were significant differences between these samples and control non-fat yogurt sample. Moisture content significantly decreased in all yogurt samples except in control full and non-fat yogurt samples after 10 days of storage period. Non-fat yogurt EPS sample had the highest percentage moisture content compared with control (full-fat and non-fat) yogurt samples and experimental non-fat yogurt (MTGase, commercial pectin and gelatin) samples throughout storage. Pro/D.M was higher in control non-fat yogurt, non-fat yogurt EPS and non-fat yogurt produced by commercial gelatin compared with non-fat yogurt (MTGase, commercial pectin) and full-fat yogurt samples at fresh. There were no significant differences in Fat /D.M % between non-fat yogurt C2 and non-fat samples containing textural improvements, while there were significant differences between them and control full-fat yogurt C1 at fresh or during storage period. There were no significant differences in acidity among all treatments. Hardness was significantly lower in the EPS non-fat yogurt compared with the other treatments throughout storage period. The non-fat yogurt containing EPS producing culture and non-fat yogurt containing commercial stabilizers had a lower level springiness, Chewiness and gumminess than the full-fat yogurt sample and non-fat yogurt containing MTGase. Adhesiveness significantly increased during storage period for all treatments. There were no significant differences in syneresis percentage among all treatments. Non-fat yogurt produced by Exopolysaccharides producing culture, microbial Transglutaminase and commercial gelatin demonstrated rheological and physical properties were closer to that of control full-fat yogurt.*

Key words: *MTGase, EPS, Pectin, Gelatin, Non-fat yogurt.*

INTRODUCTION

Low fat and non-fat dairy products have gained popularity because of consumer awareness about health concerns related to decreasing the risks connected with obesity and coronary heart diseases (Sandoval *et al.*, 2004). However, the partial or total removal of fat from dairy products decreases the overall quality perceived by the consumers (Folkenberg and Martens, 2006).

It was reported that reduction of fat content in dairy products resulted in lower gel strength and firmness than full fat products, as a consequence of lower number of fat globules embedded in the protein network (Duboc and Mollet, 2001).

To improve textural and functional properties of non-fat yogurt the use of additives has been widely investigated

(Cayot *et al.*, 2007). Fat replacers can be successfully used in the manufacture of reduced fat dairy products such as cheese, ice cream and yogurt (Barrantes *et al.*, 1994). Fat replacer is an ingredient that can be used to provide some or all of the function of fat, yielding fewer calories (Huygbaert *et al.*, 1996), to solve some physical and textural problems originating from low-fat level in the dairy products.

Enzymatic cross-linking of protein by microbial Transglutaminase (MTGase) modifies the techno-functional properties of proteins and is reported as an innovative way of producing novel milk gels (Anema *et al.*, 2005 and Jaros *et al.*, 2010). It catalyzes the acyl-transfer (acyl donor) reaction between the γ -carboxyamid group of peptide or protein-bound glutamyl residues and primary amines (acyl acceptor), is a transferencees (Dickinson, Yamamoto, 1996 and Bonisch *et al.*, 2007). Intermolecular cross-linking of proteins results in high molecular weight polymers which have different functional properties to improve the techno-functional properties of foods (Gauche *et al.*, 2009, Lorenzen, *et al.*, 2002 and Faergemand, & Qvist, 1999). Milk proteins, especially caseins, are good substrates for cross-linking with MTGase (Faergemand, and Qvist, 1997 and Lorenzen, Schlimme, 1998). It was reported that cross-linking of the proteins in milk improved gels firmness and reduced serum separation of acid-induced milk gel, mainly set-type yogurts (Lorenzen, *et al.*, 2002, Lorenzen, & Schlimme, 1998 and Şanlı, *et al.*, 2011). (Özer *et al.*, 2007) also expressed that the MTGase added into milk may be an alternative method instead of addition of extra protein and stabilizer in non-fat yogurt.

Exopolysaccharides (EPS)-producing culture are used to modify texture of yogurt (Hassan, Frank, Schmidt, & Shalabi, 1996a; Hassan & Frank, 1997; Perry, McMahon, & Oberg, 1997, 1998). Yogurt made with EPS-producing cultures is less susceptible to syneresis, more viscous, and had more

water holding capacity than that made with EPS-non-producing cultures (Hassan, Frank, Schmidt, & Shalabi, 1996a, b). The use of EPS-producing cultures produces soft cheese curd (Hassan & Frank, 1997) and higher water retention in Mozzarella cheese (Perry *et al.*, 1997, 1998 and Hassan *et al.*, 2003).

Results obtained by (Fizman *et al.*, 1999) demonstrated the suitability of using of gelatin to improve the quality of milk products. Addition of gelatin to the milk during preparation of the yogurt changed the microstructure of the product by the formation of flat sheets or surfaces which interacted with the casein matrix, enclosing granules of casein in several zones. Gelatin seemed to connect the granules and chains of milk proteins, and consequently create a continuous, fairly homogeneous double network structure with no free ends. This more interconnected network would retain the aqueous phase more efficiently, reducing the drainage of liquid.

Pectin is a component of the primary cell wall of higher plants. It consists of polymers of D-a-(1→4) anhydro-galacturonic acid (at least 65%) with a variable numbers of methyl ester groups (Wosiacki 1977). Pectins are divided into two categories depending on the degree of esterification: pectins with a degree of methylation >50% are described as high-methoxyl pectins (HMP) whereas pectins with a degree of methylation <50% are known as low-methoxyl pectins (LMP) (Lootens *et al.* 2003; Tsoga *et al.* 2004). The degree of esterification can strongly influence the gel network of pectin (Sato *et al.* 2004). High-methoxyl pectins can be obtained from citrus peels and apple pomace and is used as stabilizer, especially in acidic products, for example yogurt and acidic milk beverages. LMP can be obtained from citrus peels and is used as a gelling or thickening agent in acidic or less acidic products such as milk desserts (Williams and Phillips 2000 and Leroux *et al.* 2003).

Improvement of the rheological and physical properties of non-fat

The objectives of this study was to investigate the possibility of making a good quality non-fat yogurt by using textural improvements: Microbial Transglutaminase (MTGase), Exopolysaccharides producing culture (EPS), commercial pectin and commercial gelatin and to monitor the changed yogurt quality during storage period.

MATERIALS AND METHODS

Materials

Fresh full-fat and skim cow's milk was obtained from the farm of Faculty of Agriculture, Alexandria University. Skim milk powder (SMP) was obtained from Rucker Company, Germany.

Starter cultures, Freeze-dried lactic culture for direct vat set (DVS). (Express 0.2, thermophilic yogurt culture Yo-Flex Express), consisting of *Lactobacillus delbrueckii* ssp. *bulgaricus* and *Streptococcus thermophilus* was obtained from Chr. Hansen Lab., Denmark

Textural improvements:

Microbial Transglutaminase (MTGase): The Ca⁺²-independent microbial Transglutaminase from *Streptococcus mobaraense* (Activa® YG) was supplied from Ajinomoto Foods Europe (Hamburg, Germany) with concentration 0.02% (w/w), Exopolysaccharides producing culture (EPS): The strain *Lactobacillus delbrueckii* ssp. *Lactis* 127ST used in this study was isolated from traditional fermented milk in the Laboratory of the Biochemistry of Dairy microorganisms, Alexandria University (Egypt) with concentration 1% (v/v).

The commercial stabilizer,

Lacta 534 (based on gelatin E 441, Modified starch E 1422, Agar E 406 and Mono-and diglycerides E471) with concentration of 0.4% (w/w).

Lacta 533 (based on pectin E 440, Modified starch E 1422, and Mono-and diglycerides E471) with concentration of

0.4% (w/w) were obtained from MIFAD Company, Alexandria, Egypt.

Chemicals used for the detecting of the fat, protein, etc., were obtained from El-Gomhoria Company for chemicals and glasses.

Yogurt Manufacture:

Fresh full fat and skim cow's milk was fortified with 3% and 3.5% skim milk powder respectively and the stabilizers were added individually. The control 1 was full fat milk (3% fat, 13.2% TS), while the control 2 was non-fat milk (0.4% fat, 10.21% TS) without adding stabilizers adding. Microbial Transglutaminase was added at 0.02%, while Exopolysaccharides producing culture (EPS) was add at 1% v/v. Yogurt was manufactured according to the protocol proposed by (Tamime and Robinson, 1999).

Individual milk samples were heat treated at 90°C for 10 min, cooled to 42°C and inoculated with yogurt culture at the rate recommended by suppliers, incubation at 42°C till the pH reaching 4.9. This was followed by fast cooling to 7°C keeping the product at the same temperature overnight to represent fresh samples yogurt that were stored at 7±1°C for 10 days.

Commercial stabilizers based for gelatin and pectin were added before heat treatment, but Microbial Transglutaminase (MTGase) and Exopolysaccharides producing culture (EPS) were added with the addition of yogurt starter culture after heat treatment and cooling.

Methods of analysis:

Sampling: Yogurt samples were taken from the (fresh, 5 days and 10 days).

Chemical analysis:

All samples were analyzed for fat, total protein and dry matter according to A.O.A.C. (2000).

All samples were analyzed for acidity and pH according to Ling (1963).

Syneresis:

The syneresis % of yogurt samples was determined during storage period using centrifugation method as described by (Keogh and O’Kennedy, 1998). Yogurt (30–40 g) was centrifuged at 1100 r.p.m. for 10 min at 5°C. The clear supernatant was poured off, weighed and recorded as syneresis (%). Duplicate measures were performed for each sample. Syneresis% = (weight of supernatant /weight of yogurt sample)× 100.

Textural analysis:

Textural properties of yogurt were evaluated using a texture analyzer (FTC TMS-Pro), USA). Yogurt samples were evaluated in their cups). A two-bite penetration test was performed using the Texture Analyzer with the TA 25 mm diameter cylinder probe for yogurt and operated at a crosshead speed 50 mm/sec. Hardness, cohesiveness, springiness and chewiness were evaluated in triplicate as described by Szczesniak *et al* (1963) and Bourne, (1978).

Sensory evaluation:

Sensory evaluation of yogurt samples was conducted by panelists. The panel lists were asked to evaluate the color (10), appearance (10), texture (35) and flavor (45) and overall acceptability when fresh, after 7 days and 10 days of storage according to Farag *et al.*, (2007).

Scanning electron microscopy:

Samples of yogurt were fixed in 2.5% glutaraldehyde in cacodylate buffer (pH 7.2) for at least 1 h. After rinsing three times in cacodylate buffer, samples were postfixed in a 1% buffered osmium tetroxide for one hour. The fixed samples were dehydrated using a graded alcohol series (20%, 40%, 60%, 70%, and 90%) finishing with three changes of 100% alcohol then critical point dried from liquid Co₂. At least three dried samples of each yogurt were fractured, mounted on aluminum stubs, and coated

with gold in a K550X sputter coater (England) as described by (Puvanenthiran *et al.*, 2002). At least two images of typical structures at 1000×magnification were recorded using a Scanning Electron Microscope (JEOL JSM-5300 Scanning Microscope) by electron microscope Unite, Faculty of Science, Alexandria University.

Statistical analysis:

Data were analyzed as means, stander deviation and Duncan’s test and subject to IBM SPSS version 16 Statistics for analysis of variance (ANOVA).

RESULTS AND DISCUSSION

1. Chemical analysis

Data in Table (1) showed moisture content %, fat content/dry matter (F/DM) and total protein content/dry matter (Pro/DM). The obtained results showed that there were no significant differences between control full-fat yogurt C1 and non-fat yogurt containing textural improvements at fresh in moisture content, while there were significant differences between control non-fat yogurt C2 and all experimental treatments. During storage period moisture % significantly decreased in all samples except in C1 and C2. Also results showed that control full-fat yogurt had lower moisture% compared with other samples, while control non-fat yogurt had highest moisture%. Using different textural improvement had increase moisture% than control non-fat yogurt. Non-fat yogurt produced by EPS had the highest moisture content compared with C1, C2 and all experimental samples after 10 days. These results are in agreement with Jimenez-Guzman *et al.* (2009), Trancoso-Reyes *et al.* (2014) who mentioned that EPS producing cultures have excellent water binding properties and moisture retention, which improve the quality of low fat dairy products. Also results showed non-fat yogurt made with lacta 533 based on pectin and lacta 534 based on gelatin reduce moisture% when

Table (1): Chemical analysis of non-fat yogurt made with textural improvements at fresh and during the storage periods

Treatments	Chemical properties											
	Moisture content (%)			Fat/D.M			Pro /D.M					
	Fresh	5 days	10 days	Fresh	5 days	10 days	Fresh	5 days	10 days			
C1	84.83±0.30 ^{As}	84.72±0.28 ^{As}	83.93±1.00 ^{As}	19.77±0.40 ^{Bs}	19.63±0.37 ^{Bs}	19.27±0.96 ^{Bs}	21.25±0.10 ^{As}	21.34±0.40 ^{As}	20.26±1.02 ^{As}			
C2	86.32±1.12 ^{Bs}	85.51±0.78 ^{As}	84.51±0.35 ^{Bs}	0.74±0.06 ^{As}	0.69±0.04 ^{As}	0.66±0.01 ^{As}	24.70±2.16 ^{Bs}	23.59±0.91 ^{Bs}	22.46±0.23 ^{Bs}			
MT.Gase	85.08±0.57 ^{As}	84.80±0.06 ^{As}	84.21±0.21 ^{As}	0.67±0.00 ^{As}	0.66±0.03 ^{As}	0.63±0.00 ^{As}	22.65±0.78 ^{As}	22.70±0.76 ^{As}	21.66±0.03 ^{Bs}			
EPS	86.04±0.18 ^{As}	85.84±0.01 ^{Bs}	85.55±0.13 ^{Bs}	0.72±0.01 ^{As}	0.71±0.01 ^{As}	0.66±0.04 ^{As}	24.38±0.54 ^{Bs}	24.16±0.43 ^{As}	24.03±0.43 ^{As}			
Pectin	85.48 ±0.27 ^{As}	85.17±0.14 ^{As}	84.76±0.07 ^{As}	0.69±0.01 ^{As}	0.67±0.00 ^{As}	0.65±0.02 ^{As}	23.28±0.42 ^{As}	23.01 ± 0.28 ^{Bs}	22.83 ± 0.17 ^{Bs}			
Gelatin	85.60±0.17 ^{As}	85.34±0.78 ^{As}	84.90±0.50 ^{As}	0.70±0.01 ^{As}	0.68±0.00 ^{As}	0.66±0.00 ^{As}	23.47±0.26 ^{Bs}	23.22± 0.12 ^{Bs}	22.90 ± 0.14 ^{Bs}			

Data are means ±S.D. ^{As}C1: control yogurt made from full-fat cow's milk. ^{As}C2: control yogurt made from non-fat cow's milk. (MT.Gase): yogurt made from non-fat cow's milk + microbial Transglutaminase (EPS): yogurt made from non-fat cow's milk + Exopolysaccharides producing culture. Pectin: yogurt made from non-fat cow's milk + Lactia 533. Gelatin: yogurt made from non-fat cow's milk + Lactia 534. (A,B) Means at the column with different superscripts are different (P<0.05) by Duncan's tests multiple comparison tests. (a,b) Means at the row with different superscripts are different (P<0.05) by Duncan's multiple comparison tests.

compared to control non-fat yogurt C2 at fresh. These Results are in agreement with Yazici and Akgun (2004) who found that some protein based fat replacers in low fat yogurt led to an increase in dry matter. Total solids contents were found to be higher in yogurts with stabilizers, compared with control yogurts reflecting higher total solids content in treated yogurts due to addition of stabilizers (Mehanna, *et al.*, 2013) Changes in this parameters may affect certain other physico-chemical properties such as syneresis, water holding capacity (Wu, *et al.*, 2001). There were significant difference in Pro/D.M between controls C1, C2 and samples containing textural improvements at fresh and after 10 days of storage period. Pro/ D.M was higher in control non-fat C2, non-fat yogurt produced by EPS and non-fat yogurt produced by commercial gelatin compared with control full-fat yogurt C1 and other experimental samples at fresh. After 10 days non-fat yogurt EPS had the highest Pro/D.M compared with all experimental treatments. Control full-fat yogurt had lowest Pro/D.M, compared with all experimental treatments after 10 days. There were no significant differences in Fat /D.M% between non-fat yogurt C2 and non-fat samples containing textural improvements, while there were significant differences between them and control full-fat yogurt C1 at fresh or during storage period.

2- Acidity% and pH value

Fresh yogurt controls (full-fat and non-fat) had similar acidity levels. While, the acidity was at high level in fresh treatments with commercial (pectin, gelatin) and EPS but was low in treatment with MTGase (Table 2). The acidity wasn't significantly increased during storage in all samples except in yogurt produced by MTGase had increased at the end of storage period. There were no significant differences in pH value between all samples. While the pH value was lower in non-fat yogurt that contains EPS. There was no significant variation ($P > 0.05$) in pH value of non-fat yogurt that containing

textural improvements throughout storage period. During storage period, all non-fat yogurt samples showed gradual decrease in pH until the end of storage period. These findings are in good agreement with (Lorenzen *et al.*, 2002). Also these results are in agreement with those obtained by (Anis *et al.*, 1989) and Mehanna (1989); Abd El-salam *et al.*, (1996) who studied the use of textural improvement to improve some properties of skim cow's milk yogurt and showed that Titratable acidity of all yogurt treatments increased significantly as storage period was progressed.

3- Texture analysis

Rheological properties for foods, such as fermented dairy products, are important in the design of flow processes, quality control, storage and processing and in predicting the texture of foods (Shaker *et al.*, 2000). Table (3) shows the texture parameters (Hardness, Adhesiveness, Cohesiveness, Springiness, Chewiness and Gumminess) of the different treatment of yogurt over storage. The Hardness of yogurt is directly dependent on its total solids and specifically protein content and the type of proteins. Higher protein content would cause a higher degree of cross-linkage of the gel network, resulting in a much denser and more rigid gel structure (Tamime 2006). Hardness was significantly lower in the EPS non-fat yogurt compared with the other treatments throughout storage. This coincides with the lower intact casein content of the EPS yogurt as a consequence of its higher levels of moisture content. At the same time, the presence of the EPS within the yogurt matrix may also have helped to reduce the hardness by filling voids in the matrix and by interacting with the proteins. The non-fat yogurt containing EPS producing culture and non-fat yogurt containing stabilizers had a lower level springiness than the full fat yogurt and non-fat yogurt containing MTGase, an effect that may be attributed to the higher content of intact casein and higher moisture content. Many researchers

Improvement of the rheological and physical properties of non-fat

reported that the Exopolysaccharides could improve the texture of yogurt, because Exopolysaccharides produced by LAB interacts with the free water in the gel-like structure (Guzel *et al.*, 2005, De Vuyst and Degeest 1999, Hassan *et al.*, 2003). Chewiness and gumminess were significantly lower in non-fat yogurt containing EPS producing culture and non-fat yogurt containing stabilizers in compared with full-fat yogurt and non-fat yogurt

containing MTGase. Yogurt hardness correlated positively with Chewiness and gumminess. Adhesiveness significantly increased during storage for all treatments. No significant differences were detected in cohesiveness, the increase in the moisture content in the non-fat yogurt containing EPS producing culture and non-fat yogurt contain stabilizers had little effect on the cohesiveness of the yogurt.

Table (2): Acidity (%) and pH value of non-fat yogurt made with textural improvements at fresh and during the storage periods

Acidity (%)			
Treatments	Storage period		
	Fresh	5 days	10 days
C1	0.87±0.01 ^{Aa}	0.89±0.01 ^{Aa}	0.89±0.01 ^{Aa}
C2	0.84±0.06 ^{Aa}	0.89±0.10 ^{Aa}	1.01±0.01 ^{Aa}
MTGase	0.74±0.03 ^{Aa}	0.81±0.01 ^{Aa}	0.99±0.04 ^{Ab}
EPS	0.95±0.18 ^{Aa}	1.00±0.25 ^{Aa}	1.10±0.25 ^{Aa}
Pectin	1.04±0.14 ^{Aa}	0.87±0.78 ^{Aa}	0.84±0.36 ^{Aa}
Gelatin	1.01±0.42 ^{Aa}	0.84±0.42 ^{Aa}	0.91±0.07 ^{Aa}
pH value			
Treatments	Storage period		
	Fresh	5 days	10 days
C1	4.71±0.09 ^{Aa}	4.64±0.35 ^{Aa}	4.36±0.42 ^{Aa}
C2	4.79±0.78 ^{Aa}	4.73±0.07 ^{Aa}	4.53±0.15 ^{Aa}
MTGase	4.74±0.22 ^{Aa}	4.62±0.13 ^{Aa}	4.48±0.27 ^{Aa}
EPS	4.59±0.52 ^{Aa}	4.45±0.36 ^{Aa}	4.40±0.55 ^{Aa}
Pectin	4.66 ± 0.13 ^{Aa}	4.63 ± 0.14 ^{Aa}	4.50 ± 0.25 ^{Aa}
Gelatin	4.56 ± 0.23 ^{Aa}	4.73 ± 0.14 ^{Aa}	4.70 ± 0.16 ^{Aa}

Data are means ±S.D * C1: control yogurt made from full-fat cow's milk * C2: control yogurt made from non-fat cow's milk (MTGase): yogurt made from non-fat cow's milk + microbial Transglutaminase (EPS): yogurt made from non-fat cow's milk + Exopolysaccharides producing culture Pectin: yogurt made from non-fat cow's milk + Lacta 533 Gelatin: yogurt made from non-fat cow's milk + Lacta 534 (A.B) Means at the column with different superscripts are different (P<0.05) by Duncan's tests multiple comparison tests. (a.b) Means at the row with different superscripts are different (P<0.05) by Duncan's multiple comparison tests

Table (3): Texture analysis of non-fat yogurt made with textural improvements at fresh and during the storage periods

Texture properties	Storage period	Treatments					
		C1	C2	MTGase	EPS	Pectin	Gelatin
Hardness (g)	Fresh	134	112.50	154.50	103.50	106	113
	5 days	138	121.00	200.00	101.50	110.50	112.50
	10 days	140	123.50	174.00	99.50	110.50	110.50
Adhesiveness (g.mm)	Fresh	98.30	95.73	154.50	103.70	105.36	104
	5 days	109.90	94.48	94.47	92.33	111.28	111.11
	10 days	119.68	97.18	101.85	92.22	96.42	96.86
Adhesive force(g)	Fresh	-31.50	-23.50	-23.50	-20.50	-23	-23
	5 days	-29.50	-29.00	-30.50	-25.50	27.50	27.50
	10 days	-27.00	-28.00	-30.50	-24.50	27	27
Cohesiveness	Fresh	0.42	0.40	0.36	0.40	0.40	0.40
	5 days	0.41	0.38	0.32	0.36	0.40	0.43
	10 days	0.30	0.38	0.37	0.41	0.38	0.38
Springiness (m.m)	Fresh	6.37	5.36	5.47	5.70	5.60	5.59
	5 days	6.91	5.70	6.03	5.62	6.02	6.15
	10 days	7.15	5.67	5.92	5.72	5.95	5.82
Chewiness (mJ)	Fresh	350.40	2.40.90	300.50	235.01	236.86	252.63
	5 days	389.68	262.16	386.31	222.08	279.76	269.93
	10 days	379.56	262.38	381.16	218.62	249.46	241.04
Gumminess (N)	Fresh	55.50	45	55	41	42.50	45.25
	5 days	58.80	46	64	39.50	44	48
	10 days	53.50	48.50	64.50	38.00	42	41.50

Data are means ±S.D *C1: control yogurt made from full-fat cow's milk *C2: control yogurt made from non-fat cow's milk (MTGase): yogurt made from non-fat cow's milk + microbial Transglutaminase (EPS): yogurt made from non-fat cow's milk + Exopolysaccharides producing culture Pectin: yogurt made from non-fat cow's milk + Lacta 533 Gelatin: yogurt made from non-fat cow's milk + Lacta 534

4. Syneresis values:

Serum separation occurs in fermented milk products due to the aggregation and sedimentation of casein particles during storage. Textural improvements were found to be necessary to prevent serum separation in fermented milk (Lucey *et al.*, 1999;

Towler, 1984). Syneresis an undesirable property in yogurt products (Wu, *et al.*, 2001).

Values of syneresis was 22.55% in fresh non-fat yogurt, it was increased to 27.45 % after 10 days throughout storage (Table 4). There were differences in syneresis value

Improvement of the rheological and physical properties of non-fat

among all treatments samples at fresh or during storage period but these differences were not significantly. The Syneresis was lower in full-fat treatment when compared to non-fat counterpart. High fat content in yogurt has been associated with lower syneresis values (Keogh and O’Kennedy, 1998 and Isanga and Zhang, 2009). Common reasons for the occurrence of syneresis are low dry matter and protein contents. Enrichment of dry matter content is common way of avoiding syneresis (Amatayakul *et al.* 2006). Results reported that at fresh non-fat yogurt MTGase sample was lower in syneresis % than controls (full and non-fat) yogurt and non-fat yogurt with stabilizers (commercial pectin, gelatin), although non-fat yogurt EPS sample was higher. This may due to the cross-linking of protein chains by microbial Transglutaminase (MTGase) can stabilize the three dimensional network of yogurt gel and decreased syneresis% as a result of decrease whey expulsion as a result of decrease in gel porosity (Faergamand *et al.* 1999; özer *et al.*, 2007; Sanli *et al.*, 2011). Also these results were similar to (Folkenberg *et al.* 2003). Who reported that

syneresis of yogurt made with EPS-producing cultures was higher than those made with non- EPS- producing cultures due to their water-binding ability and texture promoting properties (Amatayakul *et al.* 2006).

5. Organoleptic properties:

The results of the evaluation of different non-fat yogurts through 10 days of the storage period are shown in Table (5). Results indicated that there were no significant among all samples in color and appearance. The texture of yogurt made with commercial gelatin was firmer than (C1) and (C2). Addition of stabilizers, improved the Body & Texture of the non-fat yogurt as compared with control non-fat yogurt to be similar with full fat control, non-fat yogurt produced by commercial gelatin had highest texture compared with all samples. Also results showed that flavor of all samples were improved by textural improvements as control full-fat yogurt compared with control non-fat yogurt. Color and appearance of the yogurt samples were scored most highly for treatment EPS and gelatin compared with MTGase and pectin.

Table (4): Syneresis (%) of non-fat yogurt made with of textural improvements at fresh and during the storage periods

Samples	Storage periods		
	Fresh	5 days	10 days
C1	17.25±2.36 ^{Aa}	20.08±1.05 ^{Aa}	18.91±4.55 ^{Aa}
C2	22.25±9.97 ^{Aa}	25.80±6.48 ^{Aa}	27.45±7.84 ^{Aa}
MTGase	14.23±0.64 ^{4a}	23.32±3.56 ^{Aa}	24.50±2.82 ^{Ab}
EPS	25.77±2.26 ^{Aa}	24.60±6.07 ^{Aa}	24.79±2.24 ^{Aa}
Pectin	19.26±7.23 ^{Aa}	23.37±3.91 ^{Aa}	23.70±1.13 ^{Aa}
Gelatin	16.79±7.96 ^{Aa}	22.72±1.01 ^{Aa}	22.60±0.14 ^{Aa}

Data are means ±S.D *C1: control yogurt made from full-fat cow's milk *C2: control yogurt made from non-fat cow's milk (MTGase): yogurt made from non-fat cow's milk + microbial Transglutaminase (EPS): yogurt made from non-fat cow's milk + Exopolysaccharides producing culture Pectin: yogurt made from non-fat cow's milk + Lacta 533 Gelatin: yogurt made from non-fat cow's milk + Lacta 534 (A,B) Means at the column with different superscripts are different (P<0.05) by Duncan's tests multiple comparison tests. (a,b) Means at the row with different superscripts are different (P<0.05) by Duncan's multiple comparison tests

Table (5): Organoleptic properties of non-fat yogurt made with different textural improvements at fresh and during storage periods

Treatments	Storage period	Appearance (10)	Color (10)	Texture (35)	Flavor (45)
C1	Fresh	9.50±0.71 ^{Aa}	9.00±0.00 ^{Aa}	32.50±0.71 ^{ABa}	43.50±0.71 ^{1Ba}
	5 days	8.75±0.35 ^{Aa}	8.75±0.35 ^{Aa}	33.00±0.00 ^{ABa}	42.00±0.00 ^{Aa}
	10 days	9.50±0.71 ^{Aa}	9.50±0.71 ^{Aa}	32.50±0.71 ^{Aa}	42.50±0.71 ^{Aa}
C2	Fresh	9.25±0.35 ^{Aa}	9.00±0.00 ^{Aa}	31.50±0.71 ^{Aa}	39.50±0.71 ^{Aa}
	5 days	9.25±0.35 ^{Aa}	9.75±0.35 ^{Ba}	33.00±0.00 ^{ABb}	42.50±0.71 ^{Aa}
	10 days	9.50±0.71 ^{Aa}	9.50±0.71 ^{Aa}	33.00±0.00 ^{Ab}	40.00±1.41 ^{Aa}
MTGase	Fresh	9.00±0.00 ^{Aa}	8.50±0.71 ^{Aa}	33.50±0.71 ^{BCa}	41.50±0.71 ^{ABa}
	5 days	9.50±0.71 ^{Aa}	9.75±0.35 ^{Aa}	34.50±0.71 ^{BCa}	43.50±0.71 ^{Ab}
	10 days	9.00±0.00 ^{Aa}	9.00±0.00 ^{Aa}	33.00±0.00 ^{Aa}	41.75±0.35 ^{Aab}
EPS	Fresh	9.50±0.71 ^{Aa}	9.50±0.71 ^{Aa}	33.50±0.71 ^{BCa}	41.50±2.12 ^{ABa}
	5 days	9.50±0.71 ^{Aa}	9.00±0.00 ^{Aa}	34.00±0.00 ^{BCa}	42.00±0.00 ^{Aa}
	10 days	10.00±0.00 ^{Aa}	10.00±0.00 ^{Aa}	33.50±0.71 ^{Aa}	42.00±0.00 ^{Aa}
Pectin	Fresh	9.00±0.00 ^{Aa}	9.00±1.41 ^{Aa}	32.50±0.71 ^{Aa}	41.00±1.41 ^{ABa}
	5 days	8.75±0.35 ^{Aa}	9.00±0.00 ^{Aa}	34.00±0.00 ^{Cb}	42.50±0.71 ^{Aa}
	10 days	9.00±0.00 ^{Aa}	9.00±0.00 ^{Aa}	32.25±0.35 ^{Aa}	42.00±0.00 ^{Aa}
Gelatin	Fresh	10.00±0.00 ^{Aa}	10.00±0.00 ^{Aa}	34.00±0.00 ^{Ca}	42.50±0.71 ^{ABa}
	5 days	9.00±0.00 ^{Aa}	9.25±0.35 ^{ABa}	32.50±0.71 ^{Aa}	42.50±2.12 ^{Aa}
	10 days	9.00±0.00 ^{Aa}	9.00±0.00 ^{Aa}	32.25±1.06 ^{Aa}	42.50±2.12 ^{Aa}

Data are means ±S.D *C1: control yogurt made from full-fat cow's milk *C2: control yogurt made from non-fat cow's milk (MTGase): yogurt made from non-fat cow's milk + microbial Transglutaminase (EPS): yogurt made from non-fat cow's milk + Exopolysaccharides producing culture Pectin: yogurt made from non-fat cow's milk + Lacta 533 Gelatin: yogurt made from non-fat cow's milk + Lacta 534

(A.B) Means at the column with different superscripts are different (P<0.05) by Duncan's tests multiple comparison tests.

(a.b) Means at the row with different superscripts are different (P<0.05) by Duncan's multiple comparison tests

6. Microstructure

It is well recognized that the structure of foods affects various properties including texture, functionality and appearance. Microstructure has a major impact on the texture and other physical properties of acid gels such as yogurt (Lee and Lucey 2010). The SEM images of the experimental materials can be seen in Fig (1, 2, 3, 4, 5 and

6). The full-fat yogurt had large pore sizes with smaller strands of protein in the microstructure compared to other samples. Treated yogurts (MTGase, pectin, gelatin and EPS) had more systematically and smoothly distributed casein with a bit coarse structure as well as less porosity in casein network. This might be attributed to hydrocolloids and emulsion stability

Improvement of the rheological and physical properties of non-fat

catalyzed cross-link formation between milk proteins as reported by (Lorenzen *et al.*, (2002). Control of non-fat yogurt contained fewer and larger pores than yogurt made with the EPS-producing culture. It was observed that the protein matrix of yogurt made using EPS- producing starter had a

more homogenous compared to control C1 and C2. However MTGase sample, it was possible to obtain high gel strength owing to the reduced mesh size of the protein network. This is in agreement with the result of (Lorenzen *et al.* 2002 and Asaanli *et al.* 2011).




Fig. 1. Scanning electron microscopy photographs (×1000) of control full-fat fresh yogurt.




Fig. 1.1. Scanning electron microscopy photographs (×1000) of control full-fat yogurt storage at 10 days.




Fig. 2. Scanning electron microscopy photographs (×1000) of control non-fat fresh yogurt.




Fig. 2.2. Scanning electron microscopy photographs (×1000) of control non-fat yogurt storage at 10 days.




Fig. 3. Scanning electron microscopy photographs (×1000) of control non-fat fresh yogurt with MTGase




Fig. 3.3. Scanning electron microscopy photographs (×1000) of control non-fat yogurt with MTGase storage at 10 days




Fig. 4. Scanning electron microscopy photographs (×1000) of control non-fat fresh yogurt with EPS




Fig. 4.4. Scanning electron microscopy photographs (×1000) of control non-fat yogurt with EPS storage at 10 days.




Fig. 5. Scanning electron microscopy photographs (×1000) of control non-fat fresh yogurt with pectin




Fig. 5.5 Scanning electron microscopy photographs (×1000) of control non-fat yogurt storage at 10 days with pectin




Fig. 6. Scanning electron microscopy photographs (×1000) of control non-fat fresh yogurt with gelatin




Fig. 6.6. Scanning electron microscopy photographs (×1000) of control non-fat yogurt storage at 10 days with gelatin

In comparing between fresh and 10 days old yogurt, we concluded that non-fat control underwent more extensive water

redistribution than non-fat yogurt made with EPS. The electron microscopy micrographs correlated well with the texture analysis

data. Disturbance of protein-protein interactions and the porosity of the background protein network in the non-fat yogurt seemed to be responsible for the improving the texture properties in these types of yogurt. The microstructure of yogurt has been well studied, and some data have been published on the mechanisms of the acid induction of gels in milk by *Streptococcus thermophilus* and *Lactobacillus delbrueckii subsp. bulgaricus* at 30–45 °C. However, the casein micelles are composed of different protein fractions, and are associated with one another via Ca-phosphate bridges. During the fermentation of milk, the micelle or colloidal Ca²⁺ content (and possibly to a lesser extent magnesium and citrate) increases in the serum as the pH is lowered due to the solubilization of micelle Ca-phosphate (Awad, 2011). Adding textural improvement can affect formation of casein gel structure by acting as filler. Therefore, the effect of textural improvement on protein matrix and structure formation depends on their composition, molecular properties, concentration and interactions with the protein. The appearance of casein micelles were less defined. These differences were probably due to the interactions between casein micelles and stabilizers through mainly hydrophobic interaction leading to the formation of casein stabilizers complexes (Wang *et al.*, 2012). Tamime and Robinson, (1999) reported that, in typical scanning electron micrographs of yogurt with textural improvement, a casein matrix is visible with various forms and sizes of compact area. Scanning electron microscopy (SEM) photographs of fresh skim cow milk yogurt with textural improvement which increases the cohesion flat casein compared with a control sample. The presence of EPS channels in the serum will confer a more polymer-like rheological behavior to the continuous phase, thus resulting in increased consistency index, more deviation from Newtonian behavior, and increased viscosity in the yogurt. The shear-induced microstructure in yogurt

made with EPS+ was shown to consist of compartmentalized protein aggregates between channels containing EPS. These EPS containing channels cause yogurt made with EPS+ hindered syneresis and buildup of structure after stirring

Conclusions

Obtained results suggest possibility of making a good quality low-calorie non-fat yogurt with the use EPS which were similar in quality characteristics to control made with full-fat milk. Also, results indicated that microbial Transglutaminase MTGase mediated cross-linking of milk proteins may be useful method for the production of non-fat yogurt without adversely affecting the affecting the physical properties of the product. It has been shown that cross-linking resulted in improved gel firmness and reduced serum separation in non-fat yogurt. Results also indicated that suitability of using commercial gelatin to improve characterizes of non-fat yogurt. These treatments produced the best quality of non-fat yogurt, besides these trials had low calories and less cost when compared to full fat yogurt and matched the demands of a wide sector of customers. Generally reduction of total solid causes some defects in texture of yogurt, but textural improvement can surmount some of these defects.

Acknowledgements

The authors would like to thank Mrs. Nour El-Hoda Ahmed for help in texture analysis and moisture determination of the yogurt samples. Thanks are also due to Mrs. Gehan Osman for her help in microbiological analysis.

REFERENCES

- Abd El-Salam, M.H., H.M. EL-Etriby and N. M. Shahein (1996). Influence of some stabilizers on some chemical and physical properties of yogurt. Egyptian J. Dairy Sci. 24:25-36.

- Amatayakul, T., F. Sherkat and N. P. Shah (2006). Physical characteristics of set yogurt made with altered casein to protein ratios and EPS-producing starter cultures at 9 and 14% total solids. *Food Hydrocolloids* 20: 314–324.
- Anema, S.G., S. Lauber, S.K. Lee, T. Henle and H. Klostermeyer (2005). Rheological properties of acid gels prepared from pressure and transglutaminase-treated skim milk. *Food Hydrocolloids*, 19: 879-887.
- Anis, S.M.K., E.S. Girgis, S.N. Amer and P.N.M. Salama (1989). Effect of using some stabilizers of yogurt manufacture. *Annals of agric. Sci.*, Moshtohor, 27: 397-406.
- A.O.A.C. (2000). *Official Methods of Analysis*. 14th Ed. Association of Official Analytical Chemists. Washington, DC.
- Awad, S. (2011). Texture and microstructure: in *Practical Food and Research* pp. 361-391 (ed. Rui M. S. Cru), Nova Science publishers, Inc.
- Bade, I. S., T. Bernardi and P. Michaud (2011). New perspectives for Lactobacillixopolysaccharides. *Biotechnology Advances* 29: 54–66.
- Barrantes, E., A.M. Sword, G. Davies and M.N. Barclay (1994a). Production of low-calorie yogurt using skim milk powder and fat-substitutes: Rheological properties. *Milchwissenschaft*, 49: 263-265.
- Bonisch, M.P., M. Huss, K. Weigl and U. Kulozik (2007). Transglutaminase cross-linking of milk proteins and impact on yogurt gel properties. *Int. Dairy J.* 17: 1360-1371.
- Bourne, M.C. (1978). Texture profile analysis. *Food Technology*. July. 62-72.
- Cayot, P. S., F. Chenker, G. Houze, C. Sulmont-Rosse and B. Colas (2007). Creaminess in relation to consistency and particle size in stirred fat-free yogurt. *Int. Dairy J.* 18: 303-311.
- De Vuyst, L. and B. Degeest (1999). Heteropolysaccharides from lactic acid bacteria. *FEMS Microbiology Reviews*, 23:153–177.
- Dickinson, E. and Y. Yamamoto (1996). Rheology of milk protein gels and protein-stabilized emulsion gels cross-linked with transglutaminase. *J. Agric. Food Chem.*, 44: 1371-1377.
- Duboc, P. and B. Mollet (2001). Application of Exopolysaccharides in the dairy industry. *Int. Dairy J.* 11: 759-768.
- Faergemand, M. and K.B. Qvist (1997). Transglutaminase: Effect on rheological properties, microstructure and permeability of set style acid skim milk gel. *Food Hydrocolloids*, 11:287-292.
- Faergemand, M., M. Sorensen, U. Jorgensen, G. Budolfson and K. B. Qvist (1999). Transglutaminase: effect on instrumental and sensory texture of set style yogurt. *Milchwissenschaft*, 54: 563–566.
- Farag-Seham, I., A. H.El-sonbaty and S.A. Hussein, A.F. Farrag and A.M. Shahine (2007). Effect of substituting added skim milk powder (SMP) with wheat germ (WG) on the quality of goat's milk yogurt and fermented camels milk drink. *Proc. 10th Egypt. Conf. Dairy Sci. and technol.*, 315-336.
- Fizman, S.M., M.A. Luch and A. Salvador (1999). Effect of addition of gelatin on microstructure of acidic milk gels and yogurt and on their rheological properties. *Int. Dairy J.* 9: 895-901.
- Florea, T. and G.M. Costin (2005). Exopolizaharide produse de culturile starter utilizate la fabricarea produselor lactate fermentate. In: Costin, G.M. Ed. *Academica*, Galati, Italy. p. 421-510.
- Folkenberg, D.M. and M. Martens (2003). Sensory properties of low-fat yogurts. Part A: Effect of fat content, fermentation culture and addition of non-fat dry milk on the sensory properties of plain yogurts. *Milchwissenschaft*, 58: 48-51.
- Folkenberg, D.M., P. Dejmek, A. Skliver, H. S. Guldager and R. Ipsen (2006). Sensory and rheological screening of exopolysaccharide producing strains of

Improvement of the rheological and physical properties of non-fat

- bacterial yogurt cultures. *Int. Dairy J.* 16: 111–118.
- Gauche, C., T. Tomazi, PLM. Barreto, P.J. Ogliari and MT. Bordignon-Luiz (2009). Physical properties of yogurt manufactured with milk whey and transglutaminase. *LWT-Food Sci. Technol*, 42: 239-243.
- Guzel-Seydim, Z. B., E. Sezgin and A. C. Seydin (2005). Influences of exopolysaccharide producing cultures on the quality of plain set type yogurt. *Food Control*, 16:205–209.
- Hassan, A. N. and J. F. Frank (1997). Modification of microstructure and texture of rennet curd by using a capsule-forming non-ropy lactic culture. *J. of Dairy Research*, 64: 115-121.
- Hassan, A. N., J. F. Frank and M. Elsoda (2003). Observation of bacterial exopolysaccharide in dairy products using cryo-scanning electron microscopy. *Int. Dairy J.* 13: 755–762.
- Hassan, A. N., R. Ipsen, T. Janzen and K.B. Qvist (2003). Microstructure and rheology of yogurt made with cultures differing only in their ability to produce Exopolysaccharides. *J. Dairy Sci.* 86: 1632.
- Hassan, A. N., J. F. Frank, K. A. Schmidt and S.I. Shalabi (1996a). Rheological properties of yogurt made with encapsulated non ropy lactic cultures. *J. of Dairy Sci.*, 70: 2091-2097.
- Hess, S. J., R. F. Roberts and G. R. Ziegler (1997). Rheological properties of nonfat yogurt stabilized using *Lactobacillus delbrueckii* ssp. *bulgaricus* producing exopolysaccharide or using commercial stabilizer system. *J. of Dairy Sci.* 80: 252-263.
- Huyghebaert, A., K. Dewettinck and W. de Greyt (1996). Fat replacers. In *Fat Replacers-Ripening and Quality of Cheese*. IDF Bulletin 317:10-15. Brussels: Int. Dairy Federation.
- Isanga, J. and G. Zhang (2009). Production and evaluation of some physicochemical parameters of peanut milk yogurt. *LWT – Food Sci. and Technology*, 42: 1132-1138.
- Jaros, D., M. Jacob, C. Otto and H. Rohm (2010). Excessive cross-linking of caseins by microbial transglutaminase and its impact on physical properties of acidified milk gels. *Int. Dairy J.* 20: 321-327.
- Jimenez- Guzman, J.A., A.E. Flores-Najera, Cruz-Guerrero and M. Garcia-Garibay (2009). Use of exopolysaccharide-producing strain of *Streptococcus thermophilus* in the manufacture of Mexican panela cheese. *LWT-Food Sci. and Technology*, 42: 1508-1512.
- Keogh, M. K. and B. T. O’Kennedy (1998). Rheology of stirred yogurt as affected by added milk fat, protein and hydrocolloids. *J. of Food Sci.*, 63: 108-112.
- Ling, E.R. (1963). *A Textbook of Dairy Chemistry*. Vol.11, Practical, 3rd Ed., Chapman and Hall, London, UK.
- Lee, W. J. and J. A. Lucey (2010) Formation and physical properties of yogurt. *Asian-Australian J. of Animal Sci.* 23: 1127–1136.
- Leroux, J., V. Langendorff, G. Schick, V. Vaishnav and R. J. Mazoye (2003). Emulsion stabilizing properties of pectin. *Food Hydrocolloids*, 17: 455–462.
- Lorenzen, P. C. and E. Schlimme (1998). Properties and potential fields of application of transglutaminase preparations in dairying, *Bulletin. Int. Dairy Federation*, Brussels, 33: 47-53.
- Lorenzen, P.C., H. Neve, R. A. Mautne and E. Schlimme (2002). Effect of enzymatic cross-linking of milk proteins on functional properties of set-style yogurt. *Int. J. of Dairy Technology*, 55: 152-157.
- Lootens, D., F. Capel, D. Durand, T. Nicolai, P. Boulenger and Langendorff (2003). Influence of pH, Ca concentration, temperature and amidation on the gelation of low methoxyl pectin. *Food Hydrocolloids*, 17: 237–244.
- Lucey, J. A., M. Tamehana, H. Singh and P. A. Munro (1999). Stability of model acid milk beverage: effect of pectin

- concentration, storage temperature and milk heat treatment. *J. Texture Studies*, 30: 305-318.
- Maiolino, D. (2002). Functional starches improve mouthfeel in liquid products. National Starch and Chemical Company, Bridgewater, NJ, USA.
- Mehanna, N.M., M.E. Ibrahim and I. L. El-Nawasany (2013). Impact of some hydrocolloids on the physical characteristics and quality of non-fat yogurt. *Egypt. J. Dairy Sci.*, 41: 163-170.
- Mehanna, N.M. and A.S. Mehanna (1989). Study on use of stabilizer for improving some properties of cow's milk yogurt. *Egyptian J. Dairy Sci.* 17:289-296.
- Motoki, M. and K. Seguro (1998). Transglutaminase and its use for food processing. *Food Sci. Technol.* 9: 204-210.
- O'Sullivan, M.M., A.L. Kelly and P. F. Fox (2002). Influence of transglutaminase treatment on some physicochemical properties of milk. *J Dairy Res.*, 69: 433-442.
- Ozer, B., H. A. Kirmaci, S. Oztekin, A. Hayaloglu and M. Atamer (2007). Incorporation of microbial transglutaminase into non-fat yogurt production. *Int. Dairy J.* 17: 199-207.
- Perry, D. B. and D. J. McMahon (1998). Manufacture of low-fat Mozzarella cheese using EPS-producing culture. *J. of Dairy Sci.*, 81: 563-566.
- Perry, D. B., D. J. McMahon and C. J. Oberg (1997). Effect of EPS-producing cultures on moisture retention in low-fat Mozzarella cheese. *J. of Dairy Sci.*, 80: 799-805.
- Puvanenthiran, A., R. Williams and M. Augustin (2002). Structure and viscoelastic properties of set yogurt with altered casein to whey protein ratios. *Int. Dairy J.*, 12: 383-391.
- Saffon, M., V. Richard, R. Jimenez-Flores, S.F. Gauthier, M. Britten and Y. Pouliot (2013). Behavior of heat-denatured whey: buttermilk protein aggregates during the yogurt-making process and their influence on set-type yogurt properties. *Foods* 2:444–459.
- Sandoval – Castilla, O., C. Lobato – Calleros, E. Aguirre – Mandujano and R. J. Vernon – Carter (2004). Microstructure and texture of yogurt as influence by fat replacers. *Int. Dairy J.*, 14: 51-159.
- Şanlı, T., E. Sezgin, O. Deveci, E. Şenel and M. Benli (2011). Effect of using transglutaminase on physical, chemical and sensory properties of set-type yogurt. *Food Hydrocolloids*, 25: 1477-1481.
- Sato, Y., S. Kawabuchi, Y. Irimoto and O. Miyawaki (2004). Effect of water activity and solvent-ordering on intermolecular interaction of high-methoxyl pectins in various sugar solutions. *Food Hydrocolloids*, 18: 527–534.
- Shaker, R. R., B. Abu-Jdayil and R.Y. Jumah (2002). Rheological properties of a concentrated fermented product. Labneh, produced from bovine milk: effect of production method. *Int. J. of Food properties*, 5: 667-679.
- Szczesniak, A. S., M. A. Brandt and H. Freidman (1963). Development of standard roting scales for mechanical parameters of texture and correlation between the objective and the sensory methods of texture evaluation. *J. of Food Sci.*, 28: 397-403.
- Tamime, A.Y. (2006). *Fermented Milks*. Blackwell Science Ltd, Oxford, England.
- Tamime, A. Y. and R. K. Robinison (1999). *Yogurt: Science and Technology*, 2 nd ed. Woodhead Publishing Ltd., Cambridge, UK.
- Teggatz, J. A. and H. A. Morris (1990). Changes in the rheology and microstructure of ropy yogurt during shearing. *Food Structure*, 9: 133-138.
- Trancoso-Reyes, N., N. Gutiérrez-Méndez, D. R. Sepulveda and L. R. Hernández-Ochoa (2014). Assessing the Yield, Microstructure, and Texture Properties of Miniature Chihuahua-Type Cheese Manufactured with a Phospholipase A1

Improvement of the rheological and physical properties of non-fat

- and Exopolysaccharide-Producing Bacteria. *J. Dairy Sci.*,97: 598-608.
- Tsoga, A., R. K. Richardson and E. R. Morris (2004). Role of cosolutes in gelation of high-methoxyl pectin. Part 1. Comparison of sugars and polyols. *Food Hydrocolloids*, 18: 907–919.
- Towler, C. (1984). Sedimentation in a cultured milk beverage. *New Zealand J. of Dairy Sci. and Technology*,19: 205-211.
- Wang, W., Y. Bao, G. Hendricks and M. Guo (2012). Consistency, microstructure and probiotic survivability of goats' milk yogurt using polymerized whey protein as a co-thickening agent. *Int. Dairy J.*, 24, 113-119.
- Williams, P. A. and G. O. Phillips (2000). *Handbook of Hydrocolloids*. Cambridge and Boca Raton: Woodhead Publishing Limited and CRC Press LLC.
- Wosiacki, G. (1977). Enzimas pectinolíticas de *Fusarium oxysporum* Schlecht. Fr. Isolado de frutos de café, Thesis. São Paulo, Brasil: Universidade Estadual de Campinas.
- Wszolek, M., A.Y. Tamime, D.D. Muir and M.N.I. Barclay (2001). Properties of kefir made in Scotland and Poland using bovine, caprine, and ovine milk with different starter cultures. *Lebensmittel-Wissenschaft und-Technologie* 34:251-261.
- Wu, H., G. J. Hulbert and J. R. Mount (2001). Effects of ultrasound on milk homogenization and fermentation with yogurt starter. *Innovative Food Science and Emerging Technologies*, 1: 211-218.
- Yazici, F. and A. Akgun (2004). Effect of some protein based fat replacers on physical, chemical, textural, and sensory properties of strained yogurt. *J. Food Eng.*, 62:245–254.

تأثير استخدام انزيم الترانس جلوتامينيز الميكروبي وبعض المزارع البكتيرية المنتجة للسكريات العديدة كمحسنات قوام علي الخواص الطبيعية والريولوجية للزبادي خالي الدهن

دينا احمد مرتضى عامر⁽¹⁾، محمد يحيى علي الهواري⁽¹⁾، مرسى ابو السعود السودة⁽²⁾

⁽¹⁾ قسم علوم وتكنولوجيا الاغذية (ألبان). كلية الزراعة جامعة طنطا

⁽²⁾ قسم علوم وتكنولوجيا الألبان. كلية الزراعة جامعة الإسكندرية (الشاطبي).

الملخص العربي

تهدف هذه الدراسة التحسين الخواص الريولوجية والفيزيائية والحسية للزبادى خالى الدسم باستخدام انزيم الترانس جلوتامينيز الميكروبي، واحد أنواع البكتريا المنتجة للسكريات العديدة، وإثنين من المثبتات التجارية المعتمدة على البكتين والجيلاتين. وأوضحت النتائج أنه لا توجد فروق معنوية فى محتوى الرطوبة فى العينات الطازجة بين كل من عينات الزبادى خالى الدسم المنتجة بمحسنات القوام وعينة الزبادى كامل الدسم (المرجعية)، بينما توجد فروق معنوية بين هذه العينات وعينة الزبادى خالى الدسم (المرجعية). كما أوضحت النتائج انخفاض محتوى الرطوبة انخفاضاً معنوياً فى كل عينات الزبادى ماعدا فى عينات الزبادى المرجعية (كامل وخالى الدسم) بعد 10 أيام من التخزين. توجد أعلى نسبة رطوبة فى عينة الزبادى خالى الدسم المنتجة بأحد السلالات البكتيرية المنتجة للسكريات العديدة مقارنة بكل من عينات الزبادى المرجعية (كامل وخالى الدسم) والعينات المختبرة (المنتجة بالأنزيم، والبكتين التجارى، والجيلاتين التجارى) خلال فترة التخزين. كما أوضحت النتائج أن نسبة البروتين الى المادة الجافة أعلى فى كل من عينة الزبادى خالى الدسم (المرجعية)، وعينة الزبادى خالى الدسم المنتج باحد أنواع البكتريا المنتجة للسكريات العديدة، وعينة الزبادى خالى الدسم المنتج بالجيلاتين التجارى مقارنة بكل من عينة الزبادى خالى الدسم المنتج بالأنزيم وعينة الزبادى خالى الدسم (المرجعية). أظهرت النتائج أنه لا توجد فروق معنوية فى نسبة الدهن الى المادة الجافة بين عينة الزبادى خالى الدسم المرجعية وعينات الزبادى خالى الدسم المنتجة بمحسنات القوام، بينما توجد فروق معنوية بين هذه العينات وعينة الزبادى كامل الدسم (المرجعية) وذلك فى العينات الطازجة أو خلال فترات التخزين. لا توجد فروق معنوية فى نسبة الحموضة بين كل المعاملات. أنخفضت الصلابة انخفاضاً معنوياً فى الزبادى خالى الدسم المنتج باحد انواع البكتريا المنتجة للسكريات العديدة مقارنة بباقي المعاملات خلال فترة التخزين. كما أنخفض مستوى المرونة، واللزوجة، والمضغية فى كل من الزبادى خالى الدسم المنتج باحد أنواع البكتريا المنتجة للسكريات العديدة، والمثبتات التجارية بالمقارنة بعينة الزبادى كامل الدسم (المرجعية)، وعينة الزبادى خالى الدسم المنتج بالأنزيم. كما زاد الألتصاق زيادة معنوية خلال فترة التخزين فى كل المعاملات. لا توجد فروق معنوية فى نسبة التشرشيش بين كل المعاملات. أظهر الزبادى خالى الدسم المنتج باحد أنواع البكتريا المنتجة للسكريات العديدة، و المنتج بانزيم الترانس جلوتامينيز الميكروبي وكذلك المنتج بالجلاتين التجارى صفات ريولوجية وطبيعية قريبة للزبادى الكنترول كامل الدسم.