

PREVALENCE OF *STAPHYLOCCUS* AND *AEROMONAS* IN SOME SALTED DAIRY PRODUCTS

AHMED HASSAN SAAD¹; EHAB MOHAMED SALAMA¹; HANAN A. EL DAHSHAN²
and NOHA TALAAT ASSAF¹

¹ Faculty of Veterinary Medicine, Suez Canal University, Egypt

² Food Hygiene Department, Port-Said Branch, Animal Health Research Institute

Received: 18 November 2018; **Accepted:** 9 December 2018

ABSTRACT

120 samples of pickled white soft cheese (domiata cheese) and Mish were collected from local markets in Port Said governorate, Egypt. Samples were analyzed for sodium chloride level, determination of *Staph. Spp* at 3% and 10% Na CL, isolation of *Staphylococcus aureus* at 3% and 10% NaCL and Determination of *Aeromonas spp.* at 3% and 10% NaCL. Results obtained revealed that the mean values of the sodium chloride percentage were 3.7 ± 0.13 % in pickled domiata cheese samples and 6.1 ± 0.14 % in mish samples. Incidence of *Staph. spp* in Pickled domiata cheese was 85% at 3% NaCL and 71.6% at 10% NaCL. while in mish samples was 93% at 3% NaCL and 83.3% at 10% NaCL. Incidence of *Aeromonas spp.* at 3% NaCL was nil while at 10 % NaCL was 48.3% in pickled domiata cheese and in Mish 38.3% in. Incidence of *A. hydrophila*, *A. caviae*, *A. trota* and *A. schubertii* in Pickled domiata cheese were 25.7 %, 40%, 20% and 14.3%, respectively. While in Mish were 25%, 46.4%, 21.4% and 7.1% respectively.

Key words: Salted dairy products, halophilic bacteria, *Staph. Spp.*, *Aeromonas spp.*

INTRODUCTION

Pickled white soft cheese (domiata cheese) and Mish are well-known as local types of cheese in Egypt. Handling of milk during cheese manufacture considered the main source of microbial contamination of cheese which affecting on cheese quality and render it unfits for human consumption (Yousef *et al.*, 2001).

Staphylococcus spp. are considering one of halophilic bacteria which its presence in milk or milk products can cause a public health hazard, *Staphylococcus aureus* is responsible for food poisoning (ICMSF, 1986) *S. aureus* is consider as one of important food-borne pathogen. It is a versatile pathogen of humans and animals; it causes a wide variety of diseases ranged in severity from slight skin infection to more severe diseases such as pneumonia and septicemia (Lowy, 1998). *Staph. aureus* is capable of producing several enterotoxin cause food poisoning in human with varying degree and generally characterized by nausea, diarrhea, abdominal cramps and emesis

(Normanno *et al.*, 2005) and (Brightwell *et al.*, 2006). In the last few decades' Staphylococcal food poisoning (SFP) has been reported as third cause of food-borne illnesses in the world (Zhang *et al.*, 1998).

Aeromonas spp are considered one of the most important halophilic bacteria which are widely distributed in nature. Moreover it is considered as food borne pathogen (Palumbo *et al.*, 1992). *Aeromonas spp.* is pathogens that cause food-borne gastroenteritis, extra intestinal symptoms such as septicemia, meningitis, endocarditis and osteomyelitis with a high mortality rate in immune-compromised person (Gold and Salit, 1993). The mechanisms by which *Aeromonas spp.* cause diarrhea has been known that they produce enterotoxins, certain enzymes, they are able to adhere cell membranes and invade them (Kirov *et al.*, 1994). The presence of such bacteria in milk products may be due to direct contact with contaminated sources in dairy farm environment, excretion from udder of an infected animal or during processing of cheese. (Oliver *et al.*, 2005).

MATERIALS AND METHODS

1. Collection of samples

One hundred and twenty random samples of pickled white soft cheese and Mish (60 of each) were collected from different markets in Port Said city,

Corresponding author: HANAN A. EL DAHSHAN

E-mail address: mhananamen@yahoo.com

Present address: Food Hygiene Department, Port-Said Branch, Animal Health Research Institute

and then taken aseptically to the laboratory immediately, where it is subjected to chemical and microbiological examinations.

2. Preparation of samples (A.P.H.A. 2004)

250 gm. of each sample were transferred aseptically in a sterilized polyethylene sac and thoroughly mashed in a sterile blender before being emulsified in the diluent solution under aseptic condition then divided into two subsamples for chemical examination and bacteriological examination.

3. Chemical Examination

3.1. Determination of sodium chloride content (A.O.A.C, 2000)

3 grams of prepared sample were weighted in 200 ml Erlenmeyer flask. 25 ml of N/10 silver nitrate solution, 10 ml of halogen free nitric acid and 50 ml of distilled water were added and the mixture was boiled. 15 ml of 5% potassium permanganate solution in 5 ml portion were added during boiling (till the solution become yellowish and clear). Then the solution was allowed to cool then filtered into 200 ml volumetric flask, the filter paper was washed thoroughly with distilled water at 20°C, and the filtrate was made up to standard volume. The excess of silver nitrate in 100 ml of the clear solution was titrated against 0.1 N potassium thiocyanate solution (9.71 g / liter) using 2 ml of saturated solution of iron alum as indicator. The salt content was calculated according to the following equation:

$$\text{Na CL \%} = \frac{2 (25-R) \times 0.00584 \times 100}{\text{Weight (3g)}}$$

4. Microbiological Examination

4.1. Determination of total Halophilic bacterial counts (A.P.H.A.2004)

Preparation of serial dilution:

10gm of each sample was transferred in sterile stomacher bag with 90ml synthetic sea water solution (3% NaCL for Slight or 10%NaCL for moderate Halophilic counts). Apply in a stomacher

lab-blender (2000 rpm) used to homogenize the specimen for 2 minutes to make a 1:10 dilution (wt/vol) then decimal dilution of 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} and 10^{-6} were prepared.

0.1 ml from each dilution was inoculated in duplicated plates containing standard plate count agar following by spread plate technique and the plates were incubated at 37°C for 48 hours, then the total slight and moderate halophiles were counted and expressed as CFU/g.

4.2. Identification of isolated Staphylococci organisms according to (FDA, 2001).

Suspected colonies of Staphylococcus organisms were examined morphologically, microscopically according to (Ryan and Ray 2004)

4.3 Biochemical examination: according to (FDA, 2001)

The biochemical tests were Coagulase test, Catalase test, DNase test, Acetoin production, Oxidase test and D-mannitol fermentation.

4.4. Isolation of *Aeromonas spp.*

10gm of each sample were homogenized with 90 ml alkaline peptone water (with 3% or 10% NaCL) for 2 min, and then incubated for 24hr at 30°C (Villari *et al.*, 2000). A loopful from alkaline peptone water was subsequently plated on the surface of starch ampicillin agar plate and incubated for 48 hr. at 30°C. Typical yellow colonies of *Aeromonas* species were purified on tryptone soya agar then stained by Gram's stain (A.P.H.A., 2004) and confirmed on the basis of the following test: Oxidase test, resistance to vibriostatic agent o/129, esculin hydrolysis, sugar fermentation and gas production, indole production and voges- proskaur test.

4.5. Differentiation of common motile *Aeromonas* isolated.

From clinical specimens according to Carnhan *et al.* (1991) as modified by Joseph and Carnahan (1994). All bacterial isolates would be short G -ve bacilli, oxidase positive that ferment glucose.

RESULTS

Sodium chloride content

Table 1: Statistical analytical results of salt conc. % in Pickled domiati cheese samples (n=60) and Mish samples (n=60) comparison with Egyptian Standards 2005.

samples	Min.	Max.	Mean ± SE	E.S 2005
Pickled domiati cheese	2.9%	5.3%	3.7±0.135	Not more than 7.0%
Mish	4.2%	8.3%	6.1 ± 0.142	Not more than 15%

Determination of total Slight & moderate Halophilic counts

Table 2: Statistical analytical results of Total Slight and moderate Halophilic count in examined samples.

Examined samples (n= 60)		Positive samples		Count (cfu/g.)		
		No.	%	Min.	Max.	Mean ± SE
Pickled domiati cheese	3%Nacl	49	81.6	1 x10 ³	1 x10 ⁶	3.31 x10 ⁵ ±0.81 x10 ⁵
	10% Nacl	46	76.6	1 x10 ²	3.2x10 ⁵	0.76 x10 ⁵ ±0.14 x10 ⁵
Mish	3%Nacl	56	93.3	1.6 x10 ³	2.6 x10 ⁶	1.99 x10 ⁵ ±0.42 x10 ⁵
	10% Nacl	58	96.6	5 x10 ²	9 x10 ⁵	2.09 x10 ⁵ ±0.36 x10 ⁵

Detection of *Staphylococcal* organisms at 3% and 10% NaCL:**Table 3:** prevalence of *Staphylococci* isolated from examined samples.

Examined samples	No.	Positive samples			
		3%NaCL		10%NaCL	
		No.	%	No.	%
Pickled domiati cheese	60	51	85	43	71.6
Mish	60	57	93	50	83.3

4. Isolation of *Aeromonas spp.* at 3% and 10% Na CL**Table 4:** Incidence of *Aeromonas spp.* in examined samples at 10% NaCl.

Types of Examined samples	No. of Examined samples	Positive samples		isolates		<i>Aeromonas spp</i>							
		No	%	No.	%	<i>A. hydrophila</i>		<i>A. caviae</i>		<i>A. trota</i>		<i>A. schubertii</i>	
						No.	%	No.	%	No.	%	No.	%
Pickled domiati cheese	60	29	48.31%	35	58.3%	9	25.7%	14	40%	7	20%	5	14.3%
Mish	60	23	38.3%	28	46.6%	7	25 %	13	46.4%	6	21.4%	2	7.1%

DISCUSSION

Sodium chloride content

The use of salting is one of the classical methods of food preservation. Salting is used to extend the shelf-life of foods throughout civilization. In this study, it is evident in table (1) that salt percentage of examined pickled domiati samples ranged from 2.90 % to 5.30 %. While in mish samples ranged from

4.20 % to 8.30% all results obtained were within the normal range of (Egyptian Standards, 2005). Nearly similar results were reported by Ceylan *et al.* (2003), Hassan and Afify, (2007), Nawar, (2007), EL-Ansary *et al.* (2011), El Bakry, (2012), Yasser, (2015) and EL-Refaay, (2016), while higher result were obtained by Mohamed, (2004), Patrick *et al.*, (2004), Hayaloglu and Kirbag, (2007) and EL Zahar,

(2010). While lower result were obtained by Riad, (1996) and Sadek, (2009).

Determination of total Slight & moderate Halophilic counts

Slight, moderate, and extreme halophiles as those bacteria that grow best in media containing 2 to 5%, 5 to 20% and 20 to 30% salt respectively. (Kushner 1978). in table (2) Samples examination refer that maximum value of slight halophilic count in Pickled domiati cheese samples and Mish samples were 1×10^6 and 2.6×10^6 respectively while minimum value were 1×10^3 and 1.6×10^3 respectively where salt conc. was 3%. This results are nearly similar to Freitas *et al.* (1993), and Saad and Moawad (1999). While higher results are obtained by Riad (1996) and Omer *et al.* (2007).

The maximum value of moderate halophilic count in Pickled domiati cheese samples and Mish were 3.2×10^5 and 9.0×10^5 cfu/g. respectively, while minimum value was 1.0×10^2 and 5.0×10^2 cfu/g. Respectively with mean value $0.76 \times 10^5 \pm 0.14 \times 10^5$ in pickled domiati cheese samples. And with mean value $2.09 \times 10^5 \pm 3.6 \times 10^4$ in Mish samples (Where NaCL concentration is 10%). These results are nearly similar to El-prince (1994) and Riad (1996), while lower results are obtained by Omar *et al.* (2007).

Detection of *Staphylococcal* organisms at 3% and 10% NaCL:

The presence of *Staphylococcus* is an index of contamination from operators or workers. It has a potential significance to public health due to its ability to produce enterotoxin leading to food poisoning. The results in Table (3) show that incidence of *Staphylococcus* in Pickled domiati cheese was 85% at 3% NaCL and 71.6% at 10% NaCL. While in Mish samples was 93% at 3% NaCL and 83.3% at 10% NaCL. These results are nearly similar to those obtained by Abou El-Makarem, (2009). Higher results are obtained by Sheleby(2008) and Elshafey (2011).

Isolation of *Aeromonas spp.* at 3% and 10% Na CL:

Aeromonas species are widely distributed in the aquatic environment, including raw and processed drinking water (Holmes *et al.*, 1996), and have been frequently isolated from various food products such as fish and shellfish, raw meat, vegetables and raw milk Palumbo, (1996). It is associated with travellers' diarrhea (Hänninen *et al.*, 1995 and Yamada *et al.*, 1997). In this study, *Aeromonas* species was not isolated from any examined samples either Mish or Pickled domiati cheese using 3% salt concentration while at 10% Na CL, as shown in table (4) that the incidence of *Aeromonas spp.* was 25% *A. hydrophila*, 46.4% *A. caviae*, 21.4% *A. trota* and 7.1% *A. schuberti* in Mish cheese samples while the incidence of

Aeromonas spp. was 25.7% *A. hydrophila*, 40% *A. caviae*, 20% *A. trota* and 14.3% *A. schuberti* in Pickled domiati cheese samples. This results are nearly similar to those obtained by to Freitas *et al.* (1993), Khalil (1997), Effat *et al.* (2000), Ahmed *et al.* (2014) Higher results are obtained by Yasser (2008), Nazem *et al.* (2010) and Alhazmi (2015).

REFERENCES

- Abou El-Makarem, H.S.M. (2009): Plasmid profile from several pathogenic microorganism strains isolated from market kareish cheese M.V.Sc. Thesis, Faculty of Vet. Med. Alex. University.
- Ahmed, N.I.; Abd-Aal, S.F.; Madeha, A.A. and El-Sayed, M.S. (2014): Enumeration of *Aeromonas spp.* isolated from milk and some dairy products in Sharkia Governorate Egypt. Alexandria journal of veterinary science 40 (1), 52-63.
- Alhazmi, M.I. (2015): Isolation of *Aeromonas spp.* from food products: Emerging *Aeromonas* infections and their significance on public health. Journal of AOAC international, 97(4): 927-929.
- A.P.H.A. (2004): American Public Health association: Standard Methods for the Examination of Dairy Products. 17th ed, APHA, Washington D.C. USA.
- A.O.A.C. (2000): Association of Official Analytical Chemists 17th Ed. pub. A.O.C.A. POBOX540, Benjamin Franklin station Washington, D.C.
- Bightwell, G.; Clemens, R. and Boerma, J.A. (2006): Evaluation of molecular methods to determine enterotoxigenic status and molecular genotype of bovine, ovine, human and food isolates of *Staphylococcus aureus*. International Journal of food Microbiology, 107 (2): 192-201
- Carnahan, A.M.; Behram, S. and Joseph, S.W. (1991): AerokeyII: a flexible key for identifying clinical *Aeromonas* species. Journal of clinical Microbiology 29 (12); 2843-2849.
- Ceylan, Z.; Turkoglu, H. and Dayisoğlu, K.S. (2003): The microbial and chemical quality of Sikma cheese produced in Turkey. Pakistan journal of Nutrition, 2(2): 95-97.
- EL-Refaay, D.G.A. (2016): Mycological studies on some types of cheese. M.V.Sc., Thesis., Fac. Vet. Med. Sues Canal University.
- Egyptian Standards (2005): Mish cheese. Egyptian Organization for Standardization and Quality Control, Es.4342/2008.
- Effat, B.A.; Hosny, I.M. and Dabiza, N.M. (2000): Occurrence of *Aeromonas hydrophila* and its growth in Egyptian soft cheese. Egyptian Journal of Dairy Science, 28 (1): 1-12.

- EL-Ansary, M.A.M. (2011)*: Hygienic condition of locally manufactured cheese. Ph. D. Thesis, Fac. Vet. Med. Alex. Univ., Egypt.
- El-Bakry, M. (2012)*: Salt in cheese., current research in dairy sciences 4(1):1-5. Department of dairy science and technology, faculty of Agriculture, Cairo university, Giza, Egypt.
- El shaffy, M.N. (2011)*: Genotypic characterization of *Staphylococcus aureus* as food poisoning organism from milk and dairy products., department of Bacteriology- immunity-mycology. M.V.Sc. Thesis. Fac. Vet. Med. Suez Canal Uni.
- El-Zahar, K.M.; Hendawi, M.Y. and Abdel-Zaher, A.M. (2010)*: Evaluation of biogenic amines in Egyptian cheese markets: the effects on physiological parameters of rats. Egyptian Journal of Dairy Science, 38 (2) 219-230.
- El-prince, E.M. (1994)*: Effect of potassium sorbate on the microbiological quality of milk and some milk products. Ph.D. Thesis, Fac. Vet. Med. Assiut University.
- FDA; Food Drug Administration (2001)*: Analytical manual. Chapter12, *Staphylococcus aureus*. Reginal W.B and Gayle A.I.
- Freitas, C.A.; Marly, P.N.; Arlete, M.M. and Ilvan, D. Ricciadi (1993)*: Occurance and characterization of *Aeromonas* species in pasteurized milk and white cheese in Rio de Janeiro, Brazil Journal of food protection. Vol.56(1): 62-65.
- Gold, W.L. and Salit, I.E. (1993)*: *Aeromonas hydrophila* infections of the skin and soft-tissue: Report of 11 cases and review. Clin. Infect. Dis., 16: 69-74.
- Hanninen, M.L.; Salmi, S.; Mattila, L.; Taipalinen, R. and Siitonen, A. (1995)*: Association of *Aeromonas* spp. with travellers' diarrhea in Finland. Journal of Medical microbiology 42, 26,31.
- Hassan, G.M. and Afify, S.I. (2007)*: Occurance of some pathogenic microorganisms in Kareish cheese and their puplic health significance. Beni Suif vet. Med. Journal pp.142-150.
- Hayaloglu, A.A. and Kirbag, S. (2007)*: Microbial quality and presence of moulds in Kuflu cheese. International journal of food microbiology,115(3): 376-380.
- Holmes, P.; Niccolls, L.M. and Sartory, D.P. (1996)*: The ecology of mesophilic *Aeromonas* in the aquatic environment. In the Genus *Aeromonas* ed. Austin, B., Altwegg, M., Gosling, P.J. and Joseph, S.pp.127-150. Chichester, UK: Wiley and Sons.
- ICMSF (International Commission on Microbiological Specification for Food (1986)*: Microbial Ecology of Food, Vol.1-2. University of Toronto Press.
- ISO (1999)*: International organization for standardization. Microbiology of food and animal feeding stuffs- Horizontal method for the detection of *Staphylococcus*.
- Mohamed, I.M.J. (2004)*: Biogenic amines and their forming organisms in cheese. Ph.D. Thesis, Fac. Vet. Med, Suez Canal Uni.
- Joseph, S.W. and Carnahan, A. (1994)*: The isolation, identification and systematics of the motile *Aeromonas* species. Annual review of fish disease. 4, 315-343.
- Khalil, N.G. (1997)*: Incidence of *Aeromonas hydrophila* group in raw milk and some dairy products in Assuit City. Assuit Vet. Med. J. 37(73): 100-108.
- Kushner, D.J. (1978)*: Life in high salt and solute concentrations: halophilic bacteria, in D.J. Kushner, Ed. Microbial life in Extrem Enviroments, Academic press, New York. p. 317
- Kirov, S.M.; Hudson, J.A.; Hayward, J. and Mott, S.J. (1994)*: Distribution of *Aeromonas hydrophila* hybridization groups and their virulence properties in Australasian clinical and environmental strains. Lett. Appl. Microbiol., 18: 71-3.
- Lowy, F.D. (1998)*: *Staphylococcus aureus* infection. N. Engl. J. Med. 339: 520-532.
- Nawar, D.M.M. (2007)*: Toxicoinfection organisms in milk and some street-vended dairy products. Ph. D. Thesis, Fac. Vet. Med., Alex. Univ., Egypt.
- Nazem, A.M.; Amer, A.A. and Soukayna, Abd El-Ahad (2010)*: Prevalance of some food poisoning micoorganisms in some dairy products. Alex.j. vet.30 (1): 1-6.
- Normanno, G.; Firinu, A.S.; Virilio, G.; Mule, A.D.; Poggiu, A.; Decastelli, L.; Mioni, R.; Scuta, S.; Bolzoni, G.; Di Giannatale, E.; Salinetti, A.P.; La Salandra, G.; Bartoli, M.; Zuccon, F.; Pirino, T.; Sias, S.; Parisi, A. and Celano, G.V. (2005)*: Coagulase-positive *Staphylococci* and *Staphylococcus aureus* in food products in marketed in Italy. International journal of food microbiology. 98. (1) (15): 73-79.
- Oliver, S.P.; Jayarao, B.M. and Almeida, R.A. (2005)*: Food borne pathogens in milk and the dairy farm environment. Food safety and public health implications, 2(2): 115-129.
- Omar, H.; Hoda, A.E.; Wafaa, M.B. and Naglaa F.G. (2007)*: Bacteriological quality of some dairy products (Kariesh cheese and Ice cream) in Alexandria. Microbiology Dep. Alex. University. Food hygein and control Division, Nutrition Department, Egypt puplic Health Assoc.Vol.82.No.586.
- Palumbo, S.A.; Abeyta, C. and Stelma, G. (1992)*: *Aeromonas hydrophila* group. In compendium of methods for the microbiological examination of foods (Eds Vanderzat, C. and Splittstoesser, D.F.) PP.497-515, Washington, D.C., American Public Health Association.

- Palumbo, S.A. (1996): The *Aeromonas hydrophila* group in food. in The Genus *Aeromonas* ed. Austin, B., Altwegg, M., Gosling, P.J. and Joseph. S., pp.287-310. Chichester, UK Wiley and Sons.
- Patrick, F.; Fox, M.; Cogan, T.M. and Timothy, G. (2004): Cheese Chemistry, Physics and Microbiology: Major Cheese Groups: Major Cheese Groups v. 2.
- Rayan, K.J. and Ray, C.G. (Eds). (2004): Sherris Medical Microbiology 4th ed. McGraw Hill, New York.
- Riad, A.M.A. (1996): Microbial monitoring for some dairy products as indices of sanitary quality, Ph. thesis. Fac. Vet. Med. Alex. Uni.
- Saad, A.H. and Maawad, A. (1999): Incidence of slight and moderate Halophiles in some selected food. Beni-Suef Veterinary Medical Journal, 9(3-A): 37-49.
- Sadek, S.M. (2009): Grading quality of some locally manufactured milk products. M.V.Sc. Thesis, Fac. Vet. Med., Alex Univ., Egypt.
- Sheleby, H.H.A. (2008): Indicator Organisms in Street-Vended Dairy Products. M.V.Sc. Thesis, Fac. Vet. Med., Minufiya University.
- Villari, P.; Crispino, M.; Montuori, P. and Stazione, S. (2000): Prevalance and Molecular characterization of *Aeromonas* spp. in Ready-to-Eat foods in Italy. Department of Health and Praventive Science. University Federico II. Journal of food protection. 63(12): 1754-1757.
- Yamada, S.; Matsushita, S.; Dejsirilet, S. and Kudoh, Y. (1997): Incidence and clinical symptoms of *Aeromonas*-associated Traveller's diarrhea in Tokyo. Epidemal infect. 199, 121-126.
- Yasser, A.M. (2008): Prevalance of *Aeromonas hydrophila* group in milk and milk products., Cairo university, faculty of Veterinary Medicine, department of food hygiene.
- Yasser, S.I.S. (2015): Incidence and Biocontrol of Staph. Aureus in some milk products. M.V. SC. Thesis. faculty of Vet. Med. Suez Canal University.
- Yousef, H.; Sobieh, M. and Nagedan, K. (2001): Microbial status of Domiati cheese at Al Gassiem area, Saudi Arabia., 8th sci. Cong., The fac. vet. Med. Assiut University, pp: 91-97.
- Zhang, S.; Iandolo, J.J. and Stewart, G.C. (1998): The enterotoxin D plasmid of *Staphylococcus aureus* encodes a second enterotoxin determinant (sej). FEMS Microbiology Letters, 168: 227-233.

تواجد انتشار الاستافيلوكوكس اوريس والايرومونات في الجبن الدمياطى المخزن والمش

احمد حسن علي سعد ، ايهاب محمد سلامه ، حنان امين مصطفى الدهشان ، نهى طلعت عبد الفتاح عساف

Email: mhananamen@yahoo.com Assiut University web-site: www.aun.edu.eg

اجريت هذه الدراسه على ١٢٠ عينه من الجبن الدمياطى المخزن والمش ٦٠ عينه من كل نوع. تم تجميع العينات عشوائيا من بعض المحلات والاسواق والباعة الجائلين في كل من محافظة بورسعيد وتم تقييم هذه العينات. وقد اظهرت النتائج مايلي:

الفحص الكيمايى:

متوسط نسبة ملح الطعام في عينات الجبن الدمياطى المخزن ٠,١٣ ± ٣,٧ بينما كانت في المش كانت ٠,١٤ ± ٦,١

الفحص الميكروبيولوجي:

١- متوسط العدد الكلى للبكتيريا طفيفة درجة الملوحة فى عينات الجبن الدمياطى المخزن والمش كان ١,٩٩ x ١٠^٥ ± ٤٢ x ١٠^٥ و ٣,٣١ x ١٠^٥ ± ٨١ x ١٠^٥ على التوالي

٢- متوسط العدد الكلى للبكتيريا متوسطة درجة الملوحة فى عينات الجبن الدمياطى المخزن والمش كان ١,٤ x ١٠^٤ ± ٧,٦ x ١٠^٤ و ٣,٣٦ x ١٠^٥ ± ٢,٠٩ x ١٠^٥ على التوالي

٣- مدى تواجد ميكروب المكورات العنقودية فى عينات الجبن الدمياطى المخزن عند ٣% ملح كانت ٨٥% بينما عند ١٠% ملح كانت ٧١,٦%.

٤- مدى تواجد ميكروب المكورات العنقودية فى عينات المش عند ٣% ملح كانت ٩٣% بينما عند ١٠% ملح كانت ٨٣,٣%

٥- لا وجود على الاطلاق لميكروب الايرومونات فى اى من عينات الجبن الدمياطى المخزن او المش عند تركيز ملح ٣% بينما عند تركيز ملح ١٠% كان مدى تواجد انواع ميكروب الايرومونات كما يلى: ٢٥,٧% أ.هيدروفيليا، ٤٠% أ.كافيه، ٢٠% أ.تروتا، و ١٤,٣% أ.اسكوبرتى فى عينات الجبن الدمياطى المخزن اما عينات المش فكانت ٢٥% أ.هيدروفيليا، ٤٦,٤% أ.كافيه، ٢١,٤% أ.تروتا و ٧,١% أ.اسكوبرتى.