

SOME PARASITIC CAUSES OF DIARRHEA IN CALVES AT BEHERA GOVERNORATE

REDA SAMIRAHMED FADLY

Animal Health Research Institute, Damanhour Branch.

ABSTRACT

Received at: 29/5/2012

Accepted: 14/7/2012

Three hundred faecal samples from diarrheic calves were examined using concentration flotation technique using saturated salt solution to detect *Eimeria* Spp. Oocysts and using Sheathers solution to detect *Giardia* species, Modified Ziehl-Neelsen staining technique to detect *Cryptosporidium parvum*. The results indicated that 158 (52.66%) were positive for protozoal parasites as *Eimeria* species (27%), *Cryptosporidium parvum* (14.66%) and *Giardia* species (11%). Parasitological examination revealed that isolated five species of *Eimeria* in calves were *E. bovis*, *E. zuernii*, *E. subspherica*, *E. cylindrical* and *E. ellipsoidalis*. *E. bovis* and *E. zuernii* (27.1%) and (23.4%) were the most prevalent species infecting calves followed by *E. subspherica* (14.8%), then *E. ellipsoidalis* (13.5%) finally *E. cylindrical* (9.8%). *Eimeria* Oocysts were demonstrated in all age groups of animals. However, it was evident that the prevalence of the disease was significantly higher in 16-20 weeks old (46.6%). While *Cryptosporidium parvum* Oocysts was found to be decreased with increasing age of calves. It was found the most affected age group was between 1-4 weeks old (32%) and Cysts of *Giardia* spp. were found to be higher at age group 4-8 weeks old (30%). Seasonality over all prevalence of *Eimeria* spp., *Cryptosporidium parvum* and *Giardia* spp. was higher in samples taken during winter than samples taken during summer.

Key words: *Eimeria* spp., *cryptosporidium*, *diarrhea* in calves.

INTRODUCTION

The most important cause of calf morbidity and mortality is diarrhea (Heat, 1992), Neonatal calf diarrhea continues to be the 1st cause of calf mortality in Egypt which ranges between 27.4-55% of the total death in young calves (Ahmed, 1980).

Cryptosporidium, *Giardia* and *Eimeria* species are among the most common and important enteropathogens in calves (Maddox – Hyttel *et al.*, 2006; Bartles *et al.*, 2010). The most important clinical sign they may cause is watery diarrhoea, occasionally containing blood in *Eimeria* infection. Other less obvious signs are fatigue, fever and reduced appetite (Fitzgerald, 1980). High infection rate, and lack of preventive and therapeutic measures when clinical disease manifests, can lead to outbreaks and high mortality among calves. Moreover, the sub clinical effects of infection, which often result in recovery but lower weight gain, are also important to animal health and productivity (Daugeschies *et al.*, 1986). Bovine coccidiosis has a world wide distribution and result in economical losses. More than twelve different species of *Eimeria* in cattle and buffalo have been documented up till now (Abebe *et al.*, 2008).

Most commonly pathogenic species are *E. bovis* and *E. zuernii* causing mortality and morbidity by

disturbing absorption mechanisms (Cicek *et al.*, 2007).

Coccidiosis in cattle is observed in all age groups but it is most common and important in young animals, in associations with other enteropathogens coccidia have been indicated as an important cause of diarrhea in calves. *Cryptosporidium parvum*, first reported as entero pathogen in a calf (Panciera *et al.*, 1971), has now been reported as prevalent and wide spread in calves from 4 days to 4 weeks of age by numerous investigators in many countries. Clinical neonatal bovine Cryptosporidiosis is usually a result of infection with *C. parvum*. *Cryptosporidium* has direct life cycle which is completed in 3 days and transmission is fecal – oral. Cryptosporidiosis is considered to be zoonotic disease (Miron, 1991) and the most common route of infection in close contact with diarrheic feces. Due to the limited availability of effective drugs the control of Cryptosporidiosis relies mainly on hygienic measures and good management (De Graaf *et al.*, 1999).

Giardia intestinalis (synonyms *G. duodenalis* and *G. lamblia*) is globally distributed and is a common protozoan parasite in many species of animals and man. The public health implication of Giardiasis are well known and its effect on calf performance is not conclusive. The earliest reports of *Giardia* were in India and Egypt were made by (Deshpande and

Shostri, 1981) and (Suvillan *et al.*, 1988)

There is a close association between the prevalence of the *Cryptosporidium* and *Giardia* infection and age of animal (Lassen, 2011).

The aim of this study was to:

- 1- Estimation the prevalence, age and seasonal distribution of *Cryptosporidium*, *Giardia* and *Eimeria* infection among calves.
- 2- Identification of various species of *Cryptosporidium*, *Giardia* and *Eimeria* affecting calves.

MATERIALS and METHODS

1- Animals: a total number of 300 fecal samples from cow calves from May 2011 to April 2012 were collected from different localities in Behera Governorate. Calves aged from 1 week to 20 weeks.

2- Samples: faecal samples were collected directly from the rectum using plastic hand gloves, each glove being marked to indicate animals number age and date of sampling.

Parasitological Examination.

1- Concentration flotation technique: according to Soulsby (1982), using saturated salt solution for detection of *Eimeria spp. oocysts* and Sheathers solution for detection of *Giardia spp.cysts*.

2- Sporulation of *Eimeria species* oocysts: according to Soulsby (1982) the positive fecal samples were sporulated using 2.5% potassium dichromate solution for one week.

3- Modified Ziehl- Neelsen staining technique (MZN): was used for diagnosis of *Cryptosporidium parvum* oocyst according to Henriksen and Pohlenz (1981). Thin fecal smears were prepared on glass slides, dried at room temp., fixed in methanol 96% for 2-5 min., then fixed in flame samples were stained in concentrated carbol fuchsin for 20-30 min. The discoloration procedure was realized with 3% hydrochloric acid for 20-60 seconds, then smears were washed with running water and counter stained with solution of malachite green for 3- 5 min. rinsed in tap water and air dried.

4- Cysts of *Giardia* were found by floating the sample in Sheathers solution which has been found superior in getting *Giardia* Cysts to float, dropwise application of dilute Lugol's iodine solution (10.5%) to facilitate microscopic examination of *Giardia* according to Levine (1985).

5- Smears were examined under light microscope with x40 and x 100 magnifications, Non sporulated

respectively.

oocysts, sporulated oocysts and cysts size were measured using bright field microscopy with a calibrated eye piece micrometer.

6- Identification of different oocysts & Cysts, according to Levine (1985).

RESULTS

Prevalence of *Cryptosporidium* oocyst, *Giardia* cyst. and *Eimeria* oocysts:

As shown in Table (1) out of the examined 300 faecal samples from diarrheic calves 158 (52.66%) were positive for oocysts of total *Eimeria spp.*, total *Cryptosporidium parvum* and total Cysts of *Giardia spp.* 27%, 14.66% and 11%. Table (2) Parasitological examination revealed that isolated five species of *Eimeria* in calves were *E.bovis*, *E.zuernii*, *E.subspherica*, *E.cylindrica* and *E.ellipsoidalis*. The most prevalent species infecting calves were *E.bovis* and *E.zuernii* (27.1%), (23.4%) followed by *E.subspherica* (14.8%) and *E.ellipsoidalis* (13.5%) respectively. The lowest prevalent was *E.cylindrica* (9.8%). Mixed infection involved two or three *Eimeria* species was observed in 11.11% of the samples.

Table (3) *Eimeria* oocysts were demonstrated in all age groups of animals. However, it was evident that the prevalence of the disease was significantly higher at 16 - 20 ws (46.6%). While *Cryptosporidium parvum* appeared to be inversely related to age, where the prevalence of oocysts was found to be decreased with increasing age of calves. It was found the most affected age group was between 1-4 ws (32%). Cysts of *Giardia spp.* were found high at age group 4-8 ws (30%). Table (4) for seasonal variations reveal that over all prevalence of *Eimeria spp.*, *Cryptosporidium parvum* and *Giardia* cysts, was higher in samples taken during winter than samples taken during summer.

Morphology of *Cryptosporidium* oocyst, *Giardia* cyst. and *Eimeria* oocysts:

Fig. (1) Microscopical examination of Acid-fast stained smears revealed ovoid oocysts of *Cryptosporidium*, measured of 4-6 um in diameter. The oocysts were spherical to ovoid in shape stained red to pink with granular appearance against green back ground.

Fig (2). Microscopical examination of iodine stained smears revealed *Giardia* cyst is oval cyst measured 7-11 um long and 7-9 um wid and a 0.3 um thick outer wall of cystic wall internally there are four nuclei. Figs (3-7) represent non sporulated and sporulated oocysts of *Eimeria spp.*

Table 1: Incidence of Protozoan Parasites Infect Calves at Behera Governorate.

No. Of Ex. calves	Single <i>Eimeria</i> spp.		Single <i>Crypto.</i> Spp.		Single <i>Giardia</i> Spp.		Mixed infection <i>Eimeria</i> + <i>Crypto</i> (2) <i>Eimeria</i> + <i>Giardia</i> (3) <i>Eimeria</i> + <i>Giardia</i> + <i>Crypto</i> (1) <i>Giardia</i> + <i>Crypto</i> (2)	Total <i>Eimeria</i>		Total <i>Crypto.</i>		Toal <i>Giardia</i>		Total infec. Calves	
	No	%	No	%	No	%		No	%	No	%	No	%	No	%
300	75	25	39	13	27	9		81	27	44	14.66	33	11	158	52.66

Table 2: Prevalence and Characters of Different *Eimeria* spp. Isolated from Calves

<i>Eimeria</i> spp.	No.	%	Shape	Size um	Micropyle	Sporulation Time days
<i>E.bovis</i>	22	27.1	Ovoid	27x 20.3	Present	2 – 3
<i>E.zuernii</i>	19	23.4	Spherical	17.8x 15.6	No	2 – 3
<i>E.subspherica</i>	12	14.8	Round or Subspherical	11 x 10.4	No	4 – 5
<i>E. cylindrica</i>	8	9.8	Cylindrical	15 x 12	No	2 – 3
<i>E.ellipsoidalls</i>	11	13.5	Ellipsiod to slightovoid	12 x 8	No	2 – 3
Mixed in fection	9	11.11				

Table 3: Prevalence of Different Protozoan Infection in Relation to Age of Calves

Age	No. of exam calves	<i>Eimeria</i>		<i>Cryptosporidia</i>		<i>Giardia</i>	
		No.	%	No.	%	No.	%
1 – 4 ws	50	7	14	16	32	3	6
4 – 8 ws	50	8	16	14	28	15	30
8 – 12 ws	75	26	34.6	8	10.6	6	8
12 – 16 ws	50	5	6.1	5	10	5	10
16 – 20 ws	75	35	46.6	1	1.3	4	9
Total	300	81	27	44	14.66	33	11

Table 4: Seasonal Variations of Prevalence of Different Protozoal Infection

Seasons	No. of exam calves	<i>Eimeria</i>		<i>Crypto.</i>		<i>Giardia</i>	
		No.	%	No.	%	No.	%
Spring	75	12	16	8	10.6	3	4
Summer	75	10	13.3	4	5.3	4	5.3
Autumn	75	25	33.3	13	17.3	11	14.6
Winter	75	34	45.3	19	25.3	15	20




Fig (1)



Fig (2)




Fig (3)




Fig (4)




Fig (5)



Fig (6)



Fig (7)

Fig (1) Oocysts of *Cryptosporidium parvum* stained with Modified Ziehl-Neelsen stain X100

Fig (2) Cysts of *Giardia spp.* stained with iodine stain X40

Fig (3) Non sporulated and sporulated oocysts of *E. bovis* X40

Fig (4) Non sporulated and sporulated oocysts of *E. zuernii* X40

Fig (5) Non sporulated and sporulated oocysts of *E. subspherica* X40

Fig (6) Non sporulated and sporulated oocysts of *E. cylindrica* X40

Fig (7) Sporulated oocysts of *E. Ellipsoidalisis* X40

DISCUSSION

Cryptosporidium, *Giardia* and *Eimeria* infections have been reported from calves in many parts of the world (Hannes *et al.*, 2006).

The parasitological examination as shown in Table (1) revealed that 158 (52.66 %) of calves examined were infected with one or more parasitic protozoan agents. The result of this study confirmed the fact that, protozoal infection were wide spread in diarrhoeic calves, this is in accordance with previous studies (Goz. *et al.*, 2006 and Gul *et al.*, 2008) in Turkey.

Eimeria oocysts were identified in (27 %) calves, these results are nearly to that recorded by Gul *et al.* (2008) 22.53% in Turkey. Over all prevalence of *Eimeria* in present study is lower than perviously reported by Gasmir *et al.* (2011) 47.59% in Sudan and Yu *et al.* (2011) 34. 94% in China while our results was higher than this reported by El. Sherif *et al.* (2000) 12 . 05% in Egypt. This variation is most likely attributed to the differences in management and hasbandry practice of the study animals in different countries. Table (2) Five *Eimeria* species were identified in calves in this study namely *E. bovis*, *E. Zuernii*, *E. Subspheica*, *E. cylindrica* and *E. ellipsoidalis*, similar number of species has been described in surveys performed by El Sheriff *et al.* (2000) and Mary (2003).

Table (2) demonstrared that the prevalence of *E. bovis* and *E zuernii*, (27.1%) and (23.4%) which are most prevalent species for calves this concurs with the findings of Harfoush *et al.* (2005), Cicek *et al.* (2007), Abebe *et al.* (2008) and lassen *et al.* (2011).

Table (4) The prevalence of *Eimeria* spp. oocysts were highest in the age group of the 16-20 weeks old, this results were in accordance with the finding of Cicek *et al.* (2007); Abebe *et al.* (2008) and Priti *et al.* (2008), this data is in contrast with Sanchez *et al.* (2008) who found high percentage of infection 21–45 day.

Seasonal prevalence of coccidiosis among calves was higher in Winter (45.3%) Followed by Autumn (33.3%), Spring (16%) and Summer (13.3%) Table (4). Similar results were recorded by El Sheriff *et al.* (2000) and Mary (2003) in Egypt who reported higher prevalence during month of rain, while Sanchez *et al.* (2008) found that highest incidence in spring and autumn than winter and summer. This difference may be due to the different localities, seasonal and climate variation.

Cryptosporidium and *Giardia* infections have been reported for cattle in many parts of the world, prevalence data have often varied markedly. The high prevalence of *Cryptosporidium* and *Giardia* in neonates and young calves is well known and unambiguously age dependent (Xiao and Herd 2004). The findings of the present study showed of *Cryptosporidium*.oocysts were identified in (14.66%) in neonat calves by Modified Ziehl Neelsen staining technique, these results were relatively Similar to the

values recorded by EL-Shemy, (2006), El.Khodery and Osman (2008), Amal and Reda (2009) and El Kelesh *et al.* (2009) (14%,14.19%, 14% and 14%). While our result is relatively lower than the values recorded by, El sherif *et al.* (2000) 21.92 % and Majeed *et al.* (2011) 38.8% Table (1). On other hand our result was higher that obtined by Cokline *et al.* (2009) 6.2%. These differences might be attributed to the difference in localities, breeding and management system. *Cryptosporidium* oocysts were found in all age categories but with high prevalence in 1-4ws (35%), these results revealed that Cryptosporidiosis is a disease of young ages and gradual decrease of infection rate occurred with age progression, these finding were emphasized by Bartels *et al.* (2010). Mean while, O'Handley *et al.* (1999) stated that *Cryptosporidium* was important pathogen when calves were less than one month, also EL Sherif *et al.* (2000); El. Dessouky and Nabila (2005); Amal and Reda (2009) and Majeed *et al.* (2011) said that calves less than one month recorded the highest degree of infection.

As shown in Table (4), the highest incidence of infection was recorded in winter (25.3%), followed by spring (17.3 %), autumn (10.8 %) and summer 8%, Bendali *et al.* (1999) and Lefay *et al.* (2001) found that the highest incidence of Cryptosporidiosis was in winter and lower infection rate was in summer.

From the presnt study *Giardia* spp.cysts were identified in (9%) of calves, these result was similar to Iburg *et al.* (1996) 10.4% in Denmark and Gul *et al.* (2008) 9.3%, While higher result observed by, Ruest *et al.* (1998) 26.6% in swedan, and Olson *et al.* (2004) 73 % British Columbia, While lower result observed by Xiao (1994) 5.6% These difference may be due to the different localities , seasonal and climate variation Table (1).

This highest rate of prevalence of *Giardia* in the age group of the 4-8 weeks old these result similar to Davoudi *et al.* (2011) who found the highest dgree of infection in 8 weeks old 54.38% Table (3). Seasonal prevalence of *Giardia* among calves highest rate of incidence was detected in winter season (20%) this result agree with Mistic *et al.* (2006) and this result disagree with Davoudi *et al.* (2011) who found the highest degree of incidence was in summer(37.6%) and least incidence was in winter (16.8%) These differences might be attributed to the difference in localities, breeding and management system Table (4).

CONCLUSION

This study reveals that *cryptosporidium* spp., *Giardia* spp., and *Eimeria* spp., are the most Frequent parasites in calves at Behera Governorate These infections play an important role in animal production as it has been demonstrating that these parasites are able to cause diarrhea which negatively inflnuce growth,impair feed conversion and reduce milk production.

Giardia and *Cryptosporidium* may also act as an infection source for humans by direct contact with contaminated faces or water.

REFERENCES

- Abebe, R.; Wossene, A. and Kumsa, B. (2008): Epidemiology of *Eimeria* infections in calves in Addis Ababa and Debre Zeit dairy farms, Ethiopia. Intern. J. Appl. Res. Vet. Med 6: 24–30.
- Amal, A.E. and Reda, S.F. (2009): Studies on *Cryptosporidium* infection in calves in Behera Province. EVMS PJ, 5:37-47.
- Ahmed, A.A. (1980): Calf scoures in Egyptian buffalo - Cows. Egyptian German Seminar on the Mortality of Newly Born Calves: 19–21.
- Bartels, C.J.; Holzhausr, M.; Jorristma, R.; Swart, W.A. and Lam, T.J. (2010): Prevalence, prediction and risk factors of entero pathogens in normal and non – normal faces of young Dutch dairy calves. 93: 162–169.
- Bendali, F.; Sanaa, M.; Bichet, H. and Schelcher, F. (1999): Risk factors associated with diarrhoea in newborn calves. Veterinary Research 30: 509-522.
- Cicek, H.; Sevimli, F.; Kozan, E.; Kose, M.; Eser, M. and Dogan, N. (2007): Prevalence of Coccidian in beef cattle in Western Turkey. Parasitol. Res., 101: 1239–1243.
- Cokline, T.; Uehinger, F.D.; Farber, J.M.; Barkema, H.W.; O'handley, R.M. and Dixon, B.R. (2009): Prevalence and molecular characterization of *Cryptosporidium* spp. in diary calves from 11 farms in PrinceEdward Island, Canada. J. Vet. Parasitol. 160: 323-326.
- Dauguschies, A.; Akimaru, M. and Burger, H.J. (1986): Experimentelle *Eimeria bovis* – infektionen beim kalb: 1. Parasitologishche und Klinische Befumle. Dtsch. Tieraztle. wschr., 93: 393–397.
- Davoudi, Y.; Garedaghi, Y. and Safarmashaei, S. (2011): Epidemiological study of Giardiasis in Diarrheic calves in East-Azerbaijjan Province, Iran. J. Aim. Vet. Adv., 10: 2508-2510.
- De Graaf, D.C.; Vanopdenbosch, E.; Ortega–More, L.M.; Abbassi, H. and Peters, J.E. (1999): A review of the importance of Cryptosporidiosis in farm animals. int. J. parasitol., 29: 1269–1287.
- Deshpande, P.D. and Shostri, U.V. (1981): Incidence of *Giardia* infection in calves in Mahashira state, India Trop. Anim. Health prod., 13: 43.
- El. Dessouky, S.A. and Nabila, M.E. (2005): Effect of *Cryptosporidium parvum* infection on the haematological and blood biochemical changes of buffalo calves with special reference to prevalence of infection among buffaloes. Assiut Vet. Med. J., 51: 108-123.
- El. Kelesh, E.A. Abdel-Maogood, S.Z. and Abdel-Wahab, A.M. (2009): Comparative studies on some *cryptosporidium* species infecting different animals. EVMS PJ, 5: 37-47.
- El Khodery, S.A. and Osman, S.A. (2008): Cryptosporidiosis in buffalo calves (Bubaus–bubalis): Prevalence and potential risk factors. Top. Ainm. Hlth. pred., 40(6): 419-426.
- El Shemy, T.M. (2006): Recent advances in diagnosis and treatment of neonatal calf diarrhoea. PhD thesis, Fac. Vet. Med., Alex. Univ.
- El Sherif, A.M.; Abed ElGawad, M.A.A.; Lotfy, H.S. and Shokier, K.A.M. (2000): Impact of gastrointestinal nematodes and some enteric protozoal affections on the health of buffalo calves. Assiut Vet. Med. J., 43(86): 260-270.
- Fitzgerald, P.R. (1980): The economic impact of coccidiosis in domestic animals. Adv. Vet. Sci Comp. Med., 24: 121–143.
- Gasmir, G.S.; Osman, A.Y.; El–Amin, F.A. and Halima, L.O. (2011): A survey of enteric *Coccidia* of cattle in the sudan. Sudan J. Vet. Res., 15: 43–48.
- Goz, Y.; Atug, N.; Yuksek, N. and Ozkan, C. (2006): Parasites detected in neo natal and young calves with diarrhea Bull. Vet. Inst. Pulawy, 50: 345–348.
- Gul, A.; Cicek, M. and Kilinic, O. (2008): Prevalence of *Eimeria* spp., *Cryptosporidium* spp. and *Giardia* spp. in calves in the Van province. Trukiye Parazitol Derg., 3: 202–204.
- Hamnes, I.S.; Gjerde, B. and Robertson, L. (2006): Prevalance of *Giardia* and *Cryptosporidium* in dairy calves in three areas of Norway. Vet. Parasitol., 140: 204–216.
- Harfoush, A.M.; Soliman, A.H. and Amer, A.S. (2005): Some studeies on coccidiosis in calves and lambs in Kafr El–Sheikn Governorate. EVMS PJ, 2: 250. 261.
- Heat, S.E. (1992): Neonatal diarrhea in calves: diagnosis and investigations in problems herds. Compend contin. Educ. Pract. Vet., 14: 995-1002.
- Henriksen, S.A. and Pohlenz, J.F.L. (1981): Staining of cryptosporidia by a modified Ziehl Neelsen technique. Acta.Vet. Scand., 22: 594-596.
- Hompson, R.C. (2000): Giardiasis as are – emerging infections disease and its zoonotic potential. Int. J. Parasitol., 30: 1259–1267.
- Iburg, T.; Gasser, R.B. and Henriksen, S.A. (1996): First record of *Giardia* in cattle in Denmark. ActaVest Scand, 77: 337–341.
- Lassen, B. (2011): The prevalences of *Eimeria* and *Cryptosporidium* in large Latvian cattle Herds. Vet. Med. Zoot., 54: 47–52.
- Lefay, D.; Naciri, M.; Pairier, P. and Chermette, R. (2001): Prevalence of *cryptosporidium* infection in calves in France. Vet. Parasitol, 8: 1-9.
- Levine, N.D. (1985): Veterinary Protozoology. Iowa StateUniversity Press. Ames p., 139.
- Maddox–Hyttel, C.; Langkjaer, R.B.; Enemark, H.L. and Vigre, H. (2006): *Cryptosporidium* and

- Giardia* in different age groups of Danish cattle and pigs: occurrence and management associated risk factors. *Vet. Parasitol.*, 141: 48–59.
- Majeed, H.A.; Maha, K.A.; Nadra Elwgoud, M.I.; El-Azazy, M.E.; Sami, M.A. and El Said, H. (2011): Infectious causes of neonatal diarrhea in cattle in Kuwait with special refernce to cryptosporidiosis. *J. of Anim and Vet. Advances.*, 17: 2282–2286.
- Mary, B. (2003): Anticoccidial effects of clindamycin and chloramphenicol on bovine *Eimeria*. *EVMSPJ*, 1: 13-27.
- Matjila, P.T. and Penzhorn, B.L. (2002): Occurance and diversity bovine Coccidian at three localities in South Africa. *Vet. Parasitol*, 2: 93–102.
- Miron, D. (1991): Calves a source of an out break of cryptosporidiosis among young children in an agricultural close community pediatric *Infect. Dis. J.*, 10: 438–441.
- Misic, Z.; Katic, R.S. and Kulisic (2006): *Cryptosporidium* infection In lambs and goats in Serbia. *Acta veterinaria*, 56, (1): 49-54.
- O'Handley, R.M.; Cockwill, C.; Mcallister, T.A.; Jelinski, M.D.; Morck, D.W. and Olson, M.E. (1999): Duration of naturally acquired giardiosis and cryptosporidiosis in dairy calves and their association with diarrhea. *J. Am. Vet. Med. Assoc.*, 214: 391-396.
- Olson, M.E.; O'Handley, R.M.; Ralston, B.J.; McAllister, T.A. and Thompson, R.C. (2004): Update on *Cryptosporidium* and *Giardia* infections in cattle. *Trends parasitol.*, 20: 185–191.
- Pancieria, R.J.; Thomassen, R.W. and Garner, F.M. (1971): Cryptosporidial infection in a calf. *Vet Pathol.*, 8: 479–484.
- Priti, M.; Sinha, S.R.; Sucheta, S.; Verma, S.B.; Sharma, S.K. and Mandal, K.G. (2008): Prevalence of bovine coccidiosis at Patna. *J. Vet. Parasitol* 22: 5–12.
- Ruest, N.; Faubert, G.M. and Couture, Y. (1998): Prevalence and geographic distribution of *Giardia* spp and *Cryptosporidium* spp in dairy farms In Quebec. *can. Vet. J.*, 39: 697–700.
- Sanchez, R.O.; Romero, J.R. and Founroge, R.O. (2008): Dynamics of *Eimeria* oocyst excretion in dairy calves in the province of Buenos Aire (Argentina), during their first 2 months of age. *Vet. Parasitol.*, 151: 133–138.
- Soulsby, J.L. (1982): *Helminthes, Arthropods and protozoa of domesticated animals* 6th ed. published by Balliere Tindall and cassel L T.d London.
- Suvillan, P.S.; Dupont, H.I.; Arafat, R.R.; Thorton, S.A; Selwyn, B.J.; Elalamy, M.A. and Zaki, A.M. (1988): Illness and reservoirs associated with *Giardia lamblia* infection in rural Egypt: the case against treatment in devolping world environment *Am. J. Epidem.*, 127: 1272-1281.
- Xiao, L. (1994): *Giardia* infection in farm animals. *Parasitol.*, Today, 10: 436–438.
- Xiao, L. and Herd, R.P. (2004): Infection pattern on *Cryptosporidium* and *Giardia* in calves. *Vet. Parasitol.*, 55: 257–262.
- Yu, K.S.; Gao, M.; Huang, N.; Jia, O.Y. and Line, Q. (2011): Prevalence of Coccidial infection in cattle in Shaanxi Province, NorthWestern China. *J. Anim. Vet. Adv*, 20: 2716–2719.

بعض الاسباب الطفيلية للاسهال في العجول بمحافظة البحيرة

رضا سمير أحمد فضلي

اجريت هذه الدراسة على ٣٠٠ عينة من البراز من العجول البقرى المصابة بالاسهال تراوحت أعمارها من أسبوع حتى ٢٠ أسبوع في محافظة البحيرة خلال الفترة من مايو الى ٢٠١١ أبريل ٢٠١٢ للوقوف على نسبة الاصابة بالأوليات وكانت نسبة الإصابة الكلية ٥٢,٦٦%. بينما حويصلات الأيميريا وحويصلات الكريبتوسبورديا والجيارديا (٢٧% و ١٤,٦٦% و ١١%) وذلك باستخدام طريقة التعويم باستخدام محلول ملح مشبع لعزل حويصلات الأيميريا و باستخدام محلول سكر مشبع لعزل حويصلات الجياريا وصبغة الزيل نيلسون المحورة لحويصلات الكريبتوسبورديا وتم تصنيف خمسة أنواع من الحويصلات البوغية لايميريا العجول وهي ايميريا (بوفس -زيرنى- صيبسريكا - سيلندريكا- البيسويدالس) باستخدام ٢.٥% محلول النوتاسيوم داي كرومات وقد وجد ان اكثر الأنواع شيوعا هما ايميريا (بوفس -زيرنى) بنسبة ٢٧,١% و ٢٣,٤% يليها صيبسريكا بنسبة ١٤,٨% ثم البيسويدالس بنسبة ١٣,٥% ، أخيرا سيلندريكا بنسبة ٩,٨%. حويصلات الأيميريا وجدت في كل الفئات العمرية للحيوانات وكان معدل الإصابة أعلى في الفئة العمرية من (١٦- ٢٠) اسبوع بنسبة ٤٦,٦% بينما حويصلات الكريبتوسبورديا لوحظ انها تقل كلما زاد العمر وكان معدل الإصابة أعلى في الفئة العمرية من (١- ٤) اسبوع بنسبة ٣٢% . وأخيرا حويصلات الجيارديا أعلى في الفئة العمرية من (٤- ٨) اسبوع بنسبة ٣٠% وان معدل الإصابة (وحويصلات الأيميريا وحويصلات الكريبتوسبورديا و الجيارديا) كان أعلى في فصل الشتاء مقارنة ببقية فصول السنة.