

**BIOACTIVITY OF LEMON EXTRACTS ON THE COTTON  
LEAFWORM, *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae)  
I- ANTIFEEDANT EFFECT OF LEMON LEAF EXTRACTS  
AGAINST *S.littoralis* larvae.**

**Ahmed A.M. Etman and Ibrahim H.H. Ali  
Department of Plant Protection, El-Fayoum Faculty of Agriculture,  
University of Fayoum, Egypt**

**ABSTRACT**

Five lemon leaf extracts were tested as antifeedants for the cotton leafworm, *Spodoptera littoralis* (Boisduval), where Chloroform, methanol, carbontetrachloride, ether and n-hexane were used. In choice tests; where larvae were offered treated and un-treated cotton leaf discs, extracts of the three latter solvents were very effective on 3<sup>rd</sup> instar larvae. The chloroform extract was effective, while the methanol extract had no such effect. In non-choice tests; where larvae were offered only treated cotton leaf discs, more than 97.0% feeding inhibition was observed with chloroform (weakly polar); 98.6, 83.0 and 95.7% for carbontetrachloride, ether, and n-hexane, (non-polar), respectively.

These results indicate that *Citrus limon* L. leaves contain chemicals that serve as antifeedants for *S.littoralis* larvae

**Key words:** *Spodoptera littoralis*, lemon extracts and Antifeedant effect.

**INTRODUCTION**

Lemon trees, *Citrus limon* L., are not known as host plants of the cotton leafworm, *Spodoptera littoralis* (Boisduval). They are well known as more resistant plants to citrus pests than any other citrus plants. On the other hand, several authors used fine-chopped lemon, or lemon juice in poison baits for the control of many insects such as locusts, grass-hoppers, the corn earworm, *Heliothis armigera* (Hubner), the fall armyworm, *S.frugiperda*, e.g., **Swenk (1918), Davis (1919), and Smith (1921). Roberto et al. (2002)** used limonoids obtained from *C. limon* seeds as antifeedants against larvae of *S. frugiperda*.

Moreover, various parts of citrus fruit trees, shrubs, are sources of citronella oil (**Bagdat, 2006 and Nhu-Trang et al., 2006**), which is well known to be a repellent to some insects such as mosquitoes. **Attaway et al. (1966)** identified 19 compounds in the leaf oil of *C. limon*, among them were some insect repellents.

Several authors used various extracts of different plants with low insect associations for antifeedant activity tests against many insect pests, e.g.; **Etman (1974)** for the native bud-worm of Australia, *Heliothis punctigera* Wallg., and the cluster caterpillar, *S.litura* (F.); **Widson et al. (1983)** for the corn earworm, *H.zea* (Boddie); **Abdel-Rahman and Al-Mozini (2007); Ballesta-Acosta et al. (2008)** for *S.littoralis* larvae.

Other authors, in intensive studies by using plant isolated chemical compounds for antifeeding activity tests, e.g.; **Kubo et al. (1976)** for the armyworms, *S.exempta*, and *S.littoralis*; **Bringmann et al. (1992), and Manna and Attia (1992)** for *S.littoralis*.

The objective of this investigation was to determine the feeding responses of the larvae of *S.littoralis*, to lemon leave extracts, and to assess the probability of the presence of insect antifeedants in lemon leaves.

## MATERIALS AND METHODS

### 1- Insect culture:

Insects used in this study were obtained from laboratory colonies maintained in our Department, and reared on artificial medium by established procedures (Etman *et al.*, 1983). Rearing took place in a controlled environment room maintained at  $28 \pm 2$  °C, with a natural light supplemented by artificial light for a 14 h photo phase, and natural dusk period.

### 2- Plant extraction:

Solvents used were methanol as a polar solvent; n-hexane, carbontetrachloride, and ether represent the non-polar solvents; and chloroform as a weakly polar solvent. Lemon leaves were collected fresh from 12-year-old trees, and dropped in the boiled solvent used to inhibit enzymatic activities. The lemon leaves were cut into small pieces and then minced with a food mincer, and thereafter the material (100 g) was homogenized in a Waring blender with 200 ml solvent. Solvents were removed by using a rotary evaporator at a temperature below 40 °C.

All plant crude extracts were kept below 0 °C. The crude extracts were bioassayed within a week. The final volume of the acetone extract solution was 25 ml.

### 3- Specific extraction procedures:

#### a- Extraction with a mixture of methanol, chloroform and water:

This method was originally used by Blight and Dyer (1959) for the rapid extraction of lipids from fish. In 1970 it was used by Wayte and Southcott in isolating natural products from algae. The chloroform extract was evaporated and the residue dried with dry nitrogen. The residue was dissolved in acetone and transferred to a volumetric flask. The aqueous methanolic layer was filtered, and the extract evaporated at 50 °C to remove the methanol. Then the aqueous solution left was freeze dried. The residue dissolved in distilled water and the volume adjusted to 25 ml with acetone.

#### b- Carbontetrachloride extract:

The minced lemon leaves sample was extracted with a mixture of 250 ml carbon tetrachloride and 250 ml propanol 2, and 150 ml distilled water. The carbon tetrachloride phase was filtered, the solvent evaporated, and the residue dissolved in 25 ml acetone. The aqueous propanol solution was discarded.

#### c- Ether extract by Soxhlet extraction apparatus:

The plant sample was minced, 100 ml of water was added, and the plant material was freeze-dried. The dry residue was scrapped out and placed in the apparatus, which was left overnight until complete extraction was effected. The ether was evaporated, and the residue dissolved in acetone.

#### d- Extraction with n-hexane:

The plant material was blended with 500 ml n-hexane. The aqueous phase was discarded, the n-hexane phase filtered, the solvent evaporated, and the residue dissolved in acetone.

### 3- Bioassay for antifeeding activity:

The leaf-disc test used by Munakata (1970) was used in testing the lemon extracts for active principles with *S.littoralis*. Since 100 g of the lemon leaves were extracted and stored in 25 ml acetone, therefore the

**BIOACTIVITY OF LEMON EXTRACTS ON THE COTTON..... 49**

concentration of the acetone solution of the extract was 4 g equivalent of fresh leaves per one ml acetone.

Round leaf-discs, 21 mm diameter, were punched out with a cork-borer, from the leaves of cotton plants. Treated (T) discs were immersed in the acetone solution of the lemon leaf extract, and the control (C) discs in pure acetone. After air drying, these discs were placed symmetrically in a polyethylene dish.

The larvae used for the experiments were removed from the rearing medium and placed onto cotton leaves for one hour to enable them to adapt more easily under the test conditions to the new type of food. The larvae then were starved for 2 h before the tests. Then 3<sup>rd</sup>-instar larvae were introduced in the center of the dish, covered, and left for 3 h. The experiments were conducted at 22 ± 2 °C. The "leaf-disk test" was repeated 12 times using 12 dishes for each leaf extract.

The data obtained were statistically analysed and any extract which showed significant results was repeated using two similar leaf-discs in the same dish, hence 3 different arrangements were used: Treated-control (T-C), Treated-Treated (T-T), and Control-Control (C-C). This enabled a study of the larval feeding responses where: (i) a choice between treated and untreated discs was offered, (ii) no choice between treated and untreated discs was offered, i.e., forced eaten feeding test. Each experiment was replicated 12 times. The graph paper method for measuring the leaf area consumed was used (Dethier, 1947).

**RESULTS AND DISCUSSION**

**1- Choice experiments:**

In these experiments, where larvae were given a choice between treated and control discs, Wilcoxon's signed-rank test were applied to treated and control data (actual area consumed) instead of a parametric analysis of variance. This was chose because the data did not appear to be normally distributed. Nevertheless, in the experiments where the observed difference was significant, the student's t-test was applied using the formula:

$$t = \frac{\bar{X} - \mu}{S / \sqrt{n}}$$

where  $\mu$  is 20%  
 $\bar{X}$  is the mean of the feeding ratio (FR)  
 $S$  is the S.D. of FR  
 $N$  is the number of samples (12 in each experiment)

**The feeding ratio was expressed as follows:**

$$\text{Feeding ratio} = \frac{\text{Amount consumed of T disc}}{\text{Amount consumed of C disc}} \times 100$$

When the amounts consumed of the treated discs were less than 20% of those of the control discs, the plant extracts were considered to have strong feeding inhibitory activity. The 20% criterion was chosen because this amount of was thought to be of a possible economic value. The feeding inhibitory effect was expressed as follows:

$$1- \frac{\text{Amount consumed of T discs}}{\text{Amount consumed of C discs}} \times 100$$

Since the tabulated  $t$  value for 11 degrees of freedom at 0.01 significance level is 3.11, and the calculated values for the carbon tetrachloride, ether, and n-hexane extracts are more than 3.11, it is evident that the feeding ratios in these experiments (means being 6.5%, 9.8%, and 10.5% for the three extracts, respectively) are significantly less than 20% at 1% level of probability. These highly significant differences and the results suggest that the above plant extracts have very strong feeding inhibitory activity against 3<sup>rd</sup> instar larvae of *S.littoralis* (table 1).

Since the calculated value of " $t$ " for the chloroform extracts is less than the tabulated value for 11 degrees of freedom at 5% level, there is no significant difference between the mean feeding ratio of this leaf extract (15.8%) and the 20% feeding ratio level. These results suggest that this extract has a strong feeding inhibitory activity against 3<sup>rd</sup> instar larvae of *S.littoralis*.

On the contrary, the methanol extract had no feeding inhibitory activity. This was demonstrated by the "not significant" results of Wilcoxon's signed-ranks test and the high feeding ratio mean (80.9%).

**Table 1. Feeding responses of third-instar larvae of *S.littoralis* to cotton leaf-discs treated with lemon leaf extracts, and to alternative choices (controls) processed with acetone only Each extract was replicated 12 times, with 1 dish/ replicate and 10 larvae/ dish.**

Solvent	Mean % area consumed ± S.E.		Mean feeding ratio ± S.E.	Mean feeding inhibitory effect ± S.E.	Wilcoxon's test	Student's t-test Value (11 DF)
	T	C				
Chloroform	1.7 ± 0.6	10.9 ± 1.3	15.8 ± 5.0	84.2 ± 5.0	S**	1.62
Methanol	7.5 ± 1.3	10.9 ± 1.4	80.9 ± 15.3	- -	N.S.	- -
Carbon-tet.	0.6 ± 0.2	10.5 ± 0.9	6.5 ± 1.9	93.5 ± 1.9	S**	6.15**
Ether	1.4 ± 0.5	17.0 ± 1.3	9.8 ± 3.3	90.2 ± 3.3	S**	4.32**
n-Hexane	0.7 ± 0.3	6.7 ± 1.0	10.5 ± 4.5	89.5 ± 4.5	S**	3.28**

S\*\* = Significant at 1 % level of confidence.

N.S. = Not significant

C = Untreated control leaf disc

T = Treated leaf disc

Concentration used = 4 g equivalent of fresh leaf/1 ml

## 2-No choice experiments;

In order to study the larval feeding responses where no choice between treated and untreated discs were offered, the non-parametric Wilcoxon two-sample test was applied to the feeding ratios of the T-C divisions and the feeding ratios of the random pairs of sums of T-T and C-C divisions of the experiments.

Table 2

The results of the analysis showed that there was a significant difference between the feeding responses of 3<sup>rd</sup> instar larvae of *S.littoralis* to chloroform lemon leaves extract where a choice was offered (0.0%), and their feeding responses where no choice was offered (2.7%). In this case larvae ate more of the treated discs where no choice was offered. Similarly, larvae ate more of the treated discs of the carbontet. extract, but the difference was not significant, (the mean feeding ratios were 0.0 and 1.4% for choice and no choice tests, respectively (table 2).

In the other two extracts i.e. ether and n-hexane, there were no significant differences between the feeding responses of 3<sup>rd</sup> *S.littoralis* larvae where a choice or no choice was offered. The mean feeding ratios were 5.2% and 17.0% for ether extract; 17.5% and 4.3% for n-hexane, for choice and no choice tests, respectively (table 2). In other words, in the no-choice tests more than 97% protection was offered by the two weakly polar extract used i.e. chloroform whereas 98.6, 83.0 and 95.7% protection by the non-polar extracts i.e. carbon tet., ether and n-hexane, respectively.

These results suggest that the above lemon leaf extracts have very strong feeding inhibitory activity against 3<sup>rd</sup> instar larvae of *S.littoralis*. The above results have been confirmed by applying Kruskal-Wallis test, which gave the same results as did Wilcoxon's signed-ranks test, approximately.

**Etman (1974)** reported the presence of two aldehydes, an alkaloid, a glycoside and an organic acid in lemon leaf extracts. **Attaway et al. (1966) and Nhu-Trng et al. (2006)** analysed the lemon leaf-oil and found that they contain citronellal, which is a mixture of stereo-isomeric aldehydes (**Merek, 1960**). **Kamiyama (19670)** stated that trans-2-hexenal (a leaf-aldehyde) is a compound of the leaf-oil of rough lemon. In the same analogy, Ruberto et al. (2002) confirmed that natural limonoids obtained from seeds of Citrus limon, i.e. limonin and obacunone showed highly significant antifeedant activity against larvae of *S.frugiperda*; while Chang **MingChow et al. (2004)** reported the presence of ascorbic acid in the lemon leaf methanol extract.

The inhibitory activity observed with *S.littoralis* 3<sup>rd</sup> instar larvae here may stem from the aldehydes, alkaloids, glycosides, and the organic acid compounds together, or it may be due to the action of some of these compounds individually.

## REFERENCES

- Abdel-Rahman, H.R. and Al-Mozini, R.N. (2007):** Antifeedant and toxic activity of some plant extracts against larvae of the cotton leafworm, *Spodoptera littoralis* (Lepidoptera: Noctuidae). Pakistan Journal of Biological Sciences, 10: 4467-4472.
- Attaway, J.A.; Pieringer, A.P.; Barbas, L.J. (1966):** Origin of citrus flavour components. I. Analysis of citrus leaf oils using gas-liquid chromatography, thin L.C. and mass spectrometry. Phytochemistry, 5: 141-151.
- Bagdat, R.B. (2006):** The essential oil of lemon balm, *Melissa officinalis* L., its components and using fields. Ondokuz Mays University, Ziraat Fakultesi Dergisi, Turkey, 21:116-121.
- Ballesta-Acosta, M.C.; Pascual-Villalobos, M.J.; Rodriguez, B. (2008):** The antifeedant activity of natural plant products towards the larvae of *Spodoptera littoralis*. Spanish Journal of Agricultural Research, 6: 851-891.

**BIOACTIVITY OF LEMON EXTRACTS ON THE COTTON..... 53**

- Blight, F.G. and Dyer, W.J. (1959):** A rapid method of total lipid extraction and purification. *Can. J. Biochem. and Physiol.* 37: 911-917.
- Bringmann, G.; Gramatzki, S.; Grimm, C.; and Proksch, P. (1992):** Feeding deterrence and growth retarding activity of the naphthylisoquinoline alkaloid dioncophyline A, against *Spodoptera littoralis*. *Phytochemistry*, 31: 3821-3825.
- Davis, J.J. (1919):** Grasshopper control in India. Purdue Uni. Agric. Expt. Sta., Lafayette, Ind., Circ. no. 88, 8 pp. (c.f. Review of applied entomology, RAE).
- Dethier, V.G. (1947):** Evaluation of feeding preferences. Un "Chemical Insect Attractants and Repellents", pp. 249-260, Blakiston Co., Philadelphia.
- Etman, A. A. M. (1974):** Larval feeding responses of the cluster caterpillar, *Spodoptera litura* (F) and the native budworm of Australia, *Heliothis punctigera* Wallgn., to certain extracts of host and non-host plants. M.Sc. thesis, Faculty of Biological Sciences, University of New South Wales, Australia.
- Etman, A.A.M.; El-Hemaesy, A.H.; El-Naggar, M.Z. (1983):** *Spodoptera littoralis*: Laboratory techniques for rearing and experimenting. Proc. 5<sup>th</sup> Arab Pesticide Conf. Tanta Univ. Vol. (IV): 11-20.
- Kamiyama, S. (1967):** Studies on the leaf oils of citrus species. Part I. Composition of leaf oils from *Citrus unshiu*, *Citrus natsudaidai*, *Citrus kokitsu*, and *Citrus limon*. *Agr. Biol. Chem.* 31: 1091-1096.
- Kubo, I.; Lee, Y.W.; Pettei, M.; Pilkiewicz, F.; Nakanishi, K. (1976):** Potent armyworm antifeedants from the East African *Warburgia* plants. *J. Chem. Soc., Chem. Commun.*, 24:1013-1014.
- Mannaa, S.H. and Attia, A. A. (1992):** Insecticidal properties of some lignin compounds, extracted from the root bark of *Artemisia argentea* L'her, on the cotton leafworm, *Spodoptera littoralis* (Boisd.). *Assiut Journal of Agricultural Sciences*: 119-139.
- Merck, index of chemicals and drugs (1960):** Citral, citronellal, and citronellol, 7<sup>th</sup> edition: 265-267.
- MingChow, C.; TzuChing, W.; PiChuan, L.; MingChan, W.; HungMing, C. and JiunnWang, L. (2004):** Studies on antioxidative activity of methanol extracts from lemon (*Citrus limon* Buram.) leaves. *J. of the Chinese Society for Horticultural Science*. 50: 185-198.
- Munakata, K. (1970):** Insect antifeedants in plants, In "Control of insect behaviour by natural products", edited by: Wood, D.L. *et al.*: 180-187.
- Nhu-Trang, T.; Casabianca, H.; and Grenier-Loustalot, M.F. (2006):** Authenticity control of essential oil containing citronellal and citral by chiral and stable-isotope gas-chromatographic analysis. *J. Analytical and bioanalytical chemistry*, 36: 2141-2152.
- Ruberto, G.; Renda, A.; Tringali, C.; Napoli, E.M. and Simmonds, M.S.J. (2002):** Citrus limonoids and their Semisynthetic derivatives as antifeedant agents against *Spodoptera frugiperda* larvae. A structure-activity relationship study. *J. Agric. Food Chem.*, 50: 6766-6774.
- Shonouda, M.; Osman, S.; Salama, O. and Ayoub, A. (2008):** Toxic effect of *Peganum harmala* L. leaves on the cotton leafworm, *Spodoptera littoralis* Boisd. and its parasitoids, *Microplitis rufiventris* Kok. *Pakistan Journal of Biological Sciences*, 11: 546-552.

- Smith, R.C. (1921):** Observations on the fall armyworm, *Laphygma frugiperla* Smith & Abbott and some control experiments. *J.econ.Ent.*, 14: 300-305. (c.f. RAE).
- Swenk, M.H. (1918):** Grasshopper control. *Nebraska Coll. Agric. Lincoln Emergency Bull.* no. 17, 4 pp. (c.f. RAE).
- Whyte, J.N.C.; Southcott, B.A. (1970):** An extraction procedure for plants: Extracts from the red alga, *Rhodomela larix*. *Phytochemistry*, 9: 1159-1161.

### استخدام مستخلصات أوراق نبات الليمون كمواد مانعة لتغذية يرقات دودة ورق القطن

أحمد عزمي عثمان و إبراهيم حامد حسين على  
قسم وقاية النبات - كلية الزراعة- جامعة الفيوم- مصر

تم اختبار خمسة مستخلصات لأوراق الليمون ضد العمر الثالث ليرقات دودة ورق القطن وهي مستخلصات الكلوروفورم، الكحول الميثيلي، رابع كلوريد الكربون، الإثير، وان-هكسان، لدراسة مدى تأثيرها كمستخلصات تحتوي على مركبات لمواد مانعة للتغذية، وقد أوضحت تجارب التغذية الاختيارية أن المستخلصات الخام لرابع كلوريد الكربون والإثير وان-هكسان لها خصائص مانعة للتغذية قوية جدا ضد يرقات العمر الثالث التي غذيت على أوراق قطن معاملة بتلك المستخلصات. وكذلك أظهرت التجارب أن مستخلص الكلوروفورم له خاصية مانعة قوية. في حين أن مستخلص الكحول الميثيلي لم يكن له أية خواص مانعة للتغذية.

وعندما أعيدت التجارب باستخدام طريقة التغذية الإيجابية التي لم تعطى فيها اليرقات خيارات تفضيلية بين أوراق نبات القطن المعاملة والغير معاملة فإنه تم الحصول على 97.3% وقاية لأوراق القطن المعاملة بمستخلص الكلوروفورم في حين أن مستخلصات رابع كلوريد الكربون والأثير وان-هكسان أعطت 98.6% و 83% و 95.7% وقاية على الترتيب.

وقد أدت هذه النتائج إلى الاستنتاج بأن مستخلصات أوراق الليمون بالمذيبات غير القطبية تحتوي على مواد تمنع تغذية العمر الثالث ليرقات دودة القطن وبدرجة قوية جدا.