

## Analysis of Well Log Data for the Evaluation of Gas Sand Reservoir of Alam El-Bueib (AEB) Formation, 3G Member, Jade-8 Well, Jade Field, Matruh Basin, Egypt

Basal, A.M.K<sup>1</sup> and El-Twargy, E.A.\*<sup>1</sup>

<sup>1</sup>Geology Department, Faculty of Science, Damietta University, New Damietta, Egypt

Received: 21 February 2018 /Accepted: 12 March 2018

\* Corresponding author: esraaeltwargy@du.edu.eg

---

### Abstract

The evaluation of gas reservoirs in Alam El Bueib has been dealt with through the interpretation of well log data. Alam El – Bueib (AEB) formation is monitored in Matruh basin at the Northern part of the Western Desert, Egypt encountered many gas producing reservoirs. Alam El- Bueib – 3G (AEB – 3G) member was chosen for application of the presented techniques. The available well log data are in the form of Gamma Ray, Bit size, Caliper, Resistivity, Neutron, Density, Sonic and Photoelectric. The reservoir petrophysical parameters ( $\Phi_{\text{eff}}$ ,  $S_w$ ,  $B_{vw}$ ) besides mineralogical make up were evaluated for this member in JADE – 8 well. The gas zone represents a challenge for well log interpretation technique as they normally plotted above the standard porosity lines. A graphical technique was used to get  $\Phi_{\text{eff}}$  in the zones on the  $\rho_b - \Phi_N$  cross plot. The  $S_w$  Vs  $\Phi$  was presented graphically on log – log plot for obtaining BVW constant lines. Line with slope unity discriminates between reservoir and non-reservoir for AEB – 3G member.


**Keywords:** Gas sand reservoir, Alam El-Bueib, 3G member, Matruh Basin, Well logging, Western Desert, Egypt.

---

### Introduction

The Western Desert comprises the area westwards from the Nile Valley and Delta and extended to the Libyan borders. The petroleum province of the Western Desert has a major hydrocarbon potentiality as recent discoveries indicated (Abu El Naga 1984); six different Mesozoic basins exist in the northern Western Desert; namely Matruh, Shoushan, Alamein, Qattara, Abu El Gharadig, and Kattaniya (Fig.1). Matruh Basin lies in the North Western part of the Western Desert and has a NNE-SSW direction inherited from reactivated Palaeozoic fabric (Moustafa *et al.*, 2002). The basin history shows an early rifting phase during the Jurassic and Early

Cretaceous when excellent Jurassic (Khatatba formation) and Lower Cretaceous (Alam El Bueib) source rocks were deposited with total organic carbon (TOC) values up to 10% and 5% respectively (Moustafa 2008). This article is focused on evaluation of Alam El Bueib 3G member (AEB – 3G) gas producing reservoir, Jade field. This field is located in the north Western Desert of Egypt, about 50 km south of Matruh City. It lies between latitudes  $31^{\circ} 56' 24''$  to  $31^{\circ} 57' 36''$  N and longitudes  $25^{\circ} 24' 20''$  to  $25^{\circ} 25' 45''$  E (Fig.1).




**Fig.1:** Location map showing the sedimentary basins in the Western Desert of Egypt (EGPC, 1992) and the study well (Jade-8) in Jade field, Matruh Basin, North Western Desert, Egypt.

*Stratigraphy of the Study area:*


The stratigraphic column of the Northern Western Desert comprises a sedimentary succession from Ordovician to recent unconformable overlying Pre-Cambrian basement rocks (EGPC 1992, Fig.2). This succession is subdivided into four major regressive cycles; each one is terminated by a marine transgression. Alam El-Bueib Formation is subdivided into six units (1 to 6). Unit three is further subdivided into seven members from (A to G) (EGPC 1992). This formation belongs to the second cycle which began at Early Cretaceous with the deposition of (units 6 and unit 5) shallow marine clastics at the base followed by (unit 4) with marine shale and (unit 3) which includes a succession of massive fluvial sand bodies separated by marine shale incursions. These sand bodies are overlain by (units 2 and 1) alternating shale, sandstone, and marine shelf carbonates and ends in the Alamein Dolomite. Dahab Shale marks the end of this cycle (Sultan and Halim 1988). The main gas producing reservoir is Alam El Bueib, 3G member (AEB – 3G). It is mainly clastics varying from fine to coarse grain sandstones.

*Structural features of the study area:*

The most predominant structures in the study area are the NE to ENE folding trend associated with reverse faulting. Most of those faults originated during late cretaceous to early tertiary while the E-W faults exhibit evidence of strike slip movements along the old faults (Sultan and Abd El – Halim, 1988). Bounding faults of Matruh basin orientation indicates NE-SSW trend (Fig.3) (Moustafa 2008).



**Fig.2:** Generalized lithostratigraphic column of North Western Desert, Egypt. (Schlumberger, 1995)



**Fig.3:** Schematic geological cross section along northwestern Egyptian NNW-SSE direction (Guiraud and Bosworth, 1999)

**AIM OF THE STUDY**

The main aims of this investigation can be summarized in the following:

- (i) Locating and identifying Gas zones,
- (ii) Define the reservoir matrix composition,
- (iii) Evaluation of the static reservoir parameters ( $S_w$ ,  $V_{sh}$ ,  $\Phi$ ),
- (iv) Detecting and discrimination between movable and residual hydrocarbons,
- (v) Detecting reservoir performance through evaluating Irreducible Water Saturation ( $S_{Wirr}$ ) by applying BUCKLE technique ( $\phi$  vs  $S_w$ ) Using computer program facilities in the form of Interactive Petrophysics (IP) for enhancing and accelerate computing processes with higher accuracy.

*3- Data of the Study:*

A complete set of wire line well log data of the selected wells (Jade- 8) (Fig.1) were kindly provided by Khalda Petroleum Company (KPC).

These logs are in the form of Gamma Ray (GR), Bit size (BS), Caliper (CALI), Resistivity, Neutron, Density, Sonic and Photoelectric. Enlarge this section by adding some description of the study data, like method of recording, dates, companies ..... You could also write about the use of the logs to study so ....I.e; Gamma ray logs were used to .....etc.

**VISUAL DESCRIPTION OF THE WELL LOG DATA**

The first step before any detailed well log data interpretation is to describe qualitatively, the log curve response to pick which intervals are promising to be potentially hydrocarbon bearing. The well log curve deflections reflects the effect of the contained matrix, shalines? , porosity and fluids. It is of prime importance to integrate and correlate all the curve shapes with standard reference values. The following section represent quick look interpretation for the available well log data provided in LAS format. The files are loaded to the IP program and displayed in analog format for Jade-8 (Fig.4)

*Qualitative interpretation of well logs:*

The following is a detailed qualitative interpretation for the log curve shapes for AEB – 3G member in Jade-8 well:

The clearly positive separation between resistivity curves (RD>> Rs>>RMIL) seen at the top of this member down to 10230 feet indicate the presence of f invasion profile characterizing good porosity and permeability with hydrocarbon. The low GR seen on track 3 (<30 API) reflects the clean nature of the reservoir. Presence of mud cake is evidenced as the caliper reads equal or slightly less than bit size. This mud cake confirm good reservoir quality

The abrupt decrease in all resistivity curves seen at 10230 Ft. when both (RD) and (Rs) track each other and continued decreasing down to the rest of the section. This point mark the Gas – water contact (G.W.C). The clean 100 % wet zone is separated from the above gas one by a relatively thick shale interval (10230-10270, GR > 60 API), Pe ≈2.5) and marked by low GR, PE = 1.98, Neutron – Density separation of about 4 P.u with neutron at the right.

It is very interesting to notice that, at the water zone (no gas effect) the separation between Neutron-Density is reduced to standard sandstone

matrix effect. Hence the same sandstone contained both gas and water; the effective porosity can be picked directly on the neutron scale at the mid-point between the two curves in the water zone. In this case the reservoir porosity is about 15%.

The deep resistivity reads about 0.7 Ωm<sup>2</sup>/m at (10280 – 10360 ft). Note that opposite gas interval RD reads > 200 Ωm<sup>2</sup>/m. (Caliper ≤ Bit size) this confirms the presence of mud cake and the resistivity separation as described above.

The sandstone matrix is expected to be the main lithology for the described AEB -3G as the PE curve (track 5) reads almost (1.98 – 2 B/E) of typical sandstone value (Fig.4).

The gas effect can be easily picked on the (CNCF – ZDNC) separation (track 5) (i.e. neutron and density reads too low values with CNCF on the right relative to ZDNC).

A quick look calculation for the connate water resistivity (Rw) can be conducted through the porosity and resistivity values at 100% water zone at (10280 – 10360ft). The deep resistivity read is (RD = 0.7) which represent Ro (Rt = Ro at Sw = 100%). From Archie (1942) equation:


$$R_o/R_w = a / \Phi^m \dots\dots\dots (1)$$

$$R_o = R_w / \Phi^2$$

While a =1, m = 2,

$$R_w = .7 * .15^2$$

$$R_w = 0.016$$



**Fig.4:** Well Log Data analog format as displayed on the IP program for AEB 3G Member, Jade – 8 well Jade field, North western desert, Egypt

### Graphical Interpretation of Well Log Data

The cross plot techniques are widely used in the field of well log data analysis. It represents a preliminary step for locating different mineral constituents. A stream of cross plots between the log data can be conducted. The technique presented in this article is based essentially on  $\Phi_N$ ,  $\rho_b$ , RT and GR. The following is a full description for the cross plots constructed for (Jade -8) well.

$$(\rho_b - \Phi_N - GR \text{ z plot})$$


Figure (5) represents a cross-plot between density (ZDNC) versus Neutron (CNCF) and Gamma ray presented as Z axis (right colored bar). The plotted points were chosen to represent good hole conditions. Points located North West direction possess low GR value (20 -30 API) reflect gas effect. Some points plotted just below limestone line trends higher densities and very low neutron may reflect presence of heavy minerals. According to Mud Log Description (fig.6) traces of free pyrite is occasionally present. Points with gradual increase of GR values indicate clay minerals.

It is worth mentioned here that this plot solves for two minerals matrix. Accordingly, it cannot be used for solving complex matrix constituents. To facilitate the interpretation the same plot was constructed for filtered points with low to moderate GR and (caliper  $\leq 12.5$ ) (Fig.7) which clearly indicate the pyrite trend which has included in the mineralogical model. Points between sand stone and lime stone possess moderately high GR may represent siltstone rather than carbonate which confirmed through the composite log description.


The  $\rho_b - \Phi_N$  crossplot technique can be extended to give a quick gas porosity correction (Fig.8). For sandstone reservoir we can draw a line parallel to the approximate gas correction line and intersect with the horizontal line at zero limestone porosity. This will be a pivot point to draw intermediate porosity lines (0 – 30 %). On this plot porosities for gas points located above sandstone line can be read directly on this porosity lines. These porosities can be correlated with these obtained from gas corrected porosities shown in (Table1) using:

$$\Phi = \left( \frac{\Phi_N^2 + \Phi_D^2}{2} \right)^{1/2} \dots\dots\dots(2)$$


The most important point here to notice is the very good matching between porosity ranges appear on (Fig.8) with those calculated (Table1). This means that, this graphical technique is valid.




**Fig.5:** compensated Neutron porosity ( $\Phi_N$ ) versus formation Density ( $\rho_b$  gm/cc) for AEB formation 3G Member, JADE-08 well, Jade Field, Matruh basin, North Western Desert, Egypt.



**Fig.6:** Mud log description for AEB 3G Member, Jade – 8 well Jade field, North western desert, Egypt.



**Fig.7:** compensated Neutron porosity ( $\Phi_N$ ) versus formation Density ( $\rho_b$  gm/cc) for clean AEB formation 3G Member, JADE-08 well, Jade Field, Matruh basin, North Western Desert, Egypt.



**Fig.8:** compensated Neutron porosity ( $\Phi_N$ ) versus formation Density ( $\rho_b$  gm/cc) for AEB formation 3G Member, JADE-08 well, Jade Field, Matruh basin, North Western Desert, Egypt.


**Table (1):** Calculated effective porosity (PHIE) for gas zones using Equation 2

DEPTH (FT)	PHIE
10124	0.17
10128	0.16
10132	0.157
10136	0.161
10140	0.162
10144	0.161
10148	0.124
10152	0.165
10160	0.142
10172	0.113
10180	0.102
10184	0.132
10188	0.121
10192	0.121
10196	0.173
10200	0.172
10204	0.146
10208	0.156
10212	0.158
10216	0.182
10220	0.169
10224	0.166
10228	0.164

*M – N plot*

This plot depends on three log parameters ( $\rho_b$ ,  $\Phi_N$  and  $\Delta T$ ) with GR-Z plot and has its advantages over  $\rho_b$  -  $\Phi_N$  as it can solve for more than two mineral constituents. (Fig.9) represents M – N plot for 3 G member of AEB formation for Jade – 8 well which reflect great enhancement for mineralogical model constitution, on this plot calcite is excluded as the majority of points are clustered on quartz. The gas effect is clearly seen as points clustered towards N


– E direction. It is noticed that, few points aligned S- W with low gamma ray readings which may be due to presence of pyrite. This plot confirmed the above mentioned matrix constituents.



**(Fig.9):** M N plot for AEB formation 3G unit at JADE-08 well Jade Field, – Matruh basin, North Western Desert, Egypt.

*Pickett plot:*

Figure (10) represents the Pickett plot for 3G reservoir at JADE-08 well Jade Field. The Gamma ray was presented as “Z” value in a color code. The red line represents 100% Sw (Ro line) with “m” equals 1.88 (slope of the line). The intersection of this line with vertical axis at 1 gives Rw of 0.0186  $\Omega m^2/m$ . It is very interesting to notice that this value is the same as those obtained through qualitative interpretation of 0.016  $\Omega m^2/m$ .



**Fig.10:** Pickett plot for AEB Formation 3G unit at JADE-08 well Jade Field, Matruh basin, North Western Desert, Egypt.


*Porosity-Saturation*

Buckle 1965 introduced the most important value for evaluating the reservoir performance. This is done by computing the product of porosity time the water saturation. Reservoir at irreducible water saturation (i.e  $\Phi \cdot Sw$  constnt) will produce clean oil with no water. Buckle that when plotting porosity against water saturation on linear scale


makes points representing reservoir at irreducible will follow hyperbola with lower values. In this article, logarithmic scales will be used instead. The relation will be as follows (Holmes, etal 2009):

$$\text{Log Swi} = \text{Log C} - \text{Log phi} \dots\dots\dots (3)$$

This can be represented graphically on log-log paper (Fig. 11). Points representing irreducible state will be followed straight line with slope of unity. Intersection of this line with Log phi equals one give the constant C. (Fig. 12) represents application to AEB-3G reservoir at JADE-08 well Jade Field. On this plot, red colored points plotted below the violet line ,with slope unity, reflect free water production. The constant is 0.03 which confirm the sandstone nature of this reservoir. Pointes plotted above this line that line (blue color) represent movable water production.



**Fig.11:** Holmes et.al. (2009) plot for representation of porosity versus water saturation on log-log paper for obtaining the constant “C”.




**Fig.12:** Holmes et.al. (2009) plot for AEB formation 3G unit at JADE-08 well Jade Field, Matruh basin, North Western Desert, Egypt.

**THE LITHOSATURATION CROSS PLOT (Fig.13)**

Through this plot, it is very obvious that the upper most part of AEB – 3G member (10120 – 10230

Ft) represent the main productive zone. It reflect a constant and continuous reservoir parameters with 18% average  $\Phi$  and 18% Sw. the sand stone is the main matrix constituent with some shale streaks. This reservoir can be considered clean. Presences of pyrite traces (violet) are also seen. The majority of the contained gas is movable (red area). The gas zone is separated from the underlying water zone by a relatively high Vsh unit (10230 – 10280Ft). This water zone is extended downward to the total depth. It is important here to refer to the very low BVW characterizing excellent to very good reservoir quality and expected to produce free water gas.



**Fig.13:** The litho-saturation cross-plot for AEB formation 3G unit at JADE-08 well Jade Field, Matruh basin, North Western Desert, Egypt.

**|CONCLUSION**

The presented work shed more light on the importance of the well log interpretation techniques for evaluating the AEB – 3G gas reservoir in Matruh Basin, North Western Desert, Egypt. The following are the main conclusions extracted:

1. The gas effect can be eliminated for the sandstone reservoir through graphical presentation for  $\rho_b$  Vs  $\Phi_N$ .
2. The tri porosity cross plot in the form of M – N greatly enhanced the obtained mineralogical model for AEB – 3G member.

- It includes gas effect, presence of pyrite traces and Quartz minerals.
3. The  $\Phi$  vs Sw presented on log – log paper facilitate discrimination and differentiation between gas and water zone for AEB – 3G member in JADE – 8 well. The plotted points of AEB – 3G reservoir were located below straight line with slope unity indicating that reservoir is at irreducible state with free water production.
  4. The lithosaturtion cross plot give visual and quick method for locating the producing intervals to aid the decision make where to perforate.
  5. The AEB -3G reservoir characterized by high effective porosity (18%), low Sw (<20%) and very low BVW (<0.02).
  6. The obtained results were confirmed through mud log description.

## References

- Archie, G. E. (1942): The electrical resistivity log as an aid in determining some reservoir characteristics: Petroleum Technology, v. 5 p. 54-62.
- Buckles, R.S.(1965): Correlating and averaging connate water saturation data, Journal of Canadian Petroleum Technology, v.9, no.1, p.42-52.
- EGPC. (1992): In: Western Desert, oil and gas fields, a comprehensive overview.11<sup>th</sup> Petroleum Exploration and Production Conference, Egyptian General Petroleum Corporation, Cairo. pp. 1-431.
- Guiraud, R., Bosworth,W.(1999): Phanerozoic geodynamic evolution of north- eastern Africa and the north western Arabian platform. Rev.Tectonophysics 315, 73 108.
- Holmes, M., Holmes, A., Holmes, D.,(2009): Relationship between Porosity and Water Saturation: Methodology to Distinguish Mobile from Capillary Bound Water AAPG Annual Convention, Denver, Colorado, June 7-10, 2009.
- Moustafa, A.R. (2008): Mesozoic Cenozoic basin evolution in the northern Desert of Egypt. In: Salem, M., El-Arnauti, A., Saleh, A. (Eds.), 3rd Symposium Sedimentary Basins of Libya, The Geology of East Libya, vol. 3, pp. 29-46.
- Moustafa, A.R., EL-Barkooky, A.N., Mahmoud, A., Badran, A.M., Helal, M.A., Nour EL din, M. and Fathy, H. (2002): Matruh basin: hydrocarbon plays in an inverted Jurassic- Cretaceous rift basin in the northern Western Desert of Egypt. Am. Assoc. Petrol. Geol. Int. Meeting (Cairo, Oct. 2002), Abstract.
- Schlumberger (1995) In: Geology of Egypt. Paper presented at the Well Evaluation Conference, Schlumberger, Cairo, pp 58–66.
- Shalaby MR, Hakimi MH, Abdullah WH (2011) Geochemical characteristics and hydrocarbon generation modeling of the Jurassic source rocks in the Shoushan Basin, north Western Desert, Egypt. Mar Petrol Geol 28(9):1611–1624.
- Sultan, N. and Halim, M.A. (1988). Tectonic framework of northern Western Desert, Egypt & its effect on hydrocarbon accumulations. Proc. 9<sup>th</sup> E. G.P. C. Petrol. Explor. & Prod. Conf., 2, 1-22.

## المخلص العربي

عنوان البحث: تحليل بيانات سجل الآبار لتقييم مكامن الغاز والرمل لتكوين علم البويب، عضو G<sup>3</sup>، بئر Jade-8، حقل اليشم، حوض مطروح، مصر

أحمد بصل<sup>1</sup>، اسراء التاورجي<sup>1</sup>

<sup>1</sup> قسم الجيولوجيا – كلية العلوم – جامعة دمياط

تتناول هذه الورقة البحثية تقييم خزانات الغاز من خلال تفسير بيانات تسجيلات الآبار لمتكون علم البويب في حوض مطروح شمال الصحراء الغربية، يوجد بمصر العديد من الخزانات المنتجة للغاز. تم اختيار AEB (3G) لتطبيق التقنيات المعروضة. تم تقييم المعلومات البتروفيزيائية للخزان إلى جانب التركيب المعدني في بئر Jade – 08. يمثل تقييم الخزانات الحاملة للغاز تحدياً لتقنيات تسجيلات الآبار حيث أنها ترسم عادةً فوق خطوط المسامية القياسية. تم استخدام تقنية رسومية للحصول على  $\Phi_{eff}$  في منطقة الدراسة باستخدام العلاقة بين  $\Phi_N$  -  $p_b$ . كما حددت العلاقة بين المسامية و مقدار تشبع الخزان بالماء الفرق بين الخزان الحامل للغاز و الغير حامل للغاز.