

RESPONSE OF TWO CULTIVARS OF *Ficus alii* L. TO FOLIAR APPLICATION OF SOME TRACE ELEMENT TREATMENTS

(Received : 21.3. 2014)

By
A. B. El- Attar

Ornamental Horticulture Department, Faculty of Agriculture, Cairo University, Giza, Egypt

ABSTRACT

This study was carried out in the glasshouse of the nursery of Ornamental Horticulture Department, Faculty of Agriculture, Cairo University, Giza, during the two successive years 2012 and 2013. The aim was to investigate the response of two cultivars of *Ficus alii*, cv. "Green" and "Variegata" to some treatments of foliar application of trace elements. The seedlings were sprayed with Zn, Fe and Mn at the rate of 50 mg/l/ 3 weeks, and Borax at the rate 10 mg/l/3 weeks, while the control plants were sprayed with tap water. All the plants were monthly fertilized with life green (NPK, 20-20-20) at the rate of 2 g/pot.

Foliar application treatments of Zn, Fe, Mn and B significantly increased all the vegetative growth parameters as well as the contents of total chlorophylls (a + b), total carbohydrates treatment, N, P, K, Zn, Fe, Mn and B ppm treatment in the leaves as compared with the control plants. On the other hand, trace elements treatment (in most cases) caused significant decrease in carotenoids content in the leaves compared to the untreated ones.

In both seasons, *Ficus alii* cv. "Green" had plant height, leaf area as well as carotenoids content in leaves which were significantly higher than those of cv. "Variegata". On the other hand, the cv. "Variegata" had the number of leaves, stem diameter, fresh and dry weights of shoots and roots as well as contents of total chlorophylls, total carbohydrates percentage, N, Zn, Fe Mn and B contents which were significantly higher than those of cv. "Green".

From the above results, it can be recommended that for the best vegetative growth of *Ficus alii* cvs. "Green" and "Variegata", the plants sprayed with Fe, Zn or Mn at 50 mg/l/ 3 weeks and/or B at 10 mg/l/ 3 weeks and fertilized by 2g/pots/ month of chemical fertilizer (NPK, 20-20-20), should be foliar sprayed

Key words: *Cultivars, Ficus alii, Micronutrients (zinc, iron, manganese, boron).*

1. INTRODUCTION

Ficus alii (*F. binnendijkii* "alii" or, *F. maclellandii* "alii", Family: Moraceae) is an interesting foliage plant, native to Philippines, Thailand and Costa Rica. It is a tree of dense growth and pendant habit which may grow to several meters in height. Its relatively thick leaves are lance-shaped (willow-like leaves), shiny, deep green up to 10" long and tapering to a slender point and leaf width depends on variety. New growth is sometimes tinted bronze and later matures to a deep, rich green. *Ficus alii* tolerates a wide range of light, temperature, and watering conditions without losing its leaves. It lacks the tendency to shed leaves.

In the commercial production of ornamental plants, the applications of both macro nutrients

(NPK) and one or more of the trace elements (micro nutrients) are the most recommended fertilization regime. Trace elements such as zinc (Zn), iron (Fe), manganese (Mn) and boron (B) play a vital role in the various physiological and biochemical processes that occur within the plant and they are necessary for normal plant growth and development. However, the amount needed from these elements are so small (Allen and David, 2007). Functions and mode of action for trace elements were explained by (Marschner, 1995 and Taiz and Zeiger, 2002). Zinc (Zn) is an essential element for plant that acts as a metal component of various enzymes (almost 60 enzymes) or as a functional structural or regulatory cofactor for protein synthesis, photosynthesis, carbohydrate metabolism, the

synthesis of auxin, cell division and maintains of membrane structure. Iron (Fe) involved with chlorophyll synthesis, plays an essential role in plant metabolism and complexes with proteins to form important enzymes in the plant. Manganese (Mn) participates in several important processes including photosynthesis, and metabolism of both nitrogen and carbohydrate (Taiz and Zeiger, 2002). Boron (B) plays an important role in sugar translocation and carbohydrate metabolism, also is involved with cell division, differentiation, growth, and respiration. Foliar feeding is a relatively new and controversial technique of feeding plants by applying liquid fertilizer directly to their leaves (Taiz and Zeiger, 2002). Since the trace elements are needed for plants in small amounts, applying them as foliar spraying at the right time is a correct way to save the plant requirements (Kuepper, 2003). In this respect, the effects of foliar application of trace elements on growth of ornamental and foliage plants have been studied by several investigators who reported improved vegetative growth parameters (including plant height, the number of leaves/plant, leaf area, stem diameter, fresh and dry weights of shoots and roots). In addition increments in pigment contents and total carbohydrates have been also reported due to foliar spraying of trace elements on *Dieffenbachia* (Dorgham 2005), *Philodendron* and *Syngonium* (Mazhar *et al.*, 2006), *Taxodium disticum* (Abd El-Aziz and Balbaa 2007), *Salvia farinacea* (Farahat *et al.* 2007), *Cupressus sempervirens* (Naguib *et al.* 2007), *Codiaeum variegatum* cv. "Gold Star", *Ruta graveolens* (Esmail 2008), *Dracaena marginata* (Heider 2010), *Areca lutescens* (El-Attar 2011). Treatments of Ca, Mg, Fe and Zn significantly increased plant height, stem diameter, leaf area and the number of leaves. However, the available data of published research as regards the effect of trace elements on *Ficus alii* are very limited.

Therefore, this study aimed to investigate the response of two cultivars of *Ficus alii* cv. ("Green" and "Variegata") to foliar application of trace elements on growth and biochemical composition.

2. MATERIALS AND METHODS

This study was carried out in the glasshouse of the Nursery of Ornamental Horticulture Department, Faculty of Agriculture, Cairo University, Giza, Egypt, during the two

successive years 2012 and 2013. The aim of this study was to investigate the effect of foliar application of different trace elements on the growth and chemical constituents of two cultivars of *Ficus alii* cv. "Green" and "Variegata".

On 1st of March, 2012 and 2013 (in the first and second seasons, respectively), uniform seedlings from both cultivars of *Ficus alii* (with an average plant height of 15-18 cm), were planted in 30-cm diameter plastic pots filled with the mixture of clay plus sand plus peat moss (1:1:1 v/v/v). The physical and chemical characteristics of the mixed media are shown in Table (1).

Table (1): Physical and chemical characteristics of the soil media used for growing two cultivars of *Ficus alii* plants.

Treatments	value	Treatments	value
pH	7.62	Course sand	4.3
E.C ds/m	0.8	Fine Sand	30.2
HCO ₃ +CO ₃	1.8	Silt	39.2
Cl ⁻	1.6	Clay	26.3
SO ₄ ⁻	4.6	Texture class	Loamy
Ca ⁺⁺	3.6		
Mg ⁺⁺	0.6		
Na ⁺	2.2		
K ⁺	1.6		

From 21st of March till 31st of October (in both years), the plants were sprayed every 3 weeks with the separated solutions of Zn, Fe and Mn (EDTA-Chelated form 13% , 6% and 13% respectively) at the rate of 50 mg/l. Boron was also used as a foliar spray at the rate of 10 mg/l (borax 11.4%). In addition, the control plants were sprayed with tap water. Bio-film was added at 1 ml /l to each solution as a wetting agent. Foliar application of each treatment was carried out after 21 days from transplanting, using a hand sprayer until run off point.

All plants received the soluble chemical fertilizer [life green (NPK, 20-20-20)], which was applied monthly at the rate of 2 g/pot, other common agricultural practices including regular watering, hand picking of weeds were also preformed.

The layout of the experiment was a complete randomized block design, with 3 blocks (replicates). The study included 10 treatments [5 trace elements treatments (including the control) x 2 cultivars of *Ficus alii*], with each block consisting of 50 plants (5 plants/treatment).

On 21st of November 2012 and 2013 (in both years, respectively), the experiment was terminated and the vegetative growth parameters were recorded, including plant height (cm), the number of leaves/plant, leaf area (cm²), stem diameter cm, at 5cm. above soil surface, as well as fresh and dry weights of shoots and roots/plant (g/ plant). Also, chemical analysis of fresh leaf sample was conducted to determine their total chlorophylls (a + b) and carotenoids contents (mg/g Fresh weight) using the method recommended by Saric *et al.* (1967). In addition, the total carbohydrate content (% of Dry weight) was determined in dried leaves using the colorimetric method as described by Dubois *et al.* (1956). Dried samples were digested to extract nutrients as described by Piper (1947), and the extract was chemically analyzed to determine its contents of nitrogen, phosphorus and potassium (% of Dry weight). Nitrogen percentage was determined using the modified micro-Kjeldahl method as described by Pregl (1945), while phosphorus percentage was estimated using the method recommended by Troug and Meyer (1939). The potassium percentage was determined by using a Jenway flame photometer, according to Chapman and Pratt, (1961). The contents of Fe, Mn, Zn and B (ppm) in the extract were also determined by using Flame Atomic Absorption JENWAY PFP7 with a boiling air-acetylene burner as recommended by (Isaac and Kerber, 1971). The data recorded on vegetative growth characteristics and chemical constituents were statistically analyzed. An analysis of variance (ANOVA) was carried out, and the means of the recorded data were compared using the "Least Significant Difference (L.S.D.)" test at the 5% level, as described by Steel and Torrie (1980).

3. RESULTS AND DISCUSSION

3.1. Vegetative growth characteristics

The results recorded in the two years of the experiments (Tables 2 and 3) indicated that foliar application of the different trace elements treatments (Fe, Zn, Mn and B) had a considerable effect on vegetative growth characteristics such as: plant height, the number of leaves/plant, leaf area, stem diameter, fresh and dry weights of shoots as well as fresh and dry weights of roots. Foliar spraying of trace elements promoted vegetative growth, and resulted in significant increases in the values recorded for the different growth parameters,

compared to the untreated control plants. Similar promotion of vegetative growth as a result of trace elements has been reported by El-Fadaly (1994) on *Jasminum sambac*, Said (1997) on *Codiaeum variegatum* El- Deeb (1999) on *Aspelinium nidus* and *Philodendron erubescens*, Mansour (2002) on *Scindapsus pictus*, Dorgham (2005) on *Dieffenbachia*, *Philodendron* and *Syngonium*, Gomaa and Ragab (2005) on *Codiaeum variegatum*, Al-Hadad (2006) on *Cupressus macrocarpa*, Mazhar *et al.* (2006) on *Taxodium disticum*, Abd El-Aziz and Balbaa (2007) on *Salvia farinacea*, Farahat *et al.* (2007) on *Cupressus sempervirens*, Naguib *et al.* (2007) on *Ruta graveolens*, El-Bagoury *et al.* (2008) on *Hedera helix*, Esmail (2008) on *Codiaeum variegatum* cv. "Gold Star", Heider (2010) on *Dracaena marginata*, Kumar and Haripriya (2010) on *Nerium odorum*, Eid *et al.* (2010) on *Polianthes tuberosa*, and El- Attar (2011) on *Areca lutescens*.

The favorable effect of the trace elements on the vegetative growth characteristics (compared to the control) can be explained by the important role of these elements in the different physiological and biochemical processes within the plant, which in turn affect the plant growth (Allen and David, 2007).

The data in both seasons (Tables 2 and 3) clearly revealed also that, foliar application of Mn treatments appeared to be the most effective treatment for increasing plant height, the number of leaves/plant, leaf area, fresh and dry weights of shoots as well as fresh and dry weights of roots, compared to the other treatments. On the other hand, the most effective treatment for increasing stem diameter (in both seasons) was foliar spraying of B followed by the application of Mn treatments.

Concerning the cultivar of *Ficus alii*, regardless of the effect of foliar application of trace elements the data in Tables (2) and (3) showed that, in both seasons, the values recorded on plant height and leaf area of the cv. "Green" were significantly higher than those recorded on the same parameters of the cv. "Variegata" as compared to each other. On the other hand, in both seasons the cv. "Variegata" had the number of leaves, stem diameter, fresh and dry weights of shoots as well as fresh and dry weights of roots which were significantly higher than the same values recorded for the cv. "Green".

Regarding the interaction effect between foliar application of trace elements and *Ficus alii*

Table (2): Effect of foliar application of trace elements on plant height, the number of leaves/plant, leaf area and stem diameter of *Ficus alii* cvs. "Green" and "Variegata", during the 2012 and 2013 seasons.

*Trace elements treatments (T)	First season (2012)			Second season (2013)		
	Cultivars (C)		Mean (T)	Cultivars(C)		Mean (T)
	Green	Variegata		Green	Variegata	
Plant height (cm)						
Control	35.27	28.13	31.70	38.32	30.16	34.24
Zn	46.38	32.92	39.65	50.29	33.48	41.89
Fe	44.36	36.20	40.28	54.35	39.36	46.85
Mn	48.31	37.10	42.71	58.36	38.14	48.25
B	45.33	41.16	43.24	47.70	44.10	45.90
Mean (C)	43.93	35.10	-----	49.80	37.05	-----
L.S.D. (0.05)						
T	0.66			0.42		
C	0.38			0.24		
T X C	0.94			0.60		
Number of leaves/plant						
Control	21.08	17.01	19.05	27.50	20.73	24.11
Zn	26.28	35.07	30.67	32.36	39.63	36.00
Fe	24.04	43.11	33.58	33.95	45.50	39.72
Mn	31.54	51.10	41.32	36.27	53.45	44.86
B	25.04	45.61	35.33	28.36	47.16	37.76
Mean (SL)	25.60	38.38	-----	31.69	41.29	-----
L.S.D. (0.05)						
T	1.62			2.50		
C	0.94			1.44		
T X C	2.29			3.54		
Leaf area cm²						
Control	17.67	16.16	16.92	19.50	18.39	18.94
Zn	23.92	21.06	22.49	25.60	22.73	24.16
Fe	23.55	19.80	21.67	26.51	21.90	24.20
Mn	24.87	22.67	23.77	25.71	23.16	24.43
B	21.66	23.68	22.67	24.60	20.86	22.73
Mean (SL)	22.33	20.67	-----	24.38	21.41	-----
L.S.D. (0.05)						
T	0.52			0.40		
C	0.30			0.23		
T X C	0.73			0.54		
Stem diameter (mm)						
Control	0.38	0.42	0.40	0.40	0.45	0.43
Zn	0.46	0.61	0.54	0.47	0.66	0.57
Fe	0.49	0.66	0.58	0.51	0.71	0.61
Mn	0.53	0.72	0.63	0.55	0.79	0.67
B	0.50	0.80	0.65	0.59	0.82	0.71
Mean (SL)	0.47	0.64	-----	0.50	0.69	-----
L.S.D. (0.05)						
T	0.12			0.13		
C	0.09			0.09		
T X C	0.20			0.21		

* Zn, Fe and Mn = 50 mg/l B= 10 mg/l

Table (3): Effect of foliar application of trace elements on shoot and root fresh and dry weights (g/plant) of *Ficus alii* cvs "Green" and "Variegata", during the 2012 and 2013 seasons.

*Trace elements treatments (T)	First season (2012)			Second season (2013)		
	Cultivars (C)		Mean (T)	Cultivars (C)		Mean (T)
	Green	Variegata		Green	Variegata	
Fresh weight of shoots (g/plant)						
Control	6.97	17.01	11.99	8.84	20.73	14.78
Zn	14.87	49.68	32.27	16.87	51.41	34.14
Fe	12.76	65.43	39.10	18.93	70.11	44.52
Mn	16.93	80.36	48.65	23.53	88.04	55.78
B	13.80	66.98	40.39	15.91	72.32	44.12
Mean (C)	13.07	55.89	-----	16.82	60.52	-----
L.S.D. (0.05)						
T	1.33			1.69		
C	0.77			0.97		
T X C	1.88			2.38		
Dry weight of shoots(g/plant)						
Control	1.89	3.75	2.82	2.03	4.95	3.49
Zn	3.71	12.87	8.29	4.87	13.11	8.99
Fe	3.19	14.09	8.64	5.90	16.10	11.00
Mn	5.01	20.00	12.51	7.88	24.54	16.21
B	3.55	14.81	9.18	4.98	17.23	11.11
Mean (SL)	3.47	13.10	-----	5.13	15.19	-----
L.S.D. (0.05)						
T	0.46			0.25		
C	0.27			0.14		
T X C	0.65			0.35		
Fresh weight of roots (g/plant)						
Control	2.52	4.93	3.73	1.88	10.38	6.13
Zn	4.83	18.69	11.76	3.46	12.79	8.13
Fe	6.51	28.57	17.54	2.93	20.89	11.91
Mn	9.83	49.04	29.44	9.53	24.36	16.94
B	3.12	10.84	6.98	2.18	15.93	9.06
Mean (SL)	5.36	22.41	-----	4.00	16.87	-----
L.S.D. (0.05)						
T	0.45			0.82		
C	0.26			0.47		
TX C	0.63			1.16		
Dry weight of roots(g/plant)						
Control	0.86	1.00	0.93	0.68	2.74	1.71
Zn	1.45	4.67	3.06	1.25	3.07	2.16
Fe	1.93	7.14	4.54	1.06	4.61	2.84
Mn	3.19	12.26	7.73	3.00	5.01	4.01
B	0.91	2.50	1.71	0.80	3.57	2.19
Mean (SL)	1.67	5.51	-----	1.36	3.80	-----
L.S.D. (0.05)						
T	0.23			0.43		
C	0.13			0.25		
T X C	0.32			0.61		

* Zn, Fe and Mn = 50 mg/l B= 10 mg/l

cultivars, the data recorded on the vegetative growth (Table 2) showed that in both seasons within both cultivars of *Ficus alii*, foliar application of the different trace elements treatments caused a significant increase in plant height, the number of leaves/plant, leaf area, fresh and dry weights of shoots as well as fresh and dry weights of roots as compared to the control plants. In both seasons, the cv. "Green" foliar sprayed with any of the trace element treatments had stem diameter insignificantly higher than those of the control plants, while application of these different treatments, in most cases, significantly increased stem diameter of the cv. "Variegata". In both seasons, the highest values recorded for most of the growth parameters of both studied cultivars were obtained from plants received Mn treatment, while the highest values of plant height, leaf area and stem diameter (in cv. "Green") resulted from plants sprayed with Mn treatment.

3.2. Chemical constituents

3.2.1. Total chlorophylls (a+b) and carotenoids mg/g F/W

Data presented in Table (4) indicated that foliar application of trace elements was very beneficial in terms of the total chlorophylls content in leaves of *Ficus alii* plants. In both seasons, spraying the plants with any solution of the tested trace elements significantly increased total chlorophyll content in the leaves, compared to the control. Increases in chlorophyll content as a result of the different trace element treatments were similar with the findings of El-Fadaly (1994) on *Jasminum sambac*, Said (1997) on *Codiaeum variegatum*, El – Deeb (1999) on *Aspelinium nidus* and *Philodendron erubescens*, Mansour (2002) on *Hedra helix*, Mazhar *et al.* (2006) on *Taxodium disticum*, Abd El-Aziz and Balbaa (2007) on *Salvia farinacea*, Farahat *et al.* (2007) on *Cupressus sempervirens* and Esmail (2008) on *Codiaeum variegatum* cv. "Gold Star". Kumar and Haripriya (2010) on *Nerium odorum*.

In the case of the carotenoides mg/g FW content, the data recorded in both seasons revealed that, in most cases, application of any trace elements significantly decreased carotenoid content in the leaves compared to the control.

Concerning the cultivars of *Ficus alii*, the data in Table (4) revealed that, in both seasons, the cv. "Variegata" had significantly higher

values of the total chlorophyll contents than the cv. "Green". On the other hand, the carotenoid content in the cv. "Green" was significantly higher than in cv. "Variegata"

Regarding the interaction effect between foliar application of trace elements and *Ficus alii* cultivars, the data in Table (4) showed that in both seasons within both cultivars of *Ficus alii*, foliar application of different trace elements resulted in significant increase in total chlorophyll content compared to the untreated control. While, in both seasons, the lowest values recorded on total chlorophyll content of both cultivars were the untreated control plants. The highest values in the first season (1.09 mg/g FW) were recorded on the cv. "Green" plants sprayed with Fe treatment, while the highest value in the second season (1.11 mg/g fresh weight) was recorded in the cv. "Variegata" sprayed with Mn or B treatment.

3.2.2. Total carbohydrates (% of dry weight)

The data recorded on total carbohydrate percentages (Table 4) revealed that foliar spraying of the different trace elements had a beneficial effect on promoting the synthesis and accumulation of carbohydrate percentages in the dried leaves of *Ficus alii* plants. In both seasons, the values recorded were significantly higher in the plants which received any treatment of the tested trace elements compared to the control. Similar results were reported by El-Deeb (1999) on *Aspelinium nidus* and *Philodendron erubescens*, Al-Hadad (2006) on *Cupressus macrocarpa*, Mazhar *et al.* (2006) on *Taxodium disticum*, Abd El-Aziz and Balbaa (2007) on *Salvia farinacea*, Farahat *et al.* (2007) on *Cupressus sempervirens*, Esmail (2008) on *Codiaeum variegatum* cv. "Gold Star". Kumar and Haripriya (2010) on *Nerium odorum* and Eid *et al.* (2010) on *Polianthes tuberosa*.

Among the different trace elements, in both seasons, the application of Mn or B treatment appeared to be the most effective for increasing carbohydrate percentage, as it gave higher mean values than those recorded with other treatments and the control plants.

The favorable effect of foliar spraying of the different trace elements on the total carbohydrates percentage may be indirectly attributed to the increase in the content of total chlorophylls. As the synthesis of total chlorophylls was promoted, the rate of photosynthesis increased, leading to an increase in carbohydrate synthesis.

Table (4): Effect of foliar application of trace elements treatments on the total chlorophylls (a + b), carotenoids content and total carbohydrates treatment in the leaves of *Ficus alii* cv.s "Green" and "Variegata", during the 2012 and 2013 seasons.

*Trace elements treatments (T)	First season (2012)			Second season (2013)		
	Cultivars (C)		Mean (T)	Cultivars (C)		Mean (T)
	Green	Variegata		Green	Variegata	
Total chlorophylls (a+ b) content (mg/g fresh weight)						
Control	0.47	0.44	0.46	0.43	0.62	0.73
Zn	0.74	1.02	0.89	0.71	0.91	0.81
Fe	1.09	0.91	1.00	0.74	0.91	0.82
Mn	0.73	0.95	0.84	0.98	1.11	1.05
B	0.78	1.07	0.92	0.85	1.11	0.98
Mean (C)	0.76	0.88	----	0.74	0.93	-----
L.S.D. (0.05)						
T	0.09			0.08		
C	0.05			0.04		
T X C	0.13			0.10		
Carotenoids content (mg/g fresh weight)						
Control	0.61	0.35	0.48	0.92	0.45	0.69
Zn	0.33	0.25	0.29	0.63	0.47	0.55
Fe	0.34	0.38	0.36	0.79	0.36	0.58
Mn	0.93	0.41	0.67	0.79	0.32	0.55
B	1.05	0.30	0.68	0.92	0.36	0.64
Mean (SL)	0.65	0.34	-----	0.81	0.39	----
L.S.D. (0.05)						
T	0.05			0.04		
C	0.03			0.02		
T X C	0.08			0.05		
Total carbohydrates (% of dry weight)						
Control	20.00	23.60	21.80	29.00	30.10	29.55
Zn	30.10	31.60	30.85	32.60	31.30	31.95
Fe	29.30	30.50	29.90	26.50	36.20	31.35
Mn	33.02	33.80	33.41	32.53	33.28	32.91
B	30.60	33.20	31.90	38.30	40.30	39.30
Mean (SL)	28.60	30.54	-----	31.79	34.24	-----
L.S.D. (0.05)						
T	0.92			1.29		
C	0.65			0.91		
T X C	1.30			1.82		

* Zn, Fe and Mn = 50 mg/l B= 10 mg/l

As for the cultivars of *Ficus alii*, the data recorded that the total carbohydrates percentage (Table 4) showed that, in both seasons the cv. "Variegata" had significantly higher values than those obtained with the cv. "Green".

Regarding the interaction effect between foliar application of trace elements and *Ficus alii* cultivars, the data in Table (4) showed that, in both seasons, within both cultivars of *Ficus alii*, the plants receiving a trace element treatments recorded significantly higher values than those recorded with the untreated control. The highest value in the first season (33.80 % of dry weight) was recorded on the cv. "Variegata" sprayed with Mn treatment, while the highest value in the second year (40.30 % of dry weight) was recorded on the cv. "Variegata" sprayed with B treatment. On the other hand, in both seasons, the lowest values recorded on carbohydrate percentages of both cultivars were obtained from untreated control plants.

3.2.3. N, P and K (% of dry weight)

As shown in Table (5) the data indicated that the foliar application of the different trace elements treatments had a considerable effect on the uptake and accumulation of N, P and K% in the leaves of *Ficus alii* plants. In both seasons, the plants sprayed with any trace element of the three nutrients were significantly higher than those recorded with the control. Similar increases in the N, P and K % as a result of the tested treatments have been reported by Said (1997) on *Codiaeum variegatum*, Al-Hadad (2006) on *Cupressus macrocarpa*, Mazhar *et al.* (2006) on *Taxodium disticum*, Abd El-Aziz and Balbaa (2007) on *Salvia farinacea*, Rajaie *et al.* (2009) on *Citrus aurantifolia*, Esmail (2008) on *Codiaeum variegatum* cv. "Gold Star"., Kumar and HariPriya (2010) on *Nerium odorum*, Eid *et al.*(2010) on *Polianthes tuberosa* and El- Attar (2011) on *Areca lutescens*.

Among the different trace element treatment, the applications of Mn followed by Fe treatment were the most effective for increasing the N % in the leaves in the first season, whereas the highest mean values in the second season were obtained in plants sprayed with B treatment followed by plants sprayed with Mn treatment. In both seasons, the most effective treatment for increasing the P % in the leaves was spraying the plants with Fe which gave the highest values (0.42 and 0.47 % of dry weight in the first and second seasons, respectively followed by spraying the plants with B. It is also clear from

the data in Table (5) that in both seasons, foliar application of Zn treatment was the most effective for increasing the K % in the leaves as it gave the highest values (2.10 and 2.29 % of dry weight in the first and second seasons, respectively, followed by foliar spraying of B treatment.

As for the cultivars of *Ficus alii*, the data in Table (5) also showed that in both seasons, the accumulation of N% in the leaves of the cv. "Variegata" was significantly higher than the cv. "Green". In the first season, the cv. "Variegata" had P % in their leaves significantly higher than in the cv. "Green". However, in the second season, there were no significant differences between the cv. "Green" and the cv. "Variegata" in this respect. The accumulation of K% differed from one season to another, where in the first season the cv. "Variegata" had K % in the leaves which was significantly higher than the cv. "Green". On the other hand, in the second season another trend was obtained, *i.e.* K % in leaves of the cv. "Green" was significantly higher than the cv. "Variegata".

Regarding the interaction effect of foliar application of the different trace elements on N, P and K% in the leaves of *Ficus alii* cultivars, the data in Table (5) showed that in both seasons, spraying the plants with any trace element caused significant increase in the treatments of the three nutrients in leaves of both cultivars as compared to the control. In both seasons, the highest mean values of N % in leaves of both cultivars were obtained from plants sprayed with Mn treatment followed by plants sprayed with Fe treatment, whereas the highest mean values of P % in leaves of both cultivars were resulted from plants sprayed with Fe treatment followed by plants sprayed with B treatment, while the highest mean values of K % in leaves of both cultivars were produced from plants sprayed with Zn treatment followed by plants sprayed with B treatment(in most cases). On the other hand, in both seasons the lowest values recorded on the three nutrients in leaves of both cultivars were obtained from untreated control plants.

3.2.4. Zn, Fe, Mn and B (ppm)

Results of chemical analysis of dried leaves of *Ficus alii* plants (Table 6) showed that the uptake and accumulation of the different nutrients (Zn, Fe, Mn and B) were enhanced, and exhibited a similar trend, in response to foliar application of the different treatments. In both

Table (5): Effect of foliar application of trace element treatments on N, P and K percentage (% of dry weight) in the leaves of *Ficus alii* cvs. "Green" and "Variegata", during the 2012 and 2013 seasons.

*Trace elements treatments (T)	First season (2012)			Second season (2013)		
	Cultivars (C)		Mean (T)	Cultivars (C)		Mean (T)
	Green	Variegata		Green	Variegata	
N (% dry weight)						
Control	1.88	2.03	1.96	1.32	1.73	1.53
Zn	2.21	2.48	2.35	2.33	2.57	2.45
Fe	2.31	2.60	2.46	2.40	2.81	2.61
Mn	2.80	2.90	2.85	2.75	3.11	2.93
B	2.20	2.40	2.30	2.23	2.49	3.36
Mean (C)	2.28	2.48	-----	2.21	2.54	-----
L.S.D. (0.05)						
T	0.04			0.05		
C	0.02			0.03		
T X C	0.05			0.08		
P (% dry weight)						
Control	0.14	0.18	0.16	0.17	0.22	0.20
Zn	0.28	0.29	0.29	0.31	0.31	0.31
Fe	0.39	0.44	0.42	0.46	0.48	0.47
Mn	0.23	0.26	0.25	0.29	0.32	0.31
B	0.30	0.35	0.33	0.36	0.34	0.35
Mean (SL)	0.27	0.30	-----	0.32	0.33	-----
L.S.D. (0.05)						
T	0.04			0.05		
C	0.02			0.03		
T X C	0.05			0.08		
K (% dry weight)						
Control	0.98	1.10	1.04	1.07	1.16	1.12
Zn	2.06	2.13	2.10	2.33	2.24	2.29
Fe	2.01	1.97	1.99	2.10	2.03	2.07
Mn	1.20	1.58	1.39	2.00	1.85	1.93
B	1.98	2.04	2.01	2.31	2.16	2.24
Mean (SL)	1.65	1.76	-----	1.96	1.89	-----
L.S.D. (0.05)						
T	0.04			0.05		
C	0.02			0.03		
T X C	0.06			0.08		

* Zn, Fe and Mn = 50 mg/l B= 10 mg/l

Table (6): Effect of foliar application of trace element treatments on Zn, Fe, Mn and B contents (ppm) in the leaves of *Ficus alii* cvs "Green" and "Variegata", during the 2012 and 2013 seasons.

*Trace elements treatments (T)	First season (2012)			Second season (2013)		
	Cultivars (C)		Mean (T)	Cultivars (C)		Mean (T)
	Green	Variegata		Green	Variegata	
Zn (ppm)						
Control	41	38	39.5	33	40	36.5
Zn	85	88	86.5	90	96	93.0
Fe	77	80	78.5	88	85	86.5
Mn	74	80	77.0	70	74	72.0
B	80	79	79.5	75	80	77.5
Mean (C)	71.4	79	-----	71.2	75	-----
L.S.D. (0.05)						
T	2.4			1.7		
C	1.4			1.0		
T X C	3.4			2.4		
Fe (ppm)						
Control	174	180	177.0	184	193	188.5
Zn	321	340	330.5	345	400	372.5
Fe	399	390	394.5	422	420	421.0
Mn	304	325	314.5	410	316	363.0
B	301	311	306.0	300	469	384.5
Mean (SL)	299.8	309.2	-----	332.2	359.6	-----
L.S.D. (0.05)						
T	6.1			5.1		
C	3.5			3.0		
T X C	8.6			7.8		
Mn (ppm)						
Control	25	29	27	32	38	35
Zn	44	46	45	42	50	46
Fe	35	38	36.5	45	49	47
Mn	42	50	46	62	57	59.5
B	38	43	40.5	48	51	49.5
Mean (SL)	36.8	41.2	-----	45.8	49	-----
L.S.D. (0.05)						
T	2.5			2.6		
C	1.6			1.5		
T X C	3.6			3.6		
B (ppm)						
Control	0.37	0.42	0.40	0.44	0.47	0.46
Zn	0.56	0.62	0.59	0.68	0.72	0.70
Fe	0.64	0.73	0.69	0.90	0.83	0.87
Mn	0.55	0.65	0.60	0.70	0.75	0.73
B	0.86	0.90	0.88	0.99	0.95	0.97
Mean (SL)	0.60	0.66	-----	0.74	0.74	-----
L.S.D. (0.05)						
T	0.05			0.04		
C	0.03			0.02		
T X C	0.08			0.05		

* Zn, Fe and Mn = 50 mg/l B= 10 mg/l

seasons, the foliar application of different trace elements, regardless of the type of element applied, caused significant increase in these nutrients contents in the leaves of *Ficus alii* plants, compared to the control. These results are in agreement with the findings of Mattis and Hershey (1992) on *Epipremnum aureum* and *Philodendron scandens*, Said (1997) *Codiaeum variegatum*, Stamps and Rock (2000) *Asparagus virgatus*, Al-Hadad (2006) *Cupressus macrocarpa*, Esmail (2008) *Codiaeum variegatum* cv. "Gold Star", Rajaie *et al.* (2009) *Citrus aurantifolia*, Kumar and Haripriya (2010) *Nerium odorum*, Eid *et al.* (2010) *Polianthes tuberosa* and El- Attar (2011) *Areca lutescens*.

It is also obvious from the data in Table (6) that, in both seasons the highest values of Zn, Fe Mn and B contents were obtained in the leaves of plants sprayed with treatments of the same elements. In other words, foliar application of each element was the most effective treatment in increasing and accumulating of the same element content in leaves tissues.

The increase in the nutrient content in the leaves tissues of the plant as a result of foliar application of the nutrient solution can be easily explained, since nutrient solution remains on the leaf surface as a thin film which may be accompanied by more absorption from leaf surface, and their accumulation in plant leaves tissues (Taiz and Zeiger, 2002 and Fernandez *et al.* 2013).

Concerning the cultivars of *Ficus alii*, the data in Table (6) showed that, in most cases, the uptake and accumulation of Zn, Fe Mn and B contents in the leaves of the cv. "Variegata" were significantly higher than the cv. "Green". The only one exception to this general trend was recoded in the second season, since there were no significant differences between B content in the leaves of the cv. "Green" and the cv. "Variegata".

Regarding the interaction effect of foliar application of the different trace elements treatments on Zn, Fe Mn and B contents in the eaves of *Ficus alii* cultivars, the data recorded in Table (6) indicated that in both seasons, all nutrient contents significantly increased in the leaves of both cultivars as a result of foliar application of trace elements, compared to the untreated control. In both seasons, the highest values of nutrient content in leaves of both cultivars, in most cases, were obtained from plants sprayed with the solution of the same

element. On the other hand, lowest values of nutrient content of both cultivars were recorded on the control plants.

Conclusion: Foliar application of the trace elements treatments improved growth and quality of *Ficus alii* cvs "Green" and "Variegata", the cv. "Variegata" was more responsive than the cv. "Green". The importance of the studied trace elements for both cultivars can be arranged as Mn > Fe > B > Zn.

From the above mentioned results, it can be recommended that for the best vegetative growth of *Ficus alii* cvs "Green" and "Variegata", foliar spray with Fe, Zn or Mn at 50 mg/l/ 3 weeks and/or B at 10 mg/l/ 3 weeks and supplied with 2g/plant/month of chemical fertilizer (NPK, 20-20-20) is recommended .

4. REFERENCES

- Abd El-Aziz N. G. and Balbaa L. K. (2007). Influence of tyrosine and zinc on growth, flowering and chemical constituents of *Salvia farinacea* P. Plants J. Appl. Sci. Res., 3(11): 1479-1489.
- Al-Hadad Z. Kh. (2006). Effect of some treatments on growth and chemical composition of *Cupressus macrocarpa*. M.Sc. Thesis, Fac. Agric., Cairo, Univ., Egypt, 83 p.
- Allen V. B. and David J. P. (2007). Handbook of Plant Nutrition. Taylor and Francis Publishers, NewYork USA. 662p.
- Chapman H.D. and Pratt P.F. (1961). Methods of Soil, Plants and Water Analysis. Univ., of California, USA Division of Agricultural Sciences, pp. 60-69.
- Dorgham A. H. (2005). Physiological studies on *Dieffenbachia*, *Philodendron* and *Syngonium* plants .M. Sc. Thesis, Fac. Agric., Cairo Univ. Egypt, 193 p.
- Dubois M., Smith F., Gilles K. A., Hamilton J. K. and Rebers P. A. (1956). Colorimetric method for determination of sugar and related substances. Anal. Chem., 28 (3): 350-356.
- Eid R. A., Khalifa R.Kh. M. and Shaaban S.H.A. (2010). Effect of foliar application of zinc and benzyladenine on growth, yield and chemical constituents of tuberose plants. Res. J Agric. and Biol. Sci., 6 (6): 732-743.
- El-Attar A. B. (2011). Response of *Areca lutescens*, *Caryota mitis* and *Chamaedorea elegans* palms to nutritional

- and irrigation treatments. Ph.D. Thesis, Fac. Agric., Cairo, Univ., Egypt, 157 p.
- El-Bagoury H. M., Nasr A. A. and El-Tantawy A. A. (2008). Effect of NPK fertilization, trace elements and gibberellic acid on vegetative growth and chlorophyll content of *Hedera helix*, l. plants. J. Product. & Dev., 13(1), 19-27.
- El-Deeb E. A. (1999). Effect of Mineral Nutrition, Planting Media and Light on Some Foliage Plants. M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt, 128 p.
- El-Fadaly H. G. (1994). Effect of Chemical Fertilization and Gibberellic Acid (GA₃) Treatments on Growth, Flowering and Chemical Composition of *Jasminum sambac*. L. Plants. M.Sc. Thesis, Fac. Agric.; Cairo Univ., 172 p.
- Esmail S. A. (2008). Physiological studies on croton plant. M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt, 131 p.
- Farahat M. M., Ibrahim S., Taha S. L. and El-Ouesni F. M. (2007). Response of vegetative growth and some chemical constituents of *Cupressus sempervirens* L. to foliar application of ascorbic acid and zinc at Nubaria. World, J. Agric. Sci., 3(3): 282-288.
- Fernandez V., Sotiropoulos T. and Brown P. (2013). Foliar Fertilization Scientific Principles and Field Practices. International Fertilizer Industry Association (IFA), Paris, France.
- Gomaa S.A. and Ragab A.A.M (2005). Studies on the effect of microelements and gibberellic acid on *Codiaeum variegatum* cv. Redii. Egypt. J. Appl. Sci., 20(5):272-289.
- Heider S. M. A. O. (2010). Effect of chemical fertilization and growing media on vegetative growth and chemical composition on two cultivars of *Dracena marginata*. M.Sc. Thesis, Fac. Agric., Cairo, Univ., Egypt, 113 p.
- Isaac R. A. and Kerber J. D. (1971). Atomic Absorption and Flamephotometry: Techniques and Uses in Soil, Plant, and Water Analysis. In Instrumental Methods for Analysis of Soils and Plant Tissue. Soil Sci. Soc. Amer., Madison, WI, USA, pp 17-37.
- Kuepper G. (2003). Foliar fertilization. ATTRA (Appropriate Technology Transfer for Rural Areas), US Dept. Agric., pp: 1-10.
- Kumar S. and Haripriya K. (2010). Effect of foliar application of iron and zinc on growth, flowering and yield on nerium (*Nerium odorum* L.). Plant Archives, 10 (2): 637-640.
- Mansour N. M. (2002). A study on the acclimatization of *Scindapsus pictus* and *Hedra helix* and their response to chemical fertilization. M.Sc. Thesis, Fac. Agric., Cairo, Univ., Egypt, 131 p.
- Marschner H. (1995). Mineral Nutrition of Higher Plants. 2nd ed., Academic Press, London, UK.
- Mattis P. R. and Hershey D. R. (1992). Iron deficiency stress response of *Epipremnum aureum* and *Philodendron scandens*. J. Plant Physiol., 139 (4):498-502.
- Mazhar A. A. M., Zaghoul S. and Yassen A. A. (2006). Impact of boron fertilizer on growth and chemical constituents of *Taxodium distichum* grown under water regime. World, J. Agric. Sci., 2(4): 412-420.
- Naguib N. Y., Hussein M. S., Sherbeny E. S. E., Khalil M. Y. and Lazari D. (2007). Response of *Ruta graveolens* L. to sowing dates and foliar micronutrients. J. Appl. Sci. Res., 3(11): 1534-1543.
- Piper C.S. (1947). Soil and Plant Analysis. Univ. of Adelaide, Adelaide, Australia, pp. 258-275.
- Pregl P. (1945). Quantitative Organic Microanalysis. Churchill Publishing Co., London, UK., 4th Ed. pp 78-82.
- Rajaie M., Ejraieb A. K., Owliaiee H. R. and Tavakolid A. R. (2009). Effect of zinc and boron interaction on growth and mineral composition of lemon seedlings in a calcareous soil. Int., J. Plant Prod. 3(1): 39-50.
- Said R. M. (1997). Effect of Some Chemical Fertilization on Croton Plant. M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt, 180 p.
- Saric M., Kastrori R., Curic R., Cupina T. and Geric I. (1967). Chlorophyll Determination. Univ. Unoven Sadu Par Ktikum is fiziologize Biljaka, Beogard, Hauncna, Anjiga, P. 215.
- Stamps R. and Rock D. K. (2000). Effects of four elements on color, yield and vase life of tree "fern" (*Asparagus virgatus*). Proceed. Fla. State Hort. Soc., 112:282-285.

Steel R. G. and Torrie S. H. (1980). Principles and Procedures of Statistics. Second Ed., McGraw-Hill Inc., New York. USA pp.63.
Taiz L. and Zeiger E. (2002). Plant Physiology, 3rd Ed., Sinauer Associates, Sunderland,

MA. 690 pps.

Troug E. and Meyer A. H. (1939). Improvem in deiness colorimetric methods for phosphorus and arsenic. Ind. Eng. Chem. Anal., Ed. 1: 136-139.

استجابته صنفين من الـ *Ficus alii* L. الى الرش الورقي ببعض معاملات العناصر الصغرى

أسماء بدرالدين العطار

قسم بساتين الزينه - كلية الزراعة - جامعة القاهرة - الجيزة - مصر

ملخص

أجرى هذا البحث داخل الصوبة الزجاجية بمشتمل قسم بساتين الزينه بكلية الزراعة - جامعة القاهرة خلال الموسمين المتتاليين (2012 و 2013). بهدف دراسة استجابة صنفين من *Ficus alii* (جرين وفاريجاتا) الى الرش الورقي بالعناصر الصغرى. تم رش النباتات بعناصر الزنك، الحديد والمنجنيز بمعدل 50 ملجم /لتر/ 3 أسابيع/نبات/ اصيص 30سم، وعنصر البورون بمعدل 10 ملجم /لتر/ 3 أسابيع، بينما تم رش نباتات الكنترول بماء الصنبور. سمدت كل نباتات التجربه شهريا بسماد لايف جرين (NPK 20:20:20) بمعدل 2جم /اصيص. أوضحت النتائج أن معاملات الرش الورقي بكل من الزنك، الحديد، المنجنيز والبورون ادت الى زياده معنويه فى النمو الخضرى وكذلك محتوى الكلوروفيل الكلى وتركيز الكربوهيدرات الكلية، النتروجين، الفسفور، البوتاسيوم، الزنك، الحديد، المنجنيز والبورون فى الاوراق مقارنة بنباتات الكنترول. وعلى العكس، أدت معاملات العناصر الصغرى (فى اغلب الحالات) الى نقص معنوى فى محتوى الكاروتينويدات فى الاوراق مقارنة بالنباتات غير المعامله. فى كلا الموسمين كانت، صفات ارتفاع النبات ومساحه الورق ومحتوى الكاروتينويدات فى الاوراق فى الصنف جرين اعلى معنويا من الصنف فاريجاتا. على العكس، كانت صفات عدد الاوراق وقطر الساق والوزن الطازج والجاف للسيقان والجنور وكذلك محتوى الكلوروفيل الكلى وتركيز الكربوهيدرات الكلية و النتروجين والفسفور والبوتاسيوم والزنك والحديد والمنجنيز والبورون فى الاوراق فى الصنف فاريجاتا اعلى معنويا من الصنف جرين. فى كلا الموسمين، مع اى من صنفى *Ficus alii* ادى الرش الورقي بالمعاملات المختلفه للعناصر الصغرى الى زياده معنويه فى معظم خصائص النمو الخضرى، وكذلك محتوى الكلوروفيل الكلى وتركيز الكربوهيدرات الكلية، النتروجين، الفسفور، البوتاسيوم، الزنك، الحديد، المنجنيز والبورون فى الاوراق مقارنة بنباتات الكنترول. يمكن من النتائج المتحصل عليها التوصية بانه للحصول على أفضل نمو خضرى لاصناف *Ficus alii* (جرين وفاريجاتا) وذلك برش النباتات شهريا بعناصر الزنك والحديد والمنجنيز بمعدل 50 ملجم / لتر/ 3 أسابيع، وعنصر البورون بمعدل 10 ملجم /لتر/ 3 أسابيع .

المجلة العلمية لكلية الزراعة - جامعة القاهرة - المجلد (65) العدد الثانى (ابريل 2014):170-182 .