

Journal of Fractional Calculus and Applications,
 Vol. 2. Jan. 2012, No. 9, pp. 1–15.
 ISSN: 2090-5858.
<http://www.fcaj.webs.com/>

NEW INEQUALITIES OF OSTROWSKI TYPE FOR CO-ORDINATED CONVEX FUNCTIONS VIA FRACTIONAL INTEGRALS

MUHAMMAD AMER LATIF, SABIR HUSSAIN

ABSTRACT. In this paper an identity similar to an identity proved in [31] for fractional integrals for functions of two variables is established. Some new Ostrowski type inequalities for Riemann-Liouville fractional integrals of functions of two variables are established as well. The results established in this paper generalize those results proved in [31].

1. INTRODUCTION

In 1938, A. Ostrowski proved the following interesting inequality [33]:

Theorem 1. [33] *Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a, b) whose derivative $f' : (a, b) \rightarrow \mathbb{R}$ is bounded on (a, b) , i.e., $\|f'\|_{\infty} := \sup_{t \in (a, b)} |f'(t)| < \infty$.*

The we have the inequality

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_{\infty}, \quad (1.1)$$

for all $x \in [a, b]$. The constant $\frac{1}{4}$ is the best possible.

The inequality (1.1) can be rewritten in equivalent form as:

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{(x-a)^2 + (b-x)^2}{2(b-a)} \right] \|f'\|_{\infty}.$$

Since 1938 when A. Ostrowski proved his famous inequality, many mathematicians have been working about and around it, in many different directions and with a lot of applications in Numerical Analysis and Probability, etc.

Several generalizations of the Ostrowski integral inequality for mappings of bounded variation, Lipschitzian, monotonic, absolutely continuous, convex mappings, quasi convex mappings and n -times differentiable mappings with error estimates for some

2000 *Mathematics Subject Classification.* 26A33, 26A51, 26D07, 26D10, 26D15.

Key words and phrases. Ostrowski inequality, co-ordinated convex function, Riemann-Liouville fractional integral.

Submitted Nov. 18, 2011. Published March 1, 2012.

special means and for some numerical quadrature rules are considered by many authors. For recent results and generalizations concerning Ostrowski's inequality see [2]-[4], [9], [11], [15]-[19], [27], [36]-[39], [44] and [45] and the references therein.

Let us consider now a bidimensional interval $\Delta =: [a, b] \times [c, d]$ in \mathbb{R}^2 with $a < b$ and $c < d$, a mapping $f : \Delta \rightarrow \mathbb{R}^2$ is said to be convex on Δ if the inequality

$$f(\lambda x + (1 - \lambda)z, \lambda y + (1 - \lambda)w) \leq \lambda f(x, y) + (1 - \lambda)f(z, w),$$

holds for all $(x, y), (z, w) \in \Delta$ and $\lambda \in [0, 1]$. The mapping f is said to be concave on the co-ordinates on Δ if the above inequality holds in reversed direction, for all $(x, y), (z, w) \in \Delta$ and $\lambda \in [0, 1]$.

A modification for convex (concave) functions on Δ , which are also known as co-ordinated convex (concave) functions, was introduced by S. S. Dragomir [12] as follows:

A function $f : \Delta \rightarrow \mathbb{R}$ is said to be convex (concave) on the co-ordinates on Δ if the partial mappings $f_y : [a, b] \rightarrow \mathbb{R}, f_y(u) = f(u, y)$ and $f_x : [c, d] \rightarrow \mathbb{R}, f_x(v) = f(x, v)$ are convex (concave) where defined for all $x \in [a, b], y \in [c, d]$.

A formal definition for co-ordinated convex (concave) functions may be stated in:

Definition 1. [29] A mapping $f : \Delta \rightarrow \mathbb{R}$ is said to be convex on the co-ordinates on Δ if the inequality

$$\begin{aligned} & f(tx + (1 - t)y, su + (1 - s)w) \\ & \leq tsf(x, u) + t(1 - s)f(x, w) + s(1 - t)f(y, u) + (1 - t)(1 - s)f(y, w), \end{aligned} \quad (1.2)$$

holds for all $t, s \in [0, 1]$ and $(x, u), (y, w) \in \Delta$. The mapping f is concave on the co-ordinates on Δ if the inequality (1.2) holds in reversed direction for all $t, s \in [0, 1]$ and $(x, u), (y, w) \in \Delta$.

Clearly, every convex (concave) mapping $f : \Delta \rightarrow \mathbb{R}$ is convex (concave) on the co-ordinates. Furthermore, there exists co-ordinated convex (concave) function which is not convex (concave), (see for instance [12]).

Here we also quote the following result form [12] to be used in the sequel of the paper:

Theorem 2. Suppose that $f : \Delta \rightarrow \mathbb{R}$ is co-ordinated convex on Δ . Then one has the inequalities:

$$\begin{aligned} & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \leq \frac{1}{2} \left[\frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}\right) dx + \frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y\right) dy \right] \\ & \leq \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx \\ & \leq \frac{1}{4(b-a)} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{4(d-c)} \int_c^d [f(a, y) + f(b, y)] dy \\ & \leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4}. \end{aligned} \quad (1.3)$$

The above inequalities are sharp. The inequalities in (1.3) hold in reverse direction if the mapping f is concave.

In recent years, many authors have proved several inequalities for co-ordinated convex functions. These studies include, among others, the works in [5]-[8], [12,

[13, 26], [28]-[31], [35] and [42] (see also the references therein). Alomari et al. [5]-[7], proved several Hermite-Hadamard type inequalities for co-ordinated s -convex functions. Bakula et. al [8], proved Jensen's inequality for convex functions on the co-ordinates from the rectangle from the plan. Dragomir [12], proved the Hermite-Hadamard type inequalities for co-ordinated convex functions. Hwang et. al [26], also proved some Hermite-Hadamard type inequalities for co-ordinated convex function of two variables by considering some mappings directly associated to the Hermite-Hadamard type inequality for co-ordinated convex mappings of two variables. Latif et. al [28]-[31], proved some inequalities of Hermite-Hadamard type for differentiable co-ordinated convex function, product of two co-ordinated convex mappings, for co-ordinated h -convex mappings and some Ostrowski type inequalities for co-ordinated convex mappings. Özdemir et. al [35], proved Hadamard's type inequalities for co-ordinated m -convex and (α, m) -convex functions. Sarikaya, et. al proved Hermite-Hadamard type inequalities for differentiable co-ordinated convex function. For more inequalities on co-ordinated convex functions see also the references in the above cited papers.

In the present paper, we establish new Ostrowski type inequalities for co-ordinated convex functions similar to those from [31] but via Riemann-Liouville fractional integral.

2. MAIN RESULTS

We give first some necessary definitions and mathematical preliminaries of fractional calculus theory which are used in this sections.

Definition 2. Let $f \in L_1[a, b]$. The Riemann-Liouville integrals $J_{a+}^{\alpha} f$ and $J_{b-}^{\alpha} f$ of order $\alpha > 0$ with $a \geq 0$ are defined by

$$\begin{aligned} J_{a+}^{\alpha} f(x) &= \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, x > a \\ J_{b-}^{\alpha} f(x) &= \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, x < b, \end{aligned}$$

where $\Gamma(\alpha) = \int_0^{\infty} e^{-u} u^{\alpha-1} du$. It is to be noted that $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$.

In the case $\alpha = 1$, the fractional integral reduces to the classical integral.

For further properties and results concerning this operator we refer the intrusted reader to [1], [10], [20]-[24], [32], [40] and [41].

For the sake of convenience, we will use the following notation throughout this section:

$$\begin{aligned} A &= \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)(d-c)} \left[J_{x-,y-}^{\alpha,\beta} f(a, c) + J_{x-,y+}^{\alpha,\beta} f(a, d) + J_{x+,y-}^{\alpha,\beta} f(b, c) \right. \\ &\quad \left. + J_{x+,y+}^{\alpha,\beta} f(b, d) \right] - \frac{[(x-a)^{\alpha} + (b-x)^{\alpha}]\Gamma(\beta+1)}{(b-a)(d-c)} \left[J_{y-}^{\beta} f(x, c) + J_{y+}^{\beta} f(x, d) \right] \\ &\quad - \frac{[(y-c)^{\beta} + (d-y)^{\beta}]\Gamma(\alpha+1)}{(b-a)(d-c)} \left[J_{x-}^{\alpha} f(a, y) + J_{x+}^{\alpha} f(b, y) \right], \end{aligned}$$

where

$$\begin{aligned}
 J_{a+,c+}^{\alpha,\beta} f(x,y) &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_c^y (x-u)^{\alpha-1} (y-v)^{\beta-1} f(u,v) dv du, x > a, y > c, \\
 J_{b-,d-}^{\alpha,\beta} f(x,y) &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_y^d (u-x)^{\alpha-1} (v-y)^{\beta-1} f(u,v) ddv du, x < b, y < d, \\
 J_{a+,d-}^{\alpha,\beta} f(x,y) &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_y^d (x-u)^{\alpha-1} (v-y)^{\beta-1} f(u,v) ddv du, x > a, y < d, \\
 J_{b-,c+}^{\alpha,\beta} f(x,y) &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_c^y (u-x)^{\alpha-1} (y-v)^{\beta-1} f(u,v) ddv du, x < b, y > c,
 \end{aligned}$$

and Γ is the Euler Gamma function.

To establish our main results we need the following identity:

Lemma 1. Let $f : \Delta := [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a twice partial differentiable mapping on Δ° with $a < b$, $c < d$. If $\frac{\partial^2 f}{\partial s \partial t} \in L(\Delta)$ and $\alpha, \beta > 0$, $a, c \geq 0$, then the following identity holds:

$$\begin{aligned}
 & \frac{[(b-x)^\alpha + (x-a)^\alpha] \left[(d-y)^\beta + (y-c)^\beta \right]}{(b-a)(d-c)} f(x,y) + A \\
 &= \frac{(x-a)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) ds dt \\
 & - \frac{(x-a)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) ds dt \\
 & - \frac{(b-x)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) ds dt \\
 & + \frac{(b-x)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) ds dt,
 \end{aligned} \tag{2.1}$$

for all $(x, y) \in \Delta$.

Proof. By integration by parts and by the change of the variables $u = tx + (1 - t)a$, $v = sy + (1 - s)c$, we have

$$\begin{aligned}
& \frac{(x-a)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) ds dt \\
&= \frac{(x-a)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 t^\alpha \left[s^\beta \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) ds \right] dt \\
&\quad = \frac{(x-a)^\alpha (y-c)^\beta}{(b-a)(d-c)} f(x, y) \\
&- \frac{\beta (x-a)^\alpha}{(b-a)(d-c)} \int_c^y (v-c)^{\beta-1} f(x, v) dv - \frac{\alpha (y-c)^\beta}{(b-a)(d-c)} \int_a^x (u-a)^{\alpha-1} f(u, y) du \\
&\quad + \frac{\alpha \beta}{(b-a)(d-c)} \int_a^x \int_c^y (u-a)^{\alpha-1} (v-c)^{\beta-1} f(u, v) du dv \\
&= \frac{(x-a)^\alpha (y-c)^\beta}{(b-a)(d-c)} f(x, y) - \frac{\beta (x-a)^\alpha \Gamma(\beta)}{(b-a)(d-c)} \frac{1}{\Gamma(\beta)} \int_c^y (v-c)^{\beta-1} f(x, v) dv \\
&\quad - \frac{\alpha (y-c)^\beta \Gamma(\alpha)}{(b-a)(d-c)} \frac{1}{\Gamma(\alpha)} \int_a^x (u-a)^{\alpha-1} f(u, y) du \\
&\quad + \frac{\alpha \beta \Gamma(\alpha) \Gamma(\beta)}{(b-a)(d-c) \Gamma(\alpha) \Gamma(\beta)} \frac{1}{\Gamma(\alpha) \Gamma(\beta)} \int_a^x \int_c^y (u-a)^{\alpha-1} (v-c)^{\beta-1} f(u, v) dv du \\
&\quad = \frac{(x-a)^\alpha (y-c)^\beta}{(b-a)(d-c)} f(x, y) - \frac{(x-a)^\alpha \Gamma(\beta+1)}{(b-a)(d-c)} J_{y-}^\beta f(x, c) \\
&\quad - \frac{(y-c)^\beta \Gamma(\alpha+1)}{(b-a)(d-c)} J_{x-}^\alpha f(a, y) + \frac{\Gamma(\alpha+1) \Gamma(\beta+1)}{(b-a)(d-c)} J_{x-, y-}^{\alpha, \beta} f(a, c) \quad (2.2)
\end{aligned}$$

Similarly, by integration by parts, we also have

$$\begin{aligned}
& \frac{(x-a)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) ds dt \\
&= - \frac{(x-a)^\alpha (d-y)^\beta}{(b-a)(d-c)} f(x, y) + \frac{(x-a)^\alpha \Gamma(\beta+1)}{(b-a)(d-c)} J_{y+}^\beta f(x, d) \\
&\quad + \frac{(d-y)^\beta \Gamma(\alpha+1)}{(b-a)(d-c)} J_{x-}^\alpha f(a, y) - \frac{\Gamma(\alpha+1) \Gamma(\beta+1)}{(b-a)(d-c)} J_{x-, y+}^{\alpha, \beta} f(a, d), \quad (2.3)
\end{aligned}$$

$$\begin{aligned}
& \frac{(b-x)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) ds dt \\
&= - \frac{(b-x)^\alpha (y-c)^\beta}{(b-a)(d-c)} f(x, y) + \frac{(b-x)^\alpha \Gamma(\beta+1)}{(b-a)(d-c)} J_{y-}^\beta f(x, c) \\
&\quad + \frac{(y-c)^\beta \Gamma(\alpha+1)}{(b-a)(d-c)} J_{x+}^\alpha f(b, y) - \frac{\Gamma(\alpha+1) \Gamma(\beta+1)}{(b-a)(d-c)} J_{x+, y-}^{\alpha, \beta} f(b, c) \quad (2.4)
\end{aligned}$$

and

$$\begin{aligned} & \frac{(b-x)^{\alpha+1}(d-y)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) ds dt \\ &= \frac{(b-x)^\alpha(d-y)^\beta}{(b-a)(d-c)} f(x, y) - \frac{(b-x)^\alpha \Gamma(\beta+1)}{(b-a)(d-c)} J_{y+}^\beta f(x, d) \\ & - \frac{(d-y)^\beta \Gamma(\alpha+1)}{(b-a)(d-c)} J_{x+}^\alpha f(b, y) + \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)(d-c)} J_{x+, y+}^{\alpha, \beta} f(b, d). \quad (2.5) \end{aligned}$$

From (2.2)-(2.5), we get (2.1). This completes the proof. \square

Theorem 3. Let $f : \Delta := [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a twice partial differentiable mapping on Δ° with $a < b$, $c < d$, $a, c \geq 0$ such that $\frac{\partial^2 f}{\partial s \partial t} \in L(\Delta)$. If $\left| \frac{\partial^2 f}{\partial s \partial t} \right|$ is convex on the co-ordinates on Δ and $\left| \frac{\partial^2 f}{\partial y \partial x} f(x, y) \right| \leq M$, $(x, y) \in \Delta$, then the following inequality for fractional integrals with $\alpha, \beta > 0$ holds:

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \\ & \leq \frac{(\alpha\beta + 2\alpha + 2\beta + 4) [(b-x)^{\alpha+1} + (x-a)^{\alpha+1}] [(d-y)^{\beta+1} + (y-c)^{\beta+1}]}{(b-a)(d-c)(\alpha+1)(\alpha+2)(\beta+1)(\beta+2)} M \quad (2.6) \end{aligned}$$

for all $(x, y) \in \Delta$, where Γ is the Euler Gamma function.

Proof. From Lemma 1, we have that the following inequality holds for all $(x, y) \in \Delta$:

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \\ & \leq \frac{(x-a)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right| ds dt \\ & + \frac{(x-a)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right| ds dt \\ & + \frac{(b-x)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right| ds dt \\ & + \frac{(b-x)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 s^\beta t^\alpha \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right| ds dt. \quad (2.7) \end{aligned}$$

By the convexity of $\left| \frac{\partial^2 f}{\partial s \partial t} \right|$ on the co-ordinates on Δ and $\left| \frac{\partial^2}{\partial y \partial x} f(x, y) \right| \leq M$, $(x, y) \in \Delta$, we get the following inequalities:

$$\begin{aligned} & \int_0^1 \int_0^1 s^\beta t^\alpha \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right| ds dt \\ & \leq M \int_0^1 \int_0^1 s^{\beta+1} t^{\alpha+1} ds dt + M \int_0^1 \int_0^1 t^{\alpha+1} (s^\beta - s^{\beta+1}) ds dt \\ & + M \int_0^1 \int_0^1 s^{\beta+1} (t^\alpha - t^{\alpha+1}) ds dt + M \int_0^1 \int_0^1 (t^\alpha - t^{\alpha+1}) (s^\beta - s^{\beta+1}) ds dt \\ & = \frac{\alpha\beta + 2\alpha + 2\beta + 4}{(\alpha+1)(\alpha+2)(\beta+1)(\beta+2)} M. \end{aligned} \quad (2.8)$$

Analogously, we also have the following inequalities:

$$\begin{aligned} & \int_0^1 \int_0^1 s^\beta t^\alpha \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right| ds dt \\ & \leq \frac{\alpha\beta + 2\alpha + 2\beta + 4}{(\alpha+1)(\alpha+2)(\beta+1)(\beta+2)} M, \end{aligned} \quad (2.9)$$

$$\begin{aligned} & \int_0^1 \int_0^1 s^\beta t^\alpha \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right| ds dt \\ & \leq \frac{\alpha\beta + 2\alpha + 2\beta + 4}{(\alpha+1)(\alpha+2)(\beta+1)(\beta+2)} M \end{aligned} \quad (2.10)$$

and

$$\begin{aligned} & \int_0^1 \int_0^1 s^\beta t^\alpha \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right| ds dt \\ & \leq \frac{\alpha\beta + 2\alpha + 2\beta + 4}{(\alpha+1)(\alpha+2)(\beta+1)(\beta+2)} M. \end{aligned} \quad (2.11)$$

By using (2.8)-(2.11) in (2.7), we get the desired inequality (2.6). This completes the proof of the theorem. \square

Remark 1. In Theorem 1, if we take $\alpha = \beta = 1$, then the inequality (2.6) reduces to the inequality established in [31, Theorem 3].

The next result is about the powers of the absolute value of the partial derivatives.

Theorem 4. Let $f : \Delta := [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a twice partial differentiable mapping on Δ° with $a < b$, $c < d$, $a, c \geq 0$ such that $\left| \frac{\partial^2 f}{\partial s \partial t} \right| \in L(\Delta)$. If $\left| \frac{\partial^2 f}{\partial s \partial t} \right|^q$ is convex on the co-ordinates on Δ , $p, q > 1$, $\frac{1}{p} + \frac{1}{q} = 1$ and $\left| \frac{\partial^2}{\partial y \partial x} f(x, y) \right| \leq M$, $(x, y) \in \Delta$, then

the following inequality for fractional integrals with $\alpha, \beta > 0$ holds:

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \\ & \leq \frac{1}{(\alpha p+1)^{\frac{1}{p}} (\beta p+1)^{\frac{1}{p}}} \left[\frac{(b-x)^{\alpha+1} + (x-a)^{\alpha+1}}{b-a} \right] \left[\frac{(d-y)^{\beta+1} + (y-c)^{\beta+1}}{d-c} \right] M, \end{aligned} \quad (2.12)$$

for all $(x, y) \in \Delta$.

Proof. From Lemma 1 and the Hölder inequality, we have that the following inequality holds, for all $(x, y) \in \Delta$:

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \leq \left(\int_0^1 \int_0^1 t^{\alpha p} s^{\alpha p} ds dt \right)^{\frac{1}{p}} \\ & \left[\frac{(x-a)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right|^q ds dt \right)^{\frac{1}{q}} \right. \\ & + \frac{(x-a)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right|^q ds dt \right)^{\frac{1}{q}} \\ & + \frac{(b-x)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right|^q ds dt \right)^{\frac{1}{q}} \\ & \left. + \frac{(b-x)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right|^q ds dt \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (2.13)$$

By the co-ordinated convexity of f and $\left| \frac{\partial^2}{\partial y \partial x} f(x, y) \right| \leq M$, for all $(x, y) \in \Delta$, we have that the following inequality holds:

$$\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right|^q ds dt \leq M^q.$$

Similarly, we also have the following inequalities:

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right|^q ds dt \leq M^q, \\ & \int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right|^q ds dt \leq M^q \end{aligned}$$

and

$$\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right|^q ds dt \leq M^q.$$

Using the fact

$$\int_0^1 \int_0^1 t^{\alpha p} s^{\alpha p} ds dt = \frac{1}{(\alpha p+1)(\beta p+1)}$$

and using the last four inequalities in (2.13), we obtain (2.12). This completes the proof of the theorem. \square

Remark 2. In Theorem 4, if we take $\alpha = \beta = 1$, then the inequality (2.12) becomes the inequality proved in [31, Theorem 4].

A different approach leads us to the following result:

Theorem 5. Let $f : \Delta := [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a twice partial differentiable mapping on Δ° with $a < b$, $c < d$, $a, c \geq 0$ such that $\frac{\partial^2 f}{\partial s \partial t} \in L(\Delta)$. If $\left| \frac{\partial^2 f}{\partial s \partial t} \right|^q$ is convex on the co-ordinates on Δ , $q \geq 1$ and $\left| \frac{\partial^2 f}{\partial y \partial x} f(x, y) \right| \leq M$, $(x, y) \in \Delta$, then the following inequality for fractional integrals with $\alpha, \beta > 0$ holds:

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \\ & \leq \frac{1}{(\alpha p+1)^{\frac{1}{p}} (\beta p+1)^{\frac{1}{p}}} \left[\frac{(b-x)^{\alpha+1} + (x-a)^{\alpha+1}}{b-a} \right] \left[\frac{(d-y)^{\beta+1} + (y-c)^{\beta+1}}{d-c} \right] M, \end{aligned} \quad (2.14)$$

for all $(x, y) \in \Delta$.

Proof. From Lemma 1 and the power mean inequality, we have that the following inequality holds, for all $(x, y) \in \Delta$:

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \leq \left(\int_0^1 \int_0^1 ts ds dt \right)^{1-\frac{1}{q}} \\ & \left[\frac{(x-a)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right|^q ds dt \right)^{\frac{1}{q}} \right. \\ & + \frac{(x-a)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right|^q ds dt \right)^{\frac{1}{q}} \\ & + \frac{(b-x)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right|^q ds dt \right)^{\frac{1}{q}} \\ & \left. + \frac{(b-x)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right|^q ds dt \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (2.15)$$

By the co-ordinated convexity of f and $\left| \frac{\partial^2 f}{\partial y \partial x} f(x, y) \right| \leq M$, for all $(x, y) \in \Delta$, we have that the following inequality holds:

$$\int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right|^q ds dt \leq \frac{1}{4} M^q.$$

In a similarly way, we also have the following inequalities:

$$\begin{aligned} & \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right|^q ds dt \leq \frac{1}{4} M^q, \\ & \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right|^q ds dt \leq \frac{1}{4} M^q \end{aligned}$$

and

$$\int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right|^q ds dt \leq \frac{1}{4} M^q.$$

Using the fact

$$\int_0^1 \int_0^1 ts ds dt = \frac{1}{4}$$

and the last four inequalities, we obtain from (2.15) the inequality (2.14). This completes the proof of the theorem. \square

Remark 3. In Theorem 5, if we take $\alpha = \beta = 1$, then the inequality (2.14) becomes the inequality proved in [31, Theorem 5].

Now we drive some results with co-ordinated concavity property instead of co-ordinated convexity.

Theorem 6. Let $f : \Delta \rightarrow \mathbb{R}$ be a twice partial differentiable mapping on Δ° such that $\frac{\partial^2 f}{\partial s \partial t} \in L(\Delta)$. If $\left| \frac{\partial^2 f}{\partial s \partial t} \right|^q$ is concave on the co-ordinates on Δ and $p, q > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, then the inequality

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \\ & \leq \frac{1}{(1+\alpha p)^{\frac{1}{p}} (1+\beta p)^{\frac{1}{p}} (b-a)(d-c)} \left[(x-a)^{\alpha+1} (y-c)^{\beta+1} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{x+a}{2}, \frac{y-c}{2} \right) \right| \right. \\ & \quad + (x-a)^{\alpha+1} (d-y)^{\beta+1} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{x+a}{2}, \frac{d+y}{2} \right) \right| \\ & \quad + (b-x)^{\alpha+1} (y-c)^{\beta+1} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{b+x}{2}, \frac{y+c}{2} \right) \right| \\ & \quad \left. + (b-x)^{\alpha+1} (d-y)^{\beta+1} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{b+x}{2}, \frac{d+y}{2} \right) \right| \right], \quad (2.16) \end{aligned}$$

holds for all $(x, y) \in \Delta$, where .

Proof. From Lemma 1 and using the Hölder inequality for double integrals, we have that inequality holds:

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \leq \left(\int_0^1 \int_0^1 s^p t^p ds dt \right)^{\frac{1}{p}} \\ & \quad \times \left[\frac{(x-a)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right|^q ds dt \right)^{\frac{1}{q}} \right. \\ & \quad + \frac{(x-a)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right|^q ds dt \right)^{\frac{1}{q}} \\ & \quad + \frac{(b-x)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right|^q ds dt \right)^{\frac{1}{q}} \\ & \quad \left. + \frac{(b-x)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \left(\int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right|^q ds dt \right)^{\frac{1}{q}} \right], \end{aligned} \quad (2.17)$$

for all $(x, y) \in \Delta$.

Since $\left| \frac{\partial^2 f}{\partial s \partial t} \right|^q$ is concave on the co-ordinates on Δ , so an application of (1.3) with inequalities in reversed direction, gives us the following inequalities:

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right|^q ds dt \\ & \leq \frac{1}{2} \left[\int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f \left(tx + (1-t)a, \frac{y+c}{2} \right) \right|^q dt \right. \\ & \quad \left. + \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{x+a}{2}, sy + (1-s)c \right) \right|^q ds \right] \\ & \leq \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{x+a}{2}, \frac{y+c}{2} \right) \right|^q, \end{aligned} \quad (2.18)$$

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right|^q ds dt \\ & \leq \frac{1}{2} \left[\int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f \left(tx + (1-t)a, \frac{d+y}{2} \right) \right|^q dt + \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{x+a}{2}, sy + (1-s)c \right) \right|^q ds \right] \\ & \leq \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{x+a}{2}, \frac{d+y}{2} \right) \right|^q, \end{aligned} \quad (2.19)$$

$$\begin{aligned} & \int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right|^q ds dt \\ & \leq \frac{1}{2} \left[\int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f \left(tx + (1-t)a, \frac{y+c}{2} \right) \right|^q dt + \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{b+x}{2}, sy + (1-s)c \right) \right|^q ds \right] \\ & \leq \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{b+a}{2}, \frac{y+c}{2} \right) \right|^q \end{aligned} \quad (2.20)$$

and

$$\begin{aligned}
& \int_0^1 \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right|^q ds dt \\
& \leq \frac{1}{2} \left[\int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f \left(tx + (1-t)b, \frac{d+y}{2} \right) \right|^q dt \right. \\
& \quad \left. + \int_0^1 \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{b+x}{2}, sy + (1-s)d \right) \right|^q ds \right] \\
& \leq \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{b+x}{2}, \frac{d+y}{2} \right) \right|^q. \quad (2.21)
\end{aligned}$$

By making use of (2.18)-(2.21) in (2.17), we obtain (2.16). Thus the proof of the theorem is complete. \square

Theorem 7. Let $f : \Delta \rightarrow \mathbb{R}$ be a twice partial differentiable mapping on Δ° such that $\frac{\partial^2 f}{\partial s \partial t} \in L(\Delta)$. If $\left| \frac{\partial^2 f}{\partial s \partial t} \right|^q$ is concave on the co-ordinates on Δ and $q \geq 1$, then the inequality

$$\begin{aligned}
& \left| \frac{[(b-x)^\alpha + (x-a)^\alpha][(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \\
& \leq \frac{1}{4(b-a)(d-c)} \left[(x-a)^{\alpha+1} (y-c)^{\beta+1} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{3x+a}{3}, \frac{3y+c}{3} \right) \right| \right. \\
& \quad + (x-a)^{\alpha+1} (d-y)^{\beta+1} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{3x+a}{3}, \frac{3y+d}{3} \right) \right| \\
& \quad + (b-x)^{\alpha+1} (y-c)^{\beta+1} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{3x+b}{3}, \frac{3y+c}{3} \right) \right| \\
& \quad \left. + (b-x)^{\alpha+1} (d-y)^{\beta+1} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{3x+b}{3}, \frac{3y+d}{3} \right) \right| \right], \quad (2.22)
\end{aligned}$$

holds for all $(x, y) \in \Delta$.

Proof. By the concavity of $\left| \frac{\partial^2 f}{\partial s \partial t} \right|^q$ on the co-ordinates on Δ and power mean inequality, we note that the following inequality holds:

$$\begin{aligned}
\left| \frac{\partial^2}{\partial s \partial t} f(\lambda x + (1-\lambda)y, v) \right|^q & \geq \lambda \left| \frac{\partial^2}{\partial s \partial t} f(x, v) \right|^q + (1-\lambda) \left| \frac{\partial^2}{\partial s \partial t} f(y, v) \right|^q \\
& \geq \left(\lambda \left| \frac{\partial^2}{\partial s \partial t} f(x, v) \right| + (1-\lambda) \left| \frac{\partial^2}{\partial s \partial t} f(y, v) \right| \right)^q,
\end{aligned}$$

for all $x, y \in [a, b]$, $\lambda \in [0, 1]$ and for fixed $v \in [c, d]$.

Hence,

$$\left| \frac{\partial^2}{\partial s \partial t} f(\lambda x + (1-\lambda)y, v) \right| \geq \lambda \left| \frac{\partial^2}{\partial s \partial t} f(x, v) \right| + (1-\lambda) \left| \frac{\partial^2}{\partial s \partial t} f(y, v) \right|,$$

for all $x, y \in [a, b]$, $\lambda \in [0, 1]$ and for fixed $v \in [c, d]$.

Similarly, we can show that

$$\left| \frac{\partial^2}{\partial s \partial t} f(u, \lambda z + (1-\lambda)w) \right| \geq \lambda \left| \frac{\partial^2}{\partial s \partial t} f(u, z) \right| + (1-\lambda) \left| \frac{\partial^2}{\partial s \partial t} f(u, w) \right|,$$

for all $z, w \in [c, d]$, $\lambda \in [0, 1]$ and for fixed $u \in [a, b]$, thus $\left| \frac{\partial^2 f}{\partial s \partial t} \right|$ is concave on the co-ordinates on Δ .

From Lemma 1, we have

$$\begin{aligned} & \left| \frac{[(b-x)^\alpha + (x-a)^\alpha] [(d-y)^\beta + (y-c)^\beta]}{(b-a)(d-c)} f(x, y) + A \right| \\ & \leq \frac{(x-a)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right| ds dt \\ & + \frac{(x-a)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right| ds dt \\ & + \frac{(b-x)^{\alpha+1} (y-c)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right|^q ds dt \\ & + \frac{(b-x)^{\alpha+1} (d-y)^{\beta+1}}{(b-a)(d-c)} \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right|^q ds dt, \end{aligned} \quad (2.23)$$

for all $(x, y) \in \Delta$.

Since $\left| \frac{\partial^2 f}{\partial s \partial t} \right|^q$ is concave on the co-ordinates on Δ , we have, by the Jensens's inequality for integrals, that

$$\begin{aligned} & \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right| ds dt \\ & = \int_0^1 t \left[\int_0^1 s \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)c) \right| ds \right] dt \\ & \leq \int_0^1 t \left[\left(\int_0^1 s ds \right) \left| \frac{\partial^2}{\partial s \partial t} f \left(tx + (1-t)a, \frac{\int_0^1 s (sy + (1-s)c) ds}{\int_0^1 s ds} \right) \right| \right] dt \\ & = \frac{1}{2} \int_0^1 t \left| \frac{\partial^2}{\partial s \partial t} f \left(tx + (1-t)a, \frac{3y+c}{3} \right) \right| dt \\ & \leq \frac{1}{2} \left(\int_0^1 t dt \right) \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{\int_0^1 t (tx + (1-t)a) dt}{2}, \frac{3y+c}{3} \right) \right| \\ & = \frac{1}{4} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{3x+a}{3}, \frac{3y+c}{3} \right) \right|. \end{aligned} \quad (2.24)$$

Analogously, we also have

$$\begin{aligned} & \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)a, sy + (1-s)d) \right| ds dt \\ & \leq \frac{1}{4} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{3x+a}{3}, \frac{3y+d}{3} \right) \right|, \end{aligned} \quad (2.25)$$

$$\begin{aligned} & \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)c) \right| ds dt \\ & \leq \frac{1}{4} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{3x+b}{3}, \frac{3y+c}{3} \right) \right| \quad (2.26) \end{aligned}$$

and

$$\begin{aligned} & \int_0^1 \int_0^1 st \left| \frac{\partial^2}{\partial s \partial t} f(tx + (1-t)b, sy + (1-s)d) \right| ds dt \\ & \leq \frac{1}{4} \left| \frac{\partial^2}{\partial s \partial t} f \left(\frac{3x+b}{3}, \frac{3y+d}{3} \right) \right|. \quad (2.27) \end{aligned}$$

Using (2.24)-(2.27) in (2.23), we obtain the required inequality (2.22) and hence the theorem is complete. \square

Remark 4. If we take $\alpha = \beta = 1$, in Theorem 6 and Theorem 7, we get the inequalities proved in [31, Theorem 5 and Theorem 6].

3. ACKNOWLEDGEMENT

The author would like to say thank to the Professor S. S. Dragomir, School of Engineering and Science, Victoria University, PO Box 14428 Melbourne City, MC 8001, Australia and Professor Gaston M. N'Guérékata, Morgan State University, Baltimore, Maryland, USA for their continuous encouragement and support.

REFERENCES

- [1] G. Anastassiou, M.R. Hooshmandasl, A. Ghasemi and F. Moftakharzadeh, Montogomery identities for fractional integrals and related fractional inequalities, J. Ineq. Pure and Appl. Math., 10(4) (2009), Art. 97.
- [2] M. Alomari and M. Darus, Some Ostrowski type inequalities for convex functions with applications, RGMIA **13** (1) (2010) article No. 3. Preprint.
- [3] M. Alomari, M. Darus, S.S. Dragomir, P. Cerone, Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense, Appl. Math. Lett. 23 (2010) 1071-1076.
- [4] M. Alomari, M. Darus, Some Ostrowski type inequalities for quasi-convex functions with applications to special means, RGMIA Res. Rep. Coll., 13 (2010), 2, Article 3.
- [5] M. Alomari and M. Darus, Hadamard-type inequalities for s -convex functions, International Mathematical Forum, 3 (2008), no. 40, 1965-1975.
- [6] M. Alomari and M. Darus, Co-ordinated s -convex function in the first sense with some Hadamard-type inequalities, Int. Journal Contemp. Math. Sciences, 3 (2008), no. 32, 1557-1567.
- [7] M. Alomari and M. Darus, The Hadamard's inequality for s -convex function of 2-variables on the co-ordinates, International Journal of Math. Analysis, 2 (2008), no. 13, 629-638.
- [8] M. K. Bakula and J. Pečarić, On the Jensen's inequality for convex functions on the coordinates in a rectangle from the plane, Taiwanese Journal of Math., 5, 2006, 1271-1292.
- [9] N.S. Barnett, P. Cerone, S.S. Dragomir, M.R. Pinheiro and A. Sofo, Ostrowski type inequalities for functions whose modulus of derivatives are convex and applications, RGMIA Res. Rep. Coll., 5(2) (2002), Article 1. [ONLINE: <http://rgmia.vu.edu.au/v5n2.html>]
- [10] S. Belarbi and Z. Dahmani, On some new fractional integral inequalities, J. Ineq. Pure and Appl. Math., 10(3) (2009), Art. 86.
- [11] P. Cerone and S.S. Dragomir, Ostrowski type inequalities for functions whose derivatives satisfy certain convexity assumptions, Demonstratio Math., 37 (2004), no. 2, 299–308.
- [12] S. S. Dragomir, On the Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, Taiwanese Journal of Mathematics, 5 (2001), no. 4, 775-788.
- [13] S. S. Dragomir and C. E. M. Pearce, Selected Topics on Hermite-Hadamard Type Inequalities and Applications, RGMIA (2000), Monographs. [ONLINE: http://ajmaa.org/RGMIA/monographs/hermite_hadamard.html].

- [14] S. S. Dragomir and S. Fitzpatrick, The Hadamard's inequality for s -convex functions in the second sense, *Demonstratio Math.* 32(4), (1999), 687-696.
- [15] S.S. Dragomir, On the Ostrowski's integral inequality for mappings with bounded variation and applications, *Math. Ineq. & Appl.*, 1(2) (1998).
- [16] S.S. Dragomir, The Ostrowski integral inequality for Lipschitzian mappings and applications, *Comput. Math. Appl.*, 38 (1999), 33-37.
- [17] S. S. Dragomir, S. Wang, A new inequality of Ostrowski's type in L_1 -norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. of Math.*, 28 (1997), 239-244.
- [18] S.S. Dragomir and S. Wang, A new inequality of Ostrowskis type in L_p -norm and applications to some special means and to some numerical quadrature rules, *Indian J. Math.*, 40(3) (1998), 299-304
- [19] S.S. Dragomir, P. Cerone, J. Roumeliotis, A new generalization of Ostrowski's integral inequality for mappings whose derivatives are bounded and applications in numerical integration and for special means, *Appl. Math. Lett.* 13 (2000) 19-25.
- [20] Z. Dahmani, New inequalities in fractional integrals, *International Journal of Nonlinear Scinece*, 9(4) (2010), 493-497.
- [21] Z. Dahmani, On Minkowski and Hermite-Hadamard integral inequalities via fractional integration, *Ann. Funct. Anal.* 1(1) (2010), 51-58.
- [22] Z. Dahmani, L. Tabharit, S. Taf, Some fractional integral inequalities, *Nonl. Sci. Lett. A*, 1(2) (2010), 155-160.
- [23] Z. Dahmani, L. Tabharit, S. Taf, New generalizations of Gruss inequality usin Riemann-Liouville fractional integrals, *Bull. Math. Anal. Appl.*, 2(3) (2010), 93-99.
- [24] R. Gorenflo, F. Mainardi, Fractional calculus: integral and differential equations of fractional order, Springer Verlag, Wien (1997), 223-276.
- [25] H. Hudzik and L. Maligranda, Some remarks on s -convex functions, *Aequationes Math.* 48 (1994), 100-111.
- [26] D. Y. Hwang, K. L. Tseng and G. S. Yang, Some Hadamard's inequalities for co-ordinated convex functions in a rectangle from the plane, *Taiwanese Journal of Mathematics*, 11 (2007), 63-73.
- [27] Z. Liu, Some companions of an Ostrowski type inequality and application, *J. Inequal. in Pure and Appl. Math.*, 10(2), 2009, Art. 52, 12 pp.
- [28] M. A. Latif, and M. Alomari, On Hadamard-type inequalities for h -convex functions on the co-ordinates, *International Journal of Math. Analysis*, 3 (2009), no. 33, 1645-1656.
- [29] M. A. Latif and M. Alomari, Hadamard-type inequalities for product two convex functions on the co-ordinates, *International Mathematical Forum*, 4 (2009), no. 47, 2327-2338.
- [30] M. A. latif and S. S. Dragomir, On some new inequalities for differentiable co-ordinated convex functions, *RGMIA Research. Report. Collection*, 14(2011), Article 45.
- [31] M. A. Latif, S. Hussain and S. S. Dragomir, New Ostrowski type inequalities for co-ordinated convex functions, *RGMIA Research Report Collection*, 14(2011), Article 49. [ONLINE: <http://www.ajmaa.org/RGMIA/v14.php>].
- [32] S. Miller and B. Ross, *An introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons, USA, 1993, p.2.
- [33] D.S. Mitrinović, J. E. Pečarić and A.M. Fink, *Inequalities Involving Functions and Their Integrals and Derivatives*, Kluwer Academic Publishers, Dordrecht, 1991
- [34] A. Ostrowski, *Über die Absolutabweichung einer differentiebaren Funktion von ihrem Integralmittelwert*, *Comment. Math. Helv.*, 10 (1938), 226–227. *Comp. Appl. Math.*, Vol. **22**, N. 2, 2003.
- [35] M. E. Özdemir, E. Set, and M. Z. Sarıkaya, Some new Hadamard's type inequalities for co-ordinated m -convex and (α, m) -convex functions, Accepted.
- [36] M.E. Özdemir, H. Kavurmac and E. Set, Ostrowski's type inequalities for (α, m) -convex functions, *Kyungpook Math. J.*, 50 (2010), 371-378.
- [37] B. G. Pachpatte, On an inequality of Ostrowski type in three independent variables, *J. Math.Anal. Appl.*, 249(2000), 583-591.
- [38] B. G. Pachpatte, On a new Ostrowski type inequality in two independent variables, *Tamkang J. Math.*, 32(1), (2001), 45-49
- [39] A. Rafiq, N.A. Mir and F. Ahmad, Weighted Čebyšev-Ostrowski type inequalities, *Applied Math. Mechanics (English Edition)*, 2007, 28(7), 901-906.

- [40] M.Z. Sarikaya, E. Set, H. Yaldiz and N. Başak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, Submitted.
- [41] M.Z. Sarikaya and H. Ogunmez, On new inequalities via Riemann-Liouville fractional integration, arXiv:1005.1167v1, submitted.
- [42] M. Z. Sarikaya, E. Set, M. E. Özdemir and S. S. Dragomir, New some Hadamard's type inequalities for co-ordinated convex functions, Accepted.
- [43] M. Z. Sarikaya, On the Ostrowski type integral inequality, Acta Math. Univ. Comenianae, Vol. LXXIX, 1(2010), pp. 129-134.
- [44] N. Ujević, Sharp inequalities of Simpson type and Ostrowski type, Comput. Math. Appl. 48 (2004) 145-151.
- [45] L. Zhongxue, On sharp inequalities of Simpson type and Ostrowski type in two independent variables, Comput. Math. Appl., 56 (2008) 2043-2047.

MUHAMMAD AMER LATIF
COLLEGE OF SCIENCE, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIL, HAIL 2440, SAUDI ARABIA

E-mail address: m.amer.latif@hotmail.com

SABIR HUSSAIN
UNIVERSITY OF ENGINEERING AND TECHNOLOGY, LAHORE, PAKISTAN
E-mail address: sabirhus@gmail.com