

## EXACT SOLUTIONS FOR COUPLED NONLINEAR PARTIAL DIFFERENTIAL EQUATIONS USING $G'/G$ METHOD

U. M. ABDELSALAM

ABSTRACT. ( $G'/G$ )-expansion method is examined to solve the Boiti-Leon-Pempinelli (BLP) system and the  $(2 + 1)$ -dimensional breaking soliton system. The results show that this method is a powerful tool for solving systems of nonlinear PDEs., it presents exact travelling wave solutions. The obtained solutions include rational, periodical, singular, shocklike wave and solitary wave solutions.

### 1. INTRODUCTION

Nonlinear PDEs are widely used to describe complex phenomena in various fields of sciences, such as fluid mechanics, plasma physics, astrophysics, optical fibers, solid state physics, chemical kinematics, chemical physics and geochemistry, etc., [[1]-[19]]. Studying the nonlinear waves such as soliton, breather, compacton, etc., is one of the most important problems in mathematical physics and engineering. Various mathematical methods for finding exact solutions of NLEEs have been proposed, such as tanh method [12], extended tanh method [20, 21], the symmetry method [22], sine-cosine method [23], the improved ( $G'/G$ )- expansion method [24], the ( $G'/G, 1/G$ )-expansion method [25], homogeneous balance [26], F-expansion method [27], generalized expansion method [28] and ( $G'/G$ ) method [[29]-[31]] . Recently, In [31] ( $G'/G$ )-expansion method is applied to solve ZK and CKP equations in multicomponent plasma. ( $G'/G$ )-expansion is a direct, effective and powerful method for finding analytical solutions of nonlinear partial differential equations. In [29] Wang et al. proposed the method, while Zhang et al. [30] proposed a generalized ( $G'/G$ )-expansion method to improve and extend  $G'/G$  method to solve variable coefficient and high dimensional equations. In this work the ( $G'/G$ )-expansion method with computations are performed with computer algebra system such as Mathematica to deduce many exact breather-type solutions containing rational, periodical, singular and solitary wave solutions. In this article, many exact travelling solutions are obtained for the Boiti-Leon-Pempinelli (BLP) equation and the  $(2 + 1)$ -dimensional breaking soliton equation.

---

2010 *Mathematics Subject Classification.* 35C07, 35C08, 35B10, 35C09.

*Key words and phrases.* ( $G'/G$ )-expansion method, coupled systems of PDES, exact travelling solutions.

Submitted Feb. 19, 2019. Revised July 31, 2019.

The manuscript is organised in the following fashion; In Section II, The extended  $(G'/G)$ -expansion method is described. In section III we apply extended  $(G'/G)$ -expansion method to solve the (BLP) equation and the  $(2 + 1)$ -dimensional breaking soliton equation systems. conclusions are given in section V.

## 2. DESCRIPTION OF THE METHOD

For the general NLEE

$$P(u, u_x, u_t, u_y, u_{xx}, \dots) = 0 \quad (1)$$

where  $u = u(x, y, t)$  and  $P$  is a polynomial in  $u$  and its derivatives. We seek its solutions in the form

$$u(\zeta) = \sum_{i=0}^n a_i \left( \frac{G'(\zeta)}{G(\zeta)} \right)^i, \quad (2)$$

where  $a_i$  are real constants with  $a_i \neq 0$  to be determined,  $n$  is a positive integer to be determined. The function  $G(\zeta)$  is the solution of the auxiliary linear ordinary differential equation

$$AGG'' - BGG' - C(G')^2 - E(G)^2 = 0, \quad (3)$$

where  $A, B, C$  and  $E$  are real constants to be determined, and

$$\zeta = x + y - \lambda t \quad (4)$$

where  $\lambda$  is the speed of the travelling wave.

**step 1.** Using transformation (4) we obtain an ordinary differential equation (ODE) for  $u = u(\zeta)$ :

$$E(u, u', u'', u''', \dots) = 0 \quad (5)$$

**step 2.** By balancing the highest nonlinear terms and the highest-order partial differential terms in the given NLEE we can determine  $n$ .

**step 3.** Substituting Eq. (2) and (3) into Eq. (5) and collecting coefficients of polynomial of  $\left(\frac{G'(\zeta)}{G(\zeta)}\right)$ , then setting each coefficient to zero yields a set of algebraic equations for  $a_i$  ( $i=0,1,2,\dots,n$ ),  $A, B, C, E$  and  $\lambda$ .

**step 4.** Solving the system of algebraic equations in step 2 for  $a_i, A, B, C, E$  and  $\lambda$  using Maple or Mathematica.

**step 5.** As Eq. (3) possesses the general solutions:

**Case 1.** If  $B \neq 0, \Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) > 0$ , then

$$\left( \frac{G'(\zeta)}{G(\zeta)} \right) = \frac{B}{2\Psi} + \frac{\sqrt{\Omega}}{2\Psi} \left( \frac{c_1 \sinh(\frac{\sqrt{\Omega}}{2\Psi}\zeta) + c_2 \cosh(\frac{\sqrt{\Omega}}{2\Psi}\zeta)}{c_1 \cosh(\frac{\sqrt{\Omega}}{2\Psi}\zeta) + c_2 \sinh(\frac{\sqrt{\Omega}}{2\Psi}\zeta)} \right), \quad (6)$$

**Case 2.** If  $B \neq 0, \Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) < 0$ , then

$$\left( \frac{G'(\zeta)}{G(\zeta)} \right) = \frac{B}{2\Psi} + \frac{\sqrt{-\Omega}}{2\Psi} \left( \frac{-c_1 \sin(\frac{\sqrt{-\Omega}}{2\Psi}\zeta) + c_2 \cos(\frac{\sqrt{-\Omega}}{2\Psi}\zeta)}{c_1 \cos(\frac{\sqrt{-\Omega}}{2\Psi}\zeta) + c_2 \sin(\frac{\sqrt{-\Omega}}{2\Psi}\zeta)} \right), \quad (7)$$

**Case 3.** If  $B \neq 0, \Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) = 0$ , then

$$\left( \frac{G'(\zeta)}{G(\zeta)} \right) = \frac{B}{2\Psi} + \frac{\sqrt{-\Omega}}{2\Psi} \left( \frac{c_2}{c_1 + c_2\zeta} \right), \quad (8)$$

**Case 4.** If  $B = 0$ ,  $\Psi = A - C$  and  $\Delta = \Psi E > 0$ , then

$$\left(\frac{G'(\zeta)}{G(\zeta)}\right) = \frac{\sqrt{\Delta}}{2\Psi} \left( \frac{c_1 \sinh(\frac{\sqrt{\Delta}}{2\Psi}\zeta) + c_2 \cosh(\frac{\sqrt{\Delta}}{2\Psi}\zeta)}{c_1 \cosh(\frac{\sqrt{\Delta}}{2\Psi}\zeta) + c_2 \sinh(\frac{\sqrt{\Delta}}{2\Psi}\zeta)} \right), \quad (9)$$

**Case 5.** If  $B = 0$ ,  $\Psi = A - C$  and  $\Delta = \Psi E < 0$ , then

$$\left(\frac{G'(\zeta)}{G(\zeta)}\right) = \frac{\sqrt{-\Delta}}{2\Psi} \left( \frac{-c_1 \sin(\frac{\sqrt{-\Delta}}{2\Psi}\zeta) + c_2 \cos(\frac{\sqrt{-\Delta}}{2\Psi}\zeta)}{c_1 \cos(\frac{\sqrt{-\Delta}}{2\Psi}\zeta) + c_2 \sin(\frac{\sqrt{-\Delta}}{2\Psi}\zeta)} \right), \quad (10)$$

### 3. APPLICATION OF THE METHOD

**3.1. Boiti-Leon-Pempinelli (BLP) system.** Let us consider the (2+1)-dimensional coupled Boiti-Leon-Pempinelli (BLP) system,

$$\begin{aligned} u_{ty} &= (u^2 - u_x)_{xy} + 2v_{xx}, \\ v_t &= v_{xx} + 2(uv)_x \end{aligned} \quad (11)$$

Balancing the highest derivative term with non-linear terms, hence we may assume that

$$\begin{aligned} u(x, y, t) &= k_0 + k_1\phi(\zeta) \\ v(x, y, t) &= \mu_0 + \mu_1\phi(\zeta) + \mu_2\phi^2(\zeta) \end{aligned} \quad (12)$$

where  $\zeta = x + y - \lambda t$ . Substituting Eq. (12) into Eq. (11) with aide of Eq. (3) and collecting coefficients of polynomial of  $\phi^i$  and equating them to zero, we get a system of algebraic equations for  $k_0$ ,  $k_1$ ,  $\mu_0$ ,  $\mu_1$  and  $\mu_2$

$$-\frac{2Ek_1\mu_0}{A} - \frac{BE\mu_1}{A^2} - \frac{\lambda E\mu_1}{A} - \frac{2Ek_0\mu_1}{A} - \frac{2E^2\mu_2}{A^2} = 0 \quad (13)$$

$$-\frac{2Bk_1\mu_0}{A} - \frac{B^2\mu_1}{A^2} - \frac{B\lambda\mu_0}{A} + \frac{2E\mu_0}{A} - \frac{2CE\mu_1}{A^2} - \frac{2Bk_0\mu_1}{A} - \frac{4Ek_1\mu_1}{A} - \frac{6BE\mu_2}{A^2} - \frac{2\lambda E\mu_2}{A} - \frac{4Ek_0\mu_2}{A} = 0 \quad (14)$$

$$\begin{aligned} 2k_1\mu_0 - \frac{2Ck_1\mu_0}{A} + \frac{3B\mu_1}{A} + \lambda\mu_1 - \frac{3B\lambda\mu_1}{A^2} - \frac{\lambda C\mu_1}{A} + 2k_0\mu_1 - \frac{2\lambda k_0\mu_1}{A} - \frac{4Bk_1\mu_1}{A} - \\ \frac{4B^2\mu_2}{A^2} - \frac{2B\lambda\mu_2}{A} + \frac{8E\mu_2}{A} - \frac{8\lambda E\mu_2}{A^2} - \frac{4Bk_0\mu_2}{A} - \frac{6Ek_1\mu_2}{A} = 0 \end{aligned} \quad (15)$$

$$\begin{aligned} -2\mu_1 + \frac{4C\mu_1}{A} - \frac{2C^2\mu_1}{A^2} + 4k_1\mu_1 - \frac{4Ck_1\mu_1}{A} + \frac{10B\mu_2}{A} + 2\lambda\mu_2 - \frac{10BC\mu_2}{A^2} - \\ \frac{2\lambda C\mu_2}{A} + 4k_0\mu_2 - \frac{4Ck_0\mu_2}{A} - \frac{6Bk_1\mu_2}{A} = 0 \end{aligned} \quad (16)$$

$$-6\mu_2 + \frac{12C\mu_2}{A} - \frac{6C^2\mu_2}{A^2} + 6k_1\mu_2 - \frac{6Ck_1\mu_2}{A} = 0 \quad (17)$$

Solving the last system of equations we get

$$k_0 = \frac{-B - Ac}{2A}, k_1 = \frac{A - C}{A}, \mu_2 = -k_1^2, \mu_0 = -\frac{ek_1^3}{-A + C + Ck_1}, \mu_2 = -ck_1 - 2k_0k_1 \quad (18)$$

Thus we obtain the following solutions of Eq. (12):

For  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) > 0$ , we get

$$u_1(\zeta) = \frac{B + 2Ak_0\sqrt{\Omega}\operatorname{Tanh}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right]}{2A}$$

$$v_1(\zeta) = -\frac{B^2 + 2AB(c + 2k_0) - 4A^2\mu_0 + 2(B + A(c + 2k_0))\sqrt{\Omega}\operatorname{Tanh}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right] + \Omega\operatorname{Tanh}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right]^2}{4A^2} \quad (19)$$

and

$$u_2(\zeta) = \frac{B + 2Ak_0\sqrt{\Omega}\operatorname{Coth}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right]}{2A}$$

$$v_2(\zeta) = -\frac{B^2 + 2AB(c + 2k_0) - 4A^2\mu_0 + 2(B + A(c + 2k_0))\sqrt{\Omega}\operatorname{Coth}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right] + \Omega\operatorname{Coth}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right]^2}{4A^2} \quad (20)$$

However for  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) < 0$ , we obtain periodic solutions

$$u_3(\zeta) = \frac{B + 2Ak_0\sqrt{-\Omega}\operatorname{Tan}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right]}{2A}$$

$$v_3(\zeta) = -\frac{B^2 + 2AB(c + 2k_0) - 4A^2\mu_0 + 2(B + A(c + 2k_0))\sqrt{-\Omega}\operatorname{Tan}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right] - \Omega\operatorname{Tan}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right]^2}{4A^2} \quad (21)$$

$$u_4(\zeta) = \frac{B + 2Ak_0\sqrt{-\Omega}\operatorname{Cot}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right]}{2A}$$

$$v_4(\zeta) = -\frac{B^2 + 2AB(c + 2k_0) - 4A^2\mu_0 + 2(B + A(c + 2k_0))\sqrt{-\Omega}\operatorname{Cot}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right] - \Omega\operatorname{Cot}\left[\frac{\zeta\sqrt{\Omega}}{2\psi}\right]^2}{4A^2} \quad (22)$$

For  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) = 0$ , we obtain rational solutions

$$u_5(\zeta) = k_0 + \frac{\frac{B}{2} + \frac{\psi}{\zeta}}{A}$$

$$v_5(\zeta) = \mu_0 - \frac{(c + 2k_0)(B\zeta + 2\psi)}{2A\zeta} - \frac{(B\zeta + 2\psi)^2}{4A^2\zeta^2} \quad (23)$$

For  $B = 0$ ,  $\Psi = A - C$  and  $\Delta = \Psi E > 0$ , we get

$$u_6(\zeta) = \frac{\lambda}{2} + \frac{\sqrt{\Delta}\operatorname{Tanh}\left[\frac{E\zeta}{\sqrt{\Delta}}\right]}{A}$$

$$v_6(\zeta) = \frac{\Delta\left[1 - \operatorname{Tanh}\left[\frac{E\zeta}{\sqrt{\Delta}}\right]^2\right]}{A^2} \quad (24)$$

$$\begin{aligned}
 u_7(\zeta) &= \frac{\lambda}{2} + \frac{\sqrt{\Delta} \operatorname{Coth} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]}{A} \\
 v_7(\zeta) &= \frac{-\Delta \left[ -1 + \operatorname{Coth} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right] \right]^2}{A^2}
 \end{aligned} \tag{25}$$

For  $B = 0$ ,  $\Psi = A - C$  and  $\Delta = \Psi E < 0$ , the periodic solutions

$$\begin{aligned}
 u_8(\zeta) &= \frac{-\lambda}{2} - \frac{\sqrt{-\Delta} \operatorname{Tan} \left[ \frac{E\zeta}{\sqrt{-\Delta}} \right]}{A} \\
 v_8(\zeta) &= \frac{\Delta \left[ -1 + \operatorname{Tan} \left[ \frac{E\zeta}{\sqrt{-\Delta}} \right] \right]^2}{A^2}
 \end{aligned} \tag{26}$$

$$\begin{aligned}
 u_9(\zeta) &= \frac{-\lambda}{2} - \frac{\sqrt{\Delta} \operatorname{Cot} \left[ \frac{E\zeta}{\sqrt{-\Delta}} \right]}{A} \\
 v_9(\zeta) &= \frac{\Delta \left[ -1 + \operatorname{Cot} \left[ \frac{E\zeta}{\sqrt{-\Delta}} \right] \right]^2}{A^2}
 \end{aligned} \tag{27}$$

**3.2. The (2 + 1)-dimensional breaking soliton system.** We consider the (2 + 1)-dimensional breaking soliton system,

$$\begin{aligned}
 u_t + \alpha u_{xxy} + 4\alpha(uv)_x &= 0, \\
 u_y &= v_x
 \end{aligned} \tag{28}$$

Balancing the highest derivative term with non-linear terms, hence we may assume that

$$\begin{aligned}
 u(x, y, t) &= k_0 + k_1\phi(\zeta) + k_2\phi^2(\zeta) \\
 v(x, y, t) &= \mu_0 + \mu_1\phi(\zeta) + \mu_2\phi^2(\zeta)
 \end{aligned} \tag{29}$$

where  $\zeta = x + y - \lambda t$ . By the same way we Substitute Eq. (29) into Eq. (28) and collecting coefficients of polynomial of  $\phi^i$  and equating them to zero, we get a system of algebraic equations for  $k_0, k_1, k_2, \mu_0, \mu_1$  and  $\mu_2$ , by solving this system of equations we get the first set,

$$k_1 = \frac{3B(A - C)}{2A^2}, k_2 = \frac{3(A - C)^2}{2A^2}, \mu_1 = k_1, \mu_2 = k_2 \tag{30}$$

and the second set,

$$k_1 = 0, k_2 = -\frac{3(A^2 - 2AC + C^2)}{2A^2}, \mu_2 = k_2, \mu_0 = \frac{-3Ak_0 + 3Ck_0 - 4ek_2}{3(A - C)}, \mu_1 = 0 \tag{31}$$

Thus for the first set, we obtain the following solutions of Eq. (6):

For  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) > 0$ , we get

$$\begin{aligned} u_1(\zeta) &= k_0 + \frac{3\Omega \operatorname{Tanh} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \\ v_1(\zeta) &= \mu_0 + \frac{3\Omega \operatorname{Tanh} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \end{aligned} \quad (32)$$

and

$$\begin{aligned} u_2(\zeta) &= k_0 + \frac{3\Omega \operatorname{Coth} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \\ v_2(\zeta) &= \mu_0 + \frac{3\Omega \operatorname{Coth} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \end{aligned} \quad (33)$$

However for  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) < 0$ , we obtain periodic solutions

$$\begin{aligned} u_3(\zeta) &= k_0 - \frac{3\Omega \operatorname{Tan} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \\ v_3(\zeta) &= \mu_0 - \frac{3\Omega \operatorname{Tan} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \end{aligned} \quad (34)$$

$$\begin{aligned} u_4(\zeta) &= k_0 - \frac{3\Omega \operatorname{Cot} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \\ v_4(\zeta) &= \mu_0 - \frac{3\Omega \operatorname{Cot} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \end{aligned} \quad (35)$$

For  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) = 0$ , we obtain rational solutions

$$\begin{aligned} u_5(\zeta) &= k_0 + \frac{3(3B^2\zeta^2 + 8B\zeta\psi + 4\psi^2)}{8A^2\zeta^2} \\ v_5(\zeta) &= \mu_0 + \frac{3(3B^2\zeta^2 + 8B\zeta\psi + 4\psi^2)}{8A^2\zeta^2} \end{aligned} \quad (36)$$

For  $B = 0$ ,  $\Psi = A - C$  and  $\Delta = \Psi E > 0$ , we get

$$\begin{aligned} u_6(\zeta) &= k_0 + \frac{3\Delta \operatorname{Tanh} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]^2}{2A^2} \\ v_6(\zeta) &= \mu_0 + \frac{3\Delta \operatorname{Tanh} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]^2}{2A^2} \end{aligned} \quad (37)$$

$$\begin{aligned}
 u_7(\zeta) &= k_0 + \frac{3\Delta \operatorname{Coth} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]^2}{2A^2} \\
 v_7(\zeta) &= \mu_0 + \frac{3\Delta \operatorname{Coth} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]^2}{2A^2}
 \end{aligned} \tag{38}$$

For  $B = 0$ ,  $\Psi = A - C$  and  $\Delta = \Psi E < 0$ , the periodic solutions

$$\begin{aligned}
 u_8(\zeta) &= k_0 - \frac{3\Delta \operatorname{Tan} \left[ \frac{\sqrt{-\Delta}\zeta}{\Psi} \right]^2}{2A^2} \\
 v_8(\zeta) &= \mu_0 - \frac{3\Delta \operatorname{Tan} \left[ \frac{\sqrt{-\Delta}\zeta}{\Psi} \right]^2}{2A^2}
 \end{aligned} \tag{39}$$

$$\begin{aligned}
 u_9(\zeta) &= k_0 - \frac{3\Delta \operatorname{Cot} \left[ \frac{\sqrt{-\Delta}\zeta}{\Psi} \right]^2}{2A^2} \\
 v_9(\zeta) &= \mu_0 - \frac{3\Delta \operatorname{Cot} \left[ \frac{\sqrt{-\Delta}\zeta}{\Psi} \right]^2}{2A^2}
 \end{aligned} \tag{40}$$

For the second set,

For  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) > 0$ , we get

$$\begin{aligned}
 u_1(\zeta) &= k_0 - \frac{3\Omega \operatorname{Tanh} \left[ \frac{\zeta\sqrt{\Omega}}{2\Psi} \right]^2}{8A^2} \\
 v_1(\zeta) &= -k_0 + \frac{2\Delta}{A^2} - \frac{3\Omega \operatorname{Tanh} \left[ \frac{\zeta\sqrt{\Omega}}{2\Psi} \right]^2}{8A^2}
 \end{aligned} \tag{41}$$

and

$$\begin{aligned}
 u_2(\zeta) &= k_0 - \frac{3\Omega \operatorname{Coth} \left[ \frac{\zeta\sqrt{\Omega}}{2\Psi} \right]^2}{8A^2} \\
 v_2(\zeta) &= -k_0 + \frac{2\Delta}{A^2} - \frac{3\Omega \operatorname{Coth} \left[ \frac{\zeta\sqrt{\Omega}}{2\Psi} \right]^2}{8A^2}
 \end{aligned} \tag{42}$$

However for  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) < 0$ , we obtain periodic solutions

$$\begin{aligned}
 u_3(\zeta) &= k_0 + \frac{3\Omega \operatorname{Tan} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2} \\
 v_3(\zeta) &= -k_0 + \frac{2e\psi}{A^2} + \frac{3\Omega \operatorname{Tan} \left[ \frac{\zeta\sqrt{\Omega}}{2\psi} \right]^2}{8A^2}
 \end{aligned} \tag{43}$$

$$\begin{aligned}
 u_4(\zeta) &= k_0 + \frac{3\Omega \text{Cot} \left[ \frac{\zeta\sqrt{\Omega}}{2\Psi} \right]^2}{8A^2} \\
 v_4(\zeta) &= -k_0 + \frac{2\Delta}{A^2} + \frac{3\Omega \text{Cot} \left[ \frac{\zeta\sqrt{\Omega}}{2\Psi} \right]^2}{8A^2}
 \end{aligned} \tag{44}$$

For  $B \neq 0$ ,  $\Psi = A - C$  and  $\Omega = B^2 + 4E(A - C) = 0$ , we obtain rational solutions

$$\begin{aligned}
 u_5(\zeta) &= k_0 \\
 v_5(\zeta) &= \frac{2E}{A} - \frac{2CE}{A^2} - k_0
 \end{aligned} \tag{45}$$

For  $B = 0$ ,  $\Psi = A - C$  and  $\Delta = \Psi E > 0$ , we get

$$\begin{aligned}
 u_6(\zeta) &= k_0 - \frac{3\Delta \text{Tanh} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]^2}{2A^2} \\
 v_6(\zeta) &= -\frac{-4\Delta + 2A^2k_0 + 3\Delta \text{Tanh} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]^2}{2A^2}
 \end{aligned} \tag{46}$$

$$\begin{aligned}
 u_7(\zeta) &= k_0 - \frac{3\Delta \text{Coth} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]^2}{2A^2} \\
 v_7(\zeta) &= -\frac{-4\Delta + 2A^2k_0 + 3\Delta \text{Coth} \left[ \frac{E\zeta}{\sqrt{\Delta}} \right]^2}{2A^2}
 \end{aligned} \tag{47}$$

For  $B = 0$ ,  $\Psi = A - C$  and  $\Delta = \Psi E < 0$ , the periodic solutions

$$\begin{aligned}
 u_8(\zeta) &= k_0 + \frac{3\Delta \text{Tan} \left[ \frac{\sqrt{-\Delta}\zeta}{\Psi} \right]^2}{2A^2} \\
 v_8(\zeta) &= -\frac{-4\Delta + 2A^2k_0 + 3\Delta \text{Tan} \left[ \frac{\sqrt{-\Delta}\zeta}{\Psi} \right]^2}{2A^2}
 \end{aligned} \tag{48}$$

$$\begin{aligned}
 u_9(\zeta) &= k_0 + \frac{3\Delta \text{Cot} \left[ \frac{\sqrt{-\Delta}\zeta}{\Psi} \right]^2}{2A^2} \\
 v_9(\zeta) &= -\frac{-4\Delta + 2A^2k_0 + 3\Delta \text{Cot} \left[ \frac{\sqrt{-\Delta}\zeta}{\Psi} \right]^2}{2A^2}
 \end{aligned} \tag{49}$$

#### 4. CONCLUSION

In this article,  $(G'/G)$ -expansion method was applied to give the traveling wave solutions of two Coupled  $(2 + 1)$ -Dimensional Equations, the Boiti-Leon-Pempinelli (BLP) equation and the  $(2 + 1)$ -dimensional breaking soliton equation, The  $(G'/G)$ -expansion method examined for investigating the rogue wave solutions for  $(2+1)$  dimensional real field NLEEs.

$(G'/G)$ -expansion method gives different classes of solutions. These solutions include many types like rational, periodical, soliton solutions, etc. For example,




FIGURE 1. Three-dimensional profile of the periodic solution.




FIGURE 2. Three-dimensional profile of the explosive/blowup pulse.




FIGURE 3. Three-dimensional profile of the soliton solution..

solutions (26) and (34) are examples exhibiting the sinusoidal-type periodical solutions, which develop a singularity at a finite point, i.e., for any fixed  $t = t_0$  there




FIGURE 4. Three-dimensional profile of the shocklike wave solution.

exists a value of  $a_0$  at which these solutions blow up (see figure 1). Solution  $u_7$  in (25) is in the form of explosive/blow-up solutions as depicted in figure 2 where we can see that the potential  $u$  goes to infinity at a finite point. The explosive excitation clear that there is an instability in the system, this instability can be produced due to the effect of the nonlinearity which depend on some physical parameters in the system.

Solution (23) represents the rational-type solutions, the rational solution may be a discrete joint union of manifolds. The solutions  $v_6$  in (24) is a soliton wave solution (see figure 3), the soliton wave solution produced from the balance between weak nonlinearity and dispersion and depends on the physical parameters, these physical parameters can decrease and increase the nonlinearity or the dispersion which effect on the hight and width of the soliton waves, while we obtained shocklike wave solutions like  $u_6$  in (24) as depicted in figure 4. The shocklike wave profile may change according to a variety of physical parameters. Solitons and shocklike wave solutions are very important in studying fluids and plasma physics

#### REFERENCES

- [1] A.M. Wazwaz (2009), Partial Differential Equations and Solitary Waves Theory, Springer.
- [2] M. Antonova, A. Biswas, Adiabatic parameter dynamics of perturbed solitary waves, Communications in Nonlinear Science and Numerical Simulation. Volume 14, Issue 3, 734-748. (2009).
- [3] U.M. Abdelsalam, WM Moslem, AH Khater, PK Shukla, Solitary and freak waves in a dusty plasma with negative ions, Physics of Plasmas 18 (2011), 092305.
- [4] A.G. Johnpillai, C.M. Khalique, A. Biswas, Exact solutions of KdV equation with time-dependent coefficients, Applied Mathematics and Computation. Volume 216, Issue 10, 3114-3119. (2010).
- [5] A. Biswas, M.S. Ismail, 1-Soliton solution of the coupled KdV equation and Gear-Grimshaw model, Applied Mathematics and Computation. Volume 216, Issue 12, 3662-3670. (2010).
- [6] U.M. Abdelsalam, M.S. Zobaer, Exact Traveling Wave Solutions of Further Modified Korteweg-De Vries Equation in Multicomponent Plasma, Iranian Journal of Science and Technology, Transactions A: Science, 1-8
- [7] H. Triki, A. Biswas, Soliton solutions for a generalized fifth-order KdV equation with t-dependent coefficients, Waves in Random and Complex Media. Volume 21, Number 1, 151-160. (2011).
- [8] A.M.A. El-Sayed, M. Gaber (2006), The Adomian decomposition method for solving partial differential equations of fractal order in finite domains, Physics Letters A 359 175-182.

- [9] A. G. Johnpillai, C. M. Khalique and Anjan Biswas, Exact solutions of the mKdV equation with time-dependent coefficients, *Mathematical Communications*. Volume 16, Number 2, 509-518. (2011).
- [10] U.M. Abdelsalam, S. Ali, I. Kourakis, Nonlinear electrostatic excitations of charged dust in degenerate ultra-dense quantum dusty plasmas, *Physics of Plasmas* 19 (2012), 062107
- [11] R. Sabry, W.M. Moslem, F. Haas, S. Ali, P.K. Shukla, Nonlinear structures: Explosive, soliton, and shock in a quantum electron-positron-ion magnetoplasma, *Physics of Plasmas* 15 (12), 122308 (2008)
- [12] Moslem W.M., Sabry R., Abdelsalam U.M., Kourakis I., Shukla P.K. (2009), Solitary and blow-up electrostatic excitations in rotating magnetized electron-positron-ion plasmas, *New Journal of Physics* 11 033028.
- [13] Houria Triki, Daniela Milovic, T. Hayat, Omar. M. Aldossary and Anjan Biswas, Topological Solitons solutions of the (2+1)-dimensional KdV equation with power law nonlinearity and time-dependent coefficients, *International Journal of Nonlinear Sciences and Numerical Simulation*. Volume 12, Issues 1-8, 45-50. (2011).
- [14] Houria Triki, Abdul H. Kara, Ali Bhrawy and Anjan Biswas, Soliton Solution and Conservation Law of Gear-Grimshaw Model for Shallow Water Waves, *Acta Physica Polonica A*. Volume 125, Number 5, 1099-1106. (2014).
- [15] Ali H. Bhrawy, M. A. Abdelkawy, S. Kumar, S. Johnson and Anjan Biswas, Soliton solutions of the quantum Zakharov-Kuznetsov equation which arises in quantum magneto-plasmas, *Indian Journal of Physics*. Volume 87, Issue 5, 455-463. (2013).
- [16] E. V. Krishnan, Sachin Kumar and Anjan Biswas, Solitons and other nonlinear waves of the Boussinesq equation. Volume 70, Number 2, 1213-1221. (2012).
- [17] P. Razborova, H. Triki and A. Biswas, Perturbation of dispersive shallow water waves, *Ocean Engineering*. Volume 63, 1-7. (2013).
- [18] U.M. Abdelsalam, M.S. Zobaer, Exact Traveling Wave Solutions of KdV Equation for DAWs in Superthermal Plasma, *Revista Mexicana de Astronomia y Astrofisica* 54, 363-373
- [19] U.M. Abdelsalam, F.M. Allehiany, Different Nonlinear Solutions of KP Equation in Dusty Plasmas, *Arabian Journal for Science and Engineering* 43 (1), 399-406
- [20] E. Fan, Extended tanh-function method and its applications to nonlinear equations, *Phys. Lett.A*. 277 212-218 (2000).
- [21] U.M. Abdelsalam, Exact travelling solutions of two coupled (2+ 1)-Dimensional Equations, *J. Egypt. Math. Soc.* 25 (2017), 125-128.
- [22] M.H.M. Moussa, R.M. El-Shiekh, Similarity reduction and similarity solutions of Zabolotskay-Khoklov equation with dissipative term via symmetry method, *Physica A* 371 (2006) 325-335.
- [23] A.M. Wazwaz, The sine-cosine method for obtaining solutions with compact and noncompact structures, *Applied Mathematics and Computation*. 159 559-576 (2004).
- [24] E.M.E. Zayed, Y.A. Amer, R.M.A. Shohib, Exact traveling wave solutions for nonlinear fractional partial differential equations using the improved  $G'/G$ -expansion method, *Int. J. Eng. Appl. Sci.* 7 (2014) 18-31.
- [25] E. M. E. Zayed and K. A. E. Alurr., On solving two higher-order nonlinear PDEs describing the propagation of optical pulses in optic fibers using the  $(G'/G, 1/G)$ -expansion method, *Ric. mat.*, 64 (2015) 167-194.
- [26] U.M. Abdelsalam, F.M. Allehiany, W.M. Moslem, and S.K. El-Labany, (2016), Nonlinear structures for extended Korteweg-de Vries equation in multicomponent plasma, *Pramana J. Phys.* 86 581-597; U. M. Abdelsalam, F. M. Allehiany, Different Nonlinear Solutions of KP Equation in Dusty Plasmas, *Arab J Sci Eng.* DOI 10.1007/s13369-017-2829-z.
- [27] G. Cai, Q. Wang, and J. Huang, *Inter. J. Nonl. Sci.*, A modified F-expansion method for solving breaking soliton equation 2 122 (2006).
- [28] R. Sabry, M.A. Zahran and E. Fan, A new generalized expansion method and its application in finding explicit exact solutions for a generalized variable coefficients KdV equation, *Phys. Lett. A*. 326 326-93 (2004).
- [29] M.L. Wang, X. Li, J. Zhang, The  $G'/G$ - expansion method and evolution equations in mathematical physics, *Phys. Lett. A*. 372 417-421 (2008).
- [30] J. Zhang, X. Wei and Y. Lu, A generalized  $(G'/G)$ -expansion method and its applications, *Phys. Lett. A*, **372**, 3653 (2008).

- [31] U. M. Abdelsalam and M. Selim M. (2013), Ion-acoustic waves in a degenerate multicomponent magnetoplasma, *J. Plasma Phy.* **79** 163 ; M. Selim, U.M. Abdelsalam (2014), Propagation of cylindrical acoustic waves in dusty plasma with positive dust, *Astrophysics and Space Science* 353 (2), 535-542

U. M. ABDELSALAM

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, FAYOUM UNIVERSITY, EGYPT.

*Email address:* `usama.ahmad@rub.de`, `maths_us@hotmail.com`