

COMPARING THE INTEGRAL MEANS FOR FUNCTIONS OF TWO VARIABLES WITH BOUNDED VARIATION

HÜSEYIN BUDAK, MEHMET ZEKI SARIKAYA

ABSTRACT. In this paper, we obtain an inequality for difference of the integral means using functions of two variables with bounded variation. An application to probability density functions is also given.

1. INTRODUCTION

Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable mapping on (a, b) whose derivative $f' : (a, b) \rightarrow \mathbb{R}$ is bounded on (a, b) , i.e. $\|f'\|_{\infty} := \sup_{t \in (a, b)} |f'(t)| < \infty$. Then we have the inequality

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_{\infty}, \quad (1)$$

for all $x \in [a, b]$ [23]. The constant $\frac{1}{4}$ is the best possible. This inequality is well known in the literature as the *Ostrowski inequality*.

In [15], Dragomir proved following Ostrowski type inequalities related functions of bounded variation:

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a mapping of bounded variation on $[a, b]$. Then*

$$\left| \int_a^b f(t) dt - (b-a) f(x) \right| \leq \left[\frac{1}{2} (b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b (f)$$

holds for all $x \in [a, b]$. The constant $\frac{1}{2}$ is the best possible.

Hwang and Dragomir gave the following inequality in [21].

2010 *Mathematics Subject Classification.* 26D15, 26B30, 26D10, 41A55.

Key words and phrases. Ostrowski type inequalities, Bounded variation, Riemann-Stieltjes integral, Probability density functions.

Submitted July 1, 2015.

Theorem 2. Let $f : [a, b] \rightarrow \mathbb{R}$ be a mapping with bounded variation on $[a, b]$. Then, for $a \leq x < y \leq b$, we have the inequality

$$\begin{aligned} & \left| \frac{1}{(b-a)} \int_a^b f(s) ds - \frac{1}{(y-x)} \int_x^y f(s) ds \right| \\ & \leq \frac{x-a}{b-a} \bigvee_a^x (f) + \max \left\{ \frac{x-a}{b-a}, \frac{b-y}{b-a} \right\} \bigvee_x^y (f) + \frac{b-y}{b-a} \bigvee_y^b (f) \end{aligned} \quad (2)$$

The inequality (2) is sharp.

Definition 1. (Vitali-Lebesgue-Fréchet-de la Vallée Poussin)[12]. We introduce the notation

$$\Delta_{11}f(x_i, y_j) = f(x_{i+1}, y_{j+1}) - f(x_{i+1}, y_j) - f(x_i, y_{j+1}) + f(x_i, y_j),$$

then, the function $f(x, y)$ is said to be of bounded variation if the sum

$$\sum_{i=0, j=0}^{m-1, n-1} |\Delta_{11}f(x_i, y_j)|$$

is bounded for all nets.

Therefore, one can define the concept of total variation of a function of two variables, as follows:

Let f be of bounded variation on $Q = [a, b] \times [c, d]$, and let $\sum(P)$ denote the sum $\sum_{i=1}^n \sum_{j=1}^m |\Delta_{11}f(x_i, y_j)|$ corresponding to the partition P of Q . The number

$$\bigvee_Q (f) := \bigvee_c^d \bigvee_a^b (f) := \sup \left\{ \sum(P) : P \in P(Q) \right\},$$

is called the total variation of f on Q . Here $P([a, b])$ denotes the family of partitions of $[a, b]$.

In [20], Jawarneh and Noorani gave the following Lemmas for double Riemann-Stieltjes integral:

Lemma 1. (Integrating by parts) If $f \in RS(\alpha)$ on Q , then $\alpha \in RS(f)$ on Q , and we have

$$\begin{aligned} & \int_c^d \int_a^b f(t, s) d_t d_s \alpha(t, s) + \int_c^d \int_a^b \alpha(t, s) d_t d_s f(t, s) \\ & = f(b, d) \alpha(b, d) - f(b, c) \alpha(b, c) - f(a, d) \alpha(a, d) + f(a, c) \alpha(a, c). \end{aligned} \quad (3)$$

Lemma 2. Assume that $g \in RS(\alpha)$ on Q and α is of bounded variation on Q , then

$$\left| \int_c^d \int_a^b g(x, y) d_x d_y \alpha(x, y) \right| \leq \sup_{(x,y) \in Q} |g(x, y)| \bigvee_Q (\alpha). \quad (4)$$

In [20], Jawarneh and Noorani proved the following Ostrowski type inequality or functions of two variables with bounded variation:

Theorem 3. Let $f : Q \rightarrow \mathbb{R}$ be mapping of bounded variation on Q . Then for all $(x, y) \in Q$, we have inequality

$$\begin{aligned} & \left| (b-a)(d-c)f(x,y) - \int_c^d \int_a^b f(t,s) dt ds \right| \\ & \leq \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \\ & \quad \times \left[\frac{1}{2}(d-c) + \left| y - \frac{c+d}{2} \right| \right] \bigvee_Q (f) \end{aligned} \quad (5)$$

where $\bigvee_Q (f)$ denotes the total (double) variation of f on Q .

For more information and recent developments on integral inequalities for mappings of bounded variation (single variable and two variables), please refer to ([1]-[11], [13]-[22],[24]-[29]).

In this paper, we compare the integral means using functions of two variables with bounded variation. A application to probability density functions is also given.

2. MAIN RESULTS

First, we give the following notations to simplify presentetion of some intervals.

$$Q_1 = [a, x_1] \times [c, y_1], \quad Q_2 = [a, x_1] \times [y_1, y_2], \quad Q_3 = [a, x_1] \times [y_2, d],$$

$$Q_4 = [x_1, x_2] \times [c, y_1], \quad Q_5 = [x_1, x_2] \times [y_1, y_2], \quad Q_6 = [x_1, x_2] \times [y_2, d],$$

$$Q_7 = [x_2, b] \times [c, y_1], \quad Q_8 = [x_1, x_2] \times [y_1, y_2], \quad Q_9 = [x_2, b] \times [y_2, d].$$

Theorem 4. If the function $f : Q = [a, b] \times [c, d] \rightarrow R$ is of bounded variation on Q , then, for $a \leq x_1 < x_2 \leq b$ and $c \leq y_1 < y_2 \leq d$, we have the inequality

$$\begin{aligned} & \left| \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(t,s) ds dt - \frac{1}{(b-a)(y_2-y_1)} \int_a^b \int_{y_1}^{y_2} f(t,s) ds dt \right. \\ & \quad \left. - \frac{1}{(x_2-x_1)(d-c)} \int_{x_1}^{x_2} \int_c^d f(t,s) ds dt + \frac{1}{(x_2-x_1)(y_2-y_1)} \int_{x_1}^{x_2} \int_{y_1}^{y_2} f(t,s) ds dt \right| \\ & \leq \frac{[(b-a)-(x_2-x_1)][(d-c)-(y_2-y_1)]}{(b-a)(d-c)} \left[\frac{1}{2} + \left| \frac{\frac{a+b}{2} - \frac{x_2+x_1}{2}}{(b-a)-(x_2-x_1)} \right| \right] \\ & \quad \times \left[\frac{1}{2} + \left| \frac{\frac{d+c}{2} - \frac{y_2+y_1}{2}}{(d-c)-(y_2-y_1)} \right| \right] \bigvee_a^b \bigvee_c^d (f). \end{aligned} \quad (6)$$

where $\bigvee_a^b \bigvee_c^d (f)$ denotes the total (double) variation of f on Q .

Proof. First, we define the mappings $K_{x_1, x_2}(t)$ and $L_{y_1, y_2}(s)$ by

$$K_{x_1, x_2}(t) = \begin{cases} \frac{a-t}{b-a}, & \text{if } t \in [a, x_1] \\ \frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a}, & \text{if } t \in (x_1, x_2] \\ \frac{b-t}{b-a}, & \text{if } t \in (x_2, b] \end{cases}$$

and

$$L_{y_1, y_2}(s) = \begin{cases} \frac{c-s}{d-c}, & \text{if } s \in [c, y_1] \\ \frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c}, & \text{if } s \in (y_1, y_2] \\ \frac{d-s}{d-c}, & \text{if } s \in (y_2, d] \end{cases} .$$

Using the kernels $K_{x_1, x_2}(t)$ and $L_{y_1, y_2}(s)$, we have

$$\begin{aligned} & \int_a^b \int_c^d K_{x_1, x_2}(t) L_{y_1, y_2}(s) d_s d_t f(t, s) \\ &= \int_a^{x_1} \int_c^{y_1} \frac{a-t}{b-a} \frac{c-s}{d-c} d_s d_t f(t, s) + \int_a^{x_1} \int_{y_1}^{y_2} \frac{a-t}{b-a} \left[\frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c} \right] d_s d_t f(t, s) \\ &+ \int_a^{x_1} \int_{y_2}^d \frac{a-t}{b-a} \frac{d-s}{d-c} d_s d_t f(t, s) + \int_{x_1}^{x_2} \int_c^{y_1} \left[\frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a} \right] \frac{c-s}{d-c} d_s d_t f(t, s) \\ &+ \int_{x_1}^{x_2} \int_{y_1}^{y_2} \left[\frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a} \right] \left[\frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c} \right] d_s d_t f(t, s) \\ &+ \int_{x_1}^{x_2} \int_{y_2}^d \left[\frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a} \right] \frac{d-s}{d-c} d_s d_t f(t, s) + \int_{x_2}^b \int_c^{y_1} \frac{b-t}{b-a} \frac{c-s}{d-c} d_s d_t f(t, s) \\ &+ \int_{x_2}^b \int_{y_1}^{y_2} \frac{b-t}{b-a} \left[\frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c} \right] d_s d_t f(t, s) + \int_{x_2}^b \int_{y_2}^d \frac{b-t}{b-a} \frac{d-s}{d-c} d_s d_t f(t, s) \\ &= K_1 + K_2 + \dots + K_9. \end{aligned} \tag{7}$$

Integrating the by parts, we have

$$\begin{aligned} K_1 &= \int_a^{x_1} \int_c^{y_1} \frac{a-t}{b-a} \frac{c-s}{d-c} d_s d_t f(t, s) \\ &= \frac{1}{(b-a)(d-c)} \left[(a-x_1)(c-y_1) f(x_1, y_1) - \int_a^{x_1} \int_c^{y_1} f(t, s) ds dt \right]. \end{aligned} \tag{8}$$

Similarly, we have

$$K_2 = \int_a^{x_1} \int_{y_1}^{y_2} \frac{a-t}{b-a} \left[\frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c} \right] d_s d_t f(t, s) \quad (9)$$

$$\begin{aligned} &= \frac{a-x_1}{b-a} f(x_1, y_2) + \frac{1}{(b-a)(y_2-y_1)} \int_a^{x_1} \int_{y_1}^{y_2} f(t, s) ds dt \\ &\quad + \frac{1}{(b-a)(d-c)} [(a-x_1)(c-y_2)f(x_1, y_2) \\ &\quad - (a-x_1)(c-y_1)f(x_1, y_1) - \int_a^{x_1} \int_{y_1}^{y_2} f(t, s) ds dt], \end{aligned}$$

$$K_3 = \int_a^{x_1} \int_{y_2}^d \frac{a-t}{b-a} \frac{d-s}{d-c} d_s d_t f(t, s) \quad (10)$$

$$= \frac{1}{(b-a)(d-c)} \left[-(a-x_1)(d-y_2)f(x_1, y_2) - \int_a^{x_1} \int_{y_2}^d f(t, s) ds dt \right],$$

$$K_4 = \int_{x_1}^{x_2} \int_c^{y_1} \left[\frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a} \right] \frac{c-s}{d-c} d_s d_t f(t, s) \quad (11)$$

$$\begin{aligned} &= \frac{c-y_1}{d-c} f(x_2, y_1) + \frac{1}{(x_2-x_1)(d-c)} \int_{x_1}^{x_2} \int_c^{y_1} f(t, s) ds dt \\ &\quad + \frac{1}{(b-a)(d-c)} [(a-x_2)(c-y_1)f(x_2, y_1) \\ &\quad - (a-x_1)(c-y_1)f(x_1, y_1) - \int_{x_1}^{x_2} \int_c^{y_1} f(t, s) ds dt], \end{aligned}$$

(12)

$$\begin{aligned} K_5 &= \int_{x_1}^{x_2} \int_{y_1}^{y_2} \left[\frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a} \right] \left[\frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c} \right] d_s d_t f(t, s) \\ &= f(x_2, y_2) - \frac{1}{(x_2-x_1)(y_2-y_1)} \int_{x_1}^{x_2} \int_{y_1}^{y_2} f(t, s) ds dt \end{aligned}$$

$$\begin{aligned}
& + \frac{c - y_2}{d - c} f(x_2, y_2) - \frac{c - y_1}{d - c} f(x_2, y_1) + \frac{1}{(x_2 - x_1)(d - c)} \int_{x_1}^{x_2} \int_{y_1}^{y_2} f(t, s) ds dt \\
& + \frac{a - x_2}{b - a} f(x_2, y_2) - \frac{a - x_1}{b - a} f(x_1, y_2) + \frac{1}{(b - a)(y_2 - y_1)} \int_{x_1}^a \int_{y_1}^{y_2} f(t, s) ds dt \\
& + \frac{1}{(b - a)(d - c)} [(a - x_2)(c - y_2)f(x_2, y_2) - (a - x_1)(c - y_2)f(x_1, y_2) \\
& - (a - x_2)(c - y_1)f(x_2, y_1) + (a - x_1)(c - y_1)f(x_1, y_1) - \int_{x_1}^{x_2} \int_{y_1}^{y_2} f(t, s) ds dt],
\end{aligned}$$

$$\begin{aligned}
K_6 &= \int_{x_1}^{x_2} \int_{y_2}^d \left[\frac{t - x_1}{x_2 - x_1} + \frac{a - t}{b - a} \right] \frac{d - s}{d - c} d_s d_t f(t, s) \\
&= -\frac{d - y_2}{d - c} f(x_2, y_2) + \frac{1}{(x_2 - x_1)(d - c)} \int_{x_1}^{x_2} \int_{y_2}^d f(t, s) ds dt \\
&\quad + \frac{1}{(b - a)(d - c)} [-(a - x_2)(d - y_2)f(x_2, y_2) \\
&\quad + (a - x_1)(d - y_2)f(x_1, y_2) - \int_{x_1}^{x_2} \int_{y_2}^d f(t, s) ds dt],
\end{aligned} \tag{13}$$

$$\begin{aligned}
K_7 &= \int_{x_2}^b \int_c^{y_1} \frac{b - t}{b - a} \frac{c - s}{d - c} d_s d_t f(t, s) \\
&= \frac{1}{(b - a)(d - c)} \left[-(b - x_2)(c - y_1)f(x_2, y_1) - \int_{x_2}^b \int_c^{y_1} f(t, s) ds dt \right],
\end{aligned} \tag{14}$$

$$\begin{aligned}
K_8 &= \int_{x_2}^b \int_{y_1}^{y_2} \frac{b - t}{b - a} \left[\frac{s - y_1}{y_2 - y_1} + \frac{c - s}{d - c} \right] d_s d_t f(t, s) \\
&= -\frac{b - x_2}{b - a} f(x_2, y_2) + \frac{1}{(b - a)(y_2 - y_1)} \int_{x_2}^b \int_{y_1}^{y_2} f(t, s) ds dt
\end{aligned} \tag{15}$$

$$\begin{aligned}
& + \frac{1}{(b-a)(d-c)} [- (b-x_2)(c-y_2) f(x_2, y_2) \\
& + (b-x_2)(c-y_1) f(x_2, y_1) - \int_{x_2}^b \int_{y_1}^{y_2} f(t, s) ds dt]
\end{aligned}$$

and

$$\begin{aligned}
K_9 &= \int_{x_2}^b \int_{y_2}^d \frac{b-t}{b-a} \frac{d-s}{d-c} d_s d_t f(t, s) \\
&= \frac{1}{(b-a)(d-c)} \left[(b-x_2)(d-y_2) f(x_2, y_2) - \int_{x_2}^b \int_{y_2}^d f(t, s) ds dt \right].
\end{aligned} \tag{16}$$

If we add the equalities (8)-(16) in (7), then we have

$$\begin{aligned}
& \int_a^b \int_c^d K_{x_1, x_2}(t) L_{y_1, y_2}(s) d_s d_t f(t, s) \\
&= \frac{1}{(b-a)(y_2-y_1)} \int_a^b \int_{y_1}^{y_2} f(t, s) ds dt + \frac{1}{(x_2-x_1)(d-c)} \int_{x_1}^{x_2} \int_c^d f(t, s) ds dt \\
&\quad - \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(t, s) ds dt - \frac{1}{(x_2-x_1)(y_2-y_1)} \int_{x_1}^{x_2} \int_{y_1}^{y_2} f(t, s) ds dt.
\end{aligned}$$

On the other hand, using Lemma 2, we get

$$\begin{aligned}
& \left| \int_a^b \int_c^d K_{x_1, x_2}(t) L_{y_1, y_2}(s) d_s d_t f(t, s) \right| \\
&\leq \left| \int_a^{x_1} \int_c^{y_1} \frac{a-t}{b-a} \frac{c-s}{d-c} d_s d_t f(t, s) \right| + \left| \int_a^{x_1} \int_{y_1}^{y_2} \frac{a-t}{b-a} \left[\frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c} \right] d_s d_t f(t, s) \right| \\
&\quad + \left| \int_a^{x_1} \int_{y_2}^d \frac{a-t}{b-a} \frac{d-s}{d-c} d_s d_t f(t, s) \right| + \left| \int_{x_1}^{x_2} \int_c^{y_1} \left[\frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a} \right] \frac{c-s}{d-c} d_s d_t f(t, s) \right| \\
&\quad + \left| \int_{x_1}^{x_2} \int_{y_1}^{y_2} \left[\frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a} \right] \left[\frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c} \right] d_s d_t f(t, s) \right| \\
&\quad + \left| \int_{x_1}^{x_2} \int_{y_2}^d \left[\frac{t-x_1}{x_2-x_1} + \frac{a-t}{b-a} \right] \frac{d-s}{d-c} d_s d_t f(t, s) \right| + \left| \int_{x_2}^b \int_c^{y_1} \frac{b-t}{b-a} \frac{c-s}{d-c} d_s d_t f(t, s) \right|
\end{aligned}$$

$$\begin{aligned}
& + \left| \int_{x_2}^b \int_{y_1}^{y_2} \frac{b-t}{b-a} \left[\frac{s-y_1}{y_2-y_1} + \frac{c-s}{d-c} \right] d_s d_t f(t, s) \right| + \left| \int_{x_2}^b \int_{y_2}^d \frac{b-t}{b-a} \frac{d-s}{d-c} d_s d_t f(t, s) \right| \\
& \leq \frac{x_1-a}{b-a} \frac{y_1-c}{d-c} \bigvee(f) + \frac{x_1-a}{b-a} \max \left\{ \frac{y_1-c}{d-c}, \frac{d-y_2}{d-c} \right\} \bigvee_{Q_2}(f) \\
& \quad + \frac{x_1-a}{b-a} \frac{d-y_2}{d-c} \bigvee_{Q_3}(f) + \max \left\{ \frac{x_1-a}{b-a}, \frac{b-x_1}{b-a} \right\} \frac{y_1-c}{d-c} \bigvee_{Q_4}(f) \\
& \quad + \max \left\{ \frac{x_1-a}{b-a}, \frac{b-x_1}{b-a} \right\} \max \left\{ \frac{y_1-c}{d-c}, \frac{d-y_2}{d-c} \right\} \bigvee_{Q_5}(f) \\
& \quad + \max \left\{ \frac{x_1-a}{b-a}, \frac{b-x_1}{b-a} \right\} \frac{d-y_2}{d-c} \bigvee_{Q_6}(f) + \frac{b-x_1}{b-a} \frac{y_1-c}{d-c} \bigvee_{Q_7}(f) \\
& \quad + \frac{b-x_1}{b-a} \max \left\{ \frac{y_1-c}{d-c}, \frac{d-y_2}{d-c} \right\} \bigvee_{Q_8}(f) + \frac{b-x_1}{b-a} \frac{d-y_2}{d-c} \bigvee_{Q_9}(f) \\
& \leq \max \left\{ \frac{x_1-a}{b-a}, \frac{b-x_1}{b-a} \right\} \max \left\{ \frac{y_1-c}{d-c}, \frac{d-y_2}{d-c} \right\} \bigvee_a^b \bigvee_c^d(f) \\
& = \frac{[(b-a)-(x_2-x_1)][(d-c)-(y_2-y_1)]}{(b-a)(d-c)} \left[\frac{1}{2} + \left| \frac{\frac{a+b}{2} - \frac{x_2+x_1}{2}}{(b-a)-(x_2-x_1)} \right| \right] \\
& \quad \times \left[\frac{1}{2} + \left| \frac{\frac{d+c}{2} - \frac{y_2+y_1}{2}}{(d-c)-(y_2-y_1)} \right| \right] \bigvee_a^b \bigvee_c^d(f)
\end{aligned}$$

which completes the proof. \square

2.1. Applications for Probability Density Functions. Let X and Y be two continuous random variables with $f : [a, b] \times [c, d] \rightarrow \mathbb{R}^+$ is a joint probability density function and $F : [a, b] \times [c, d] \rightarrow \mathbb{R}^+$, $F(t, s) = \int_a^t \int_c^s f(t, s) ds dt$ is its cumulative distribution function.

Proposition 1. *Let f and F be as above. Then we have*

$$\begin{aligned}
& \left| F(t, s) - \frac{t-a}{b-a} \int_a^b \int_c^s f(\xi, \eta) d\eta d\xi - \frac{s-c}{d-c} \int_a^t \int_c^d f(\xi, \eta) d\eta d\xi + \frac{(t-a)(s-c)}{(b-a)(d-c)} \right| \\
& \leq \frac{(b-t)(t-a)(d-s)(s-c)}{(b-a)(d-c)} \bigvee_a^b \bigvee_c^d(f).
\end{aligned}$$

Proof. Taking $x_1 = a$, $x_2 = t$, $y_1 = c$, $y_2 = s$ and using the fact that $\int_a^b \int_c^d f(\xi, \eta) d\eta d\xi = 1$, we have the required result. \square

REFERENCES

- [1] M. W. Alomari, *A Generalization of Weighted Companion of Ostrowski Integral Inequality for Mappings of Bounded Variation*, RGMIA Research Report Collection, 14(2011), Article 87, 11 pp.
- [2] M. W. Alomari and M.A. Latif, *Weighted Companion for the Ostrowski and the Generalized Trapezoid Inequalities for Mappings of Bounded Variation*, RGMIA Research Report Collection, 14(2011), Article 92, 10 pp.
- [3] M. W. Alomari, *A Generalization of Dragomir's generalization of Ostrowski Integral Inequality and applications in numerical integration*, Ukrainian Mathematical Journal, 2012, 64(4).
- [4] M. W. Alomari, *A companion of Ostrowski's inequality for the Riemann-Stieltjes integral* $\int_a^b f(t)du(t)$, where f is of bounded variation and u is of $r-H-Hölder$ type and applications, Applied Mathematics and Computation 219 (2013) 4792–4799.
- [5] N. S. Barnett, S. S. Dragomir, I. Gomm, *A companion for the Ostrowski and the generalized trapezoid inequalities*, Mathematical and Computer Modelling 50 (2009) 179–187.
- [6] H. Budak and M. Z. Sarikaya, *On Ostrowski type inequalities for functions of two variables with bounded variation*, RGMIA Research Report Collection, 17(2014), Article 153, 18 pp.
- [7] H. Budak and M. Z. Sarikaya, *On generalization Ostrowski type inequalities for functions of two variables with bounded variation*, Palestine Journal of Mathematics, 5(1) (2016), 86–97.
- [8] H. Budak and M. Z. Sarikaya, *On generalization of Dragomir's inequalities*, RGMIA Research Report Collection, 17(2014), Article 155, 10 pp.
- [9] H. Budak and M. Z. Sarikaya, *On generalization trapezoid inequality for functions of two variables with bounded variation and applications*, Int. J. Nonlinear Anal. Appl., 7(1), 2016, 77–85.
- [10] H. Budak and M. Z. Sarikaya, *A companion of Ostrowski type inequalities for functions of two variables with bounded variation*, Journal of Advanced Mathematical Studies, 8(2), 2015, 170–184.
- [11] P. Cerone, S. S. Dragomir, and C. E. M. Pearce, *A generalized trapezoid inequality for functions of bounded variation*, Turk J Math, 24 (2000), 147–163.
- [12] J. A. Clarkson and C. R. Adams, *On definitions of bounded variation for functions of two variables*, Bull. Amer. Math. Soc. 35 (1933), 824–854.
- [13] S. S. Dragomir, *The Ostrowski integral inequality for mappings of bounded variation*, Bull. Austral. Math. Soc., 60(1) (1999), 495–508.
- [14] S. S. Dragomir, *On the midpoint quadrature formula for mappings with bounded variation and applications*, Kragujevac J. Math. 22(2000) 13–19.
- [15] S. S. Dragomir, *On the Ostrowski's integral inequality for mappings with bounded variation and applications*, Math. Inequal. Appl. 4 (2001), no. 1, 59–66.
- [16] S. S. Dragomir, *A companion of Ostrowski's inequality for functions of bounded variation and applications*, Int. J. Nonlinear Anal. Appl. 5 (2014) No. 1, 89–97.
- [17] S. S. Dragomir, *Refinements of the generalised trapezoid and Ostrowski inequalities for functions of bounded variation*, Arch. Math. (Basel) 91 (2008), no. 5, 450–460.
- [18] S.S . Dragomir and E. Momoniat, *A Three point quadrature rule for functions of bounded bariation and applications*, RGMIA Research Report Collection, 14(2011), Article 33, 16 pp.
- [19] S. S. Dragomir, *Some perturbed Ostrowski type inequalities for functions of bounded variation*, Preprint RGMIA Res. Rep. Coll. 16 (2013), Art. 93.
- [20] Y. Jawarneh and M. S. M Noorani, *Inequalities of Ostrowski and Simpson Type for Mappings of Two Variables with Bounded Variation and Applications*, TJMM, 3 (2011), No. 2, 81–94.
- [21] D-Y. Hwang and S. S. Dragomir, *Comparing two integral means for mapping of bounded variation and applications*, RGMIA Research Report Collection, 16(2013), Article 11, 8 pp
- [22] W. Liu and Y. Sun, *A Refinement of the Companion of Ostrowski inequality for functions of bounded variation and Applications*, arXiv:1207.3861v1, 2012.
- [23] A. M. Ostrowski, *Über die absolutabweichung einer differentierbaren funktion von ihrem integralmittelwert*, Comment. Math. Helv. 10(1938), 226–227.
- [24] K-L. Tseng, G-S. Yang, and S. S. Dragomir, *Generalizations of weighted trapezoidal inequality for mappings of bounded variation and their applications*, Mathematical and Computer Modelling 40 (2004) 77–84.

- [25] K-L. Tseng, *Improvements of some inequalities of Ostrowski type and their applications*, Taiwan. J. Math. 12 (9) (2008) 2427–2441.
- [26] K-L. Tseng, S-R. Hwang, and S. S. Dragomir, *Generalizations of weighted Ostrowski type inequalities for mappings of bounded variation and applications*, Computers and Mathematics with Applications, 55(2008), 1785–1793.
- [27] K-L. Tseng, S-R. Hwang, G-S. Yang, and Y-M. Chou, *Improvements of the Ostrowski integral inequality for mappings of bounded variation I*, Applied Mathematics and Computation 217 (2010) 2348–2355.
- [28] K-L. Tseng, S-R. Hwang, G-S. Yang, and Y-M. Chou, *Weighted Ostrowski integral inequality for mappings of bounded variation*, Taiwanese J. of Math., Vol. 15, No. 2, pp. 573–585, April 2011.
- [29] K-L. Tseng, *Improvements of the Ostrowski integral inequality for mappings of bounded variation II*, Applied Mathematics and Computation 218 (2012) 5841–5847.

HÜSEYİN BUDAK, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

E-mail address: hsyn.budak@gmail.com

MEHMET ZEKİ SARİKAYA, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

E-mail address: sarikayamz@gmail.com