
Contents :

Letter from the Editor	2
Evaluation of the analgesic effect of Intraperitoneal instillation of lidocaine with and without meperidine versus non-steroidal anti-inflammatory drugs during and after gynecological laparoscopic procedures <i>Ahmed M. Radwan, Ehab F. Girbash, Mohamed I Taama.</i>	3
Use of Hysteroscopy and Pregnancy Outcomes during Assisted Reproduction by ICSI, Add on cost or certified indication! <i>Sayed Abdel-Moneim Mahmoud, Ahmed Mohamed Salama, Adel Shafik Salah El Deen, Alla Masoud Abd-Elgeid</i>	10
Assessment of Awareness, Knowledge, and Attitude of Suez University medical students towards Human Papilloma Virus vaccine (HPV): A Cross-sectional Study <i>Ahmed Sewidan, Ahmed Mohamed ElSherbiny ,Khaled Affi, Shymaa S. Ali</i>	21
Gonadotropin releasing hormone agonist (GnRHa) versus human chorionic gonadotropin (HCG) for triggering of ovulation in ovarian stimulation cycles in polycystic ovarian syndrome (PCOS) women <i>Ammar Hesham Dorra, Hamed Youssef, Ismail, Mohamed Sayed Abdelhafez, Khaled Samir, Ahmed Mousa ElMenayer</i>	28
Prevalence of Molar Pregnancy in Histopathological Examination of Products of Conception Following First Trimester Miscarriages <i>Mohamed Abdelmawla Ezzat Eldesouki, Osama Mahmoud Warda, Hanan Nabil Abdelhafez, Mahmoud Mohamed Abdelrazek</i>	39
Urine albumin/creatinine ratio for the assessment of albuminuria in hypertensive disorders of pregnancy <i>Mohamed I. Taema, Amr Sobhy Sweed, Ahmed M. Radwan</i>	51

<p>Comparison between Clomiphene Citrate / Metformin Stair Step Protocol and Traditional Protocol in Treatment of Polycystic Ovary Syndrome <i>Waleed F. Ghareeb, Nourhan E. Ahmad, Elham H. Madny, Osama E. Ashour</i></p>	57
<p>Ultrasound assessment of the anovaginal distance as a predictor of obstetric perineal tears <i>Nora Y. Elbaik, Elham H. Madny, Ahmed A. Aboelroose, Omima T. Taha</i></p>	65
<p>Platelet Indices Signaling in Early and Late Onset Preeclampsia <i>Rana Tharwat Abdelghaffar Amer, Hend Abdelrahman Shalaby, Mohamed Ibrahim Eid, Maher Elesawi Elgaly</i></p>	72
<p>Comparative study between the use of long-acting insulin versus multiple dose regimen in control of Gestational Diabetes Mellitus: a randomized controlled trial <i>Maii Nawara, Mohammad Abd El Hameed, Sara Ahmad, Tamer Borg, Noha Abdel Satar Sakna</i></p>	81
<p>Progesterone-Primed Ovarian Stimulation might be a Safe and Effective Alternative to GnRH-antagonist Protocol for Controlled-Ovarian Stimulation of Infertile PCOS Women <i>Khaled Fathy Helal, Mohamed El Sayed Rafeek, Walid Mohamed Elnagar</i></p>	95
<p>Cons and Pros of Interventions for Management of Ovarian Endometrioma in Infertile Women with Good Ovarian Reserve <i>Youssef Abdel Zaher, Ali A Morsi, Wagdy Megahed Amer</i></p>	106

Letter from the Editor:

Dear colleagues,

Very interesting subjects are included in this edition. Routine office hysteroscopy was not an added cost before ICSI even in cases with normal trans vaginal ultrasonography. It can diagnose and treat uterine cavity lesions on the same setting. An appreciable level of awareness about the HPV infection and the vaccination. Females had better awareness regarding the infection as well as the vaccine. Medical education has a positive impact on raising awareness regarding the causes of cervical cancer, the availability of the vaccine, and its protective efficacy.

The GnRH agonist could be an effective and safe alternative to the traditional HCG in ovulation triggering after sequential minimal ovarian stimulation in PCOS patients without affecting ovulation and clinical pregnancy rates. Progesterone primed ovarian stimulation is safe and effective for infertile PCOS women. It achieved appropriate LH suppression with nearly no profound suppression or moderate-to-severe OHSS. The outcomes of this protocol were comparable to the GnRH-antagonist protocol; however, it is more cost-effective.

Histopathological examination seemed to be a promising tool in the context of assessment of products of conception in which the prevalence of molar pregnancy was 7%. In addition, incidence of molar pregnancy has no significant correlations with all clinical features as well as with blood groups. Random PCR may be employed in urgent situations as a quick, simple, and accurate diagnostic for the detection of substantial proteinuria in hypertensive diseases during pregnancy. A short anovaginal distance predicted the occurrence of perineal tears significantly.

There is no clinical difference between using a multiple-dose regimen and using long-acting insulin analogues to control gestational diabetes regarding maternal and foetal outcomes. However, a multiple-dose regimen needs a shorter time for blood glucose control than long-acting insulin. It is to be noted that long-acting insulin is more expensive.

Best regards.

Aboubakr Elnashar

MD

Chief Editor of EFSSJ

Prof. obs Gyn. Benha university, Egypt

elnashar53@hotmail.com