




EGYPTIAN ACADEMIC JOURNAL OF  
**BIOLOGICAL SCIENCES**  
HISTOLOGY & HISTOCHEMISTRY


D



ISSN  
2090-0775

[WWW.EAJBS.EG.NET](http://WWW.EAJBS.EG.NET)

Vol. 15 No. 1 (2023)



## A Comparative Histological Study of Testicles and Epididymis in Pets (Dog and Cat) and Wild Animals (Genet and Mongoose)

Derouiche Louiza<sup>1</sup>, Zatra Yamina<sup>2</sup>, Ouennes Houria<sup>3</sup>, Sahi Sameh<sup>3</sup>, Ghadbane Karima<sup>2</sup> and Bouadel Amira<sup>2</sup>

<sup>1</sup>Higher School of Food Sciences & Agri-food Industries Algiers, Algeria.

<sup>2</sup>Saad Dahleb Blida1 University, Faculty of Natural and Life Sciences, Algeria.

<sup>3</sup>Laboratory of Animal Productions, Biotechnologies and Health. Institute Agronomic and Veterinary Sciences, Souk-Ahras University, Algeria.

E.Mail\* : [derouiche\\_fatma@yahoo.fr](mailto:derouiche_fatma@yahoo.fr)

### ARTICLE INFO

Article History

Received:7/5/2023

Accepted:8/6/2023

Available:12/6/2023

### Keywords:

Pets-epididymal-canals, Histology, Testis, Wild-animals.

### ABSTRACT

This paper study aims to compare morphologically and histologically the testes of the two predatory animals (mongoose and genet) with the two pets (cat and dog). The research is based on direct observations, preparation of histological sections, and accurate measurement of the seminiferous tubule diameters and epididymal ducts. For this purpose, the testes and epididymis have been removed and quickly fixed. The results of the histological comparison showed that the size is more pronounced in dogs than in other species; significant differences were found between wild and domestic species. This study requires further investigation regarding the great differences between selective breeding and implies the possibilities of using biotechnologies, such as artificial insemination, to increase their multiplication especially that of those exposed to extinction.

### INTRODUCTION

Life development of men passed by several steps, such as domesticating wild species, which represents an important period of the development of human societies. Among the very first domesticated animals that went from wild animals to pets, the dog *Canis lupus familiaris*, although it belongs to the same wolf race Canidae family, was domesticated, which is the best witness of the great development of farming. Next, men were rather able to domesticate also the cat *Felis silvestris catus* (Ollivier, 2017).

Despite domesticating and adopting several animals for our own benefit, several others are still wild, including the genet *Genetta genetta* and the mongoose *Herpestes ichneumon*, belonging to the same breeds of dogs and cats, that the *Carnivora*. These two species have always kept their wild character, which translated into damage caused to farm animals, leading to death while they are protected (IUCN, 2010).

Animals' reproduction inside farms is better than in nature, because, it has been said that wild animals may reproduce much better inside farms where they are well-fed and well overseen by veterinary care throughout their lives (Khamas *et al.*, 2014). Moreover, being aware of the reproductive biology of these species is very important for the correct management of their breeds, especially in its basic aspects, so that many studies could be carried out performed using them as a biological model for the physiology of reproduction, especially species in danger of extinction (Mehanna *et al.*, 2018).

The increased interest in breeding dogs and cats and their use as models for other canids and felids demand research to improve reproductive techniques. Among them, testicular cryopreservation stands out. Testicular cryopreservation enables the maintenance of reproductive capacity and allows the establishment of germplasm banks for several species of commercial value or at risk of extinction. Furthermore, it enables the transport of genetic material among different regions. It is noteworthy that this biotechnology represents the only possibility of preserving the fertility of prepubertal animals that have died, so it has great importance in the propagation of the genetic material of animals (Silva, 2022).

We tried to find out if there are differences in histology, morphometry and similarities between several species. This study aims to provide more

information on the intraspecific variation of male reproductive organs for two wild animals (genet and mongoose), using domestic animals (cat and dog) as a model. Knowing about the diversity in various organs and physiological and behavioural measures of these species is a prerequisite for the control of reproduction as well as the use of biotechnologies for the preservation of these species, especially those that are in danger of extinction.

## MATERIALS AND METHODS

### The Animals:

Adult testicles of four different species were used for this experiment. This is a prospective study carried out within the Pathological Anatomy Department of the Issad Hassani university hospital of Beni Messous in Algiers, Algeria. The details are shown in Table 1.

**Table 1:** Animals studied in our experiment.

Animal	Species	Region
Cat	<i>Felis catus</i>	Veterinary clinic in Dely-Brahim, Algeria
Dog	<i>Canis lupus familiaris</i>	Veterinary clinic in Kouba, Algeria
Mongoose	<i>Herpestes ichneumon</i>	Recovered dead from the forest in Souk-Ahras, Algeria
Genet	<i>Genetta genetta</i>	Recovered dead in Belezma national park in Batna, Algeria

### The Histological Study:

In the present study, we have cleaned the male genitalia and removed the scrotal and the adipose tissue. We performed two types of sections, transverse and longitudinal (Derouiche *et al.*, 2023). Dealing with these samples was carried out through a series of mandatory successive steps, the purpose of which is to obtain fine cuts ready to receive the staining of interest. The procedure used is inspired by that developed by Martoja and Martoja (1967) and Vilar *et al.* (2017).

### Dehydration and Impregnation:

The testicles and the epididymis were placed in paraffin blocks, using old histological techniques (increasing dehydration in alcohol baths, and inclusion in paraffin). To achieve dehydration, we used a series of ethyl

alcohol baths of increasing degrees (50%, 70%, 80%, 90%, 100%), for 2 hours for each bath to avoid disorganization of the structures.

The last bath is a Xylene bath to complete the dehydration and prepare the impregnation of the organ with paraffin because Ethanol and paraffin are not miscible. Immediately after the Xylene baths, the organs were immersed in three successive paraffin baths, each for 2 hours at 60°C; it is impregnation. The second and third baths contain pure paraffin, while the first is half paraffin and half Xylene.

### Inclusion:

For the inclusion operation, the organs were placed in molds which will receive paraffin. The respective cassettes, identifying each sample, were placed on the surface of the moulds.

paraffin was poured inside the molds until the sample was completely submerged. Then, the device is placed on a cooling plate of the device (-10°C to -15°C) until the block becomes solid.

### Cutting, Staining Colouring, and Observation:

Sections of 3 µm were made in the microtome and then rehydrated and stained with:

-Hematoxylin-Eosin: This stain turns the cytoplasm pink with Eosin and the nucleus purple with Hematoxylin.

-Congo Red: This staining marks amyloid deposits in red.

-Masson's Trichrome: Used to differentiate collagen fibres and muscle tissue in histological sections before staining, the rehydration was performed in a reverse sequence to that of dehydration. Mounting is the operation that consists of preserving the staining with the help of Eukitt (Merck, Darmstadt, r.f.a) which allows the adhesion between the slide and the coverslip. After mounting, the slides are dried on absorbent paper and finally observed. To draw histograms and to compare the sizes of the seminiferous tubules and epididymides of our samples, measurements were made on histological sections of the animals studied. The images were captured by a digital camera (hero cam, ma88-500, BME lab and science, st. Paul, USA) connected to a photonic microscope (Optika b 235, Italy) via TS view software (microscopes America, cumming, ga, USA).

### Morphometric Study:

The surface area, diameter (minor and major axis), perimeter for seminiferous tubules and epididymides at G×10, and epididymides contours at G×40 were measured using image analysis and processing software "Axio Vision 4.6.3.0" developed by Carl Zeiss company.

In our morphometric study on the specimens of each species, and for each section, 50 measurements were taken for the surface, the diameter and the perimeter enlarged at G×10 magnification; for the seminiferous

tubules as well as 20 magnified measurements at G×40 magnification to measure the outline of the epididymides and the heights of the main cells of the epididymis.

### Statistical Analysis:

The results obtained are presented in the form of means ± Mean Squared Error (MSE).

-Arithmetic mean (x) of individual values:

$$x = \frac{\sum_{i=1}^n xi}{n}$$

Σx : Sum of individual values

n : number of values

-Standard error of the mean (S.E.M.):

$$S.A.M = \frac{\delta}{\sqrt{n}} \quad \delta = \sqrt{\frac{\sum(xi-x)}{n-1}}$$

δ : Standard deviation, xi: Individual value

-Correlation coefficient r:

$$r = \frac{p}{\delta x \delta y} p = \frac{1}{n} \sum xi yi - xy$$

$$(\delta x)^2 = \frac{1}{n} \sum (xi - x)^2 (\delta x)^2 = \frac{1}{n} \sum (yi - y)^2$$

xi and yi: individual values compared; x and y: average of individual values compared.

### -Statistical Validity:

The statistical validity of the differences is calculated by Student's test using the statistical software. The difference between the two compared means is statistically significant if the probability "p", read as a function of the number of degrees of freedom (d.d.l. = n<sub>1</sub> + n<sub>2</sub> - 2) is equal to or less than 5%.

## RESULTS

Our study is based on the comparison of the male genitalia of the cat "*Felis catus*" and the dog "*Canis lupus familiaris*" with two other species, the genet "*Genetta genetta*" and the mongoose "*Herpestes ichneumon*".

### Morphological Study:

Before making the histological sections, we made a morphological comparison of the testicles. We noticed that the size of the dog's testicles was higher than that of the three other species, namely the mongoose, the cat, and the genet, and the same is true for the morphology of the epididymis (Fig. 1).




**Fig. 1:** Testicles of the dog (1), the cat (2), the mongoose (3) and the genet (4).

### Histo-morphometry:

#### Structural Aspect and Morphometry of The Seminiferous Tubes:


At the magnification Gx4; In *Felis catus* and *Canis lupus familiaris*: Seminiferous tubules are fused with a central lumen. In *Herpestes ichneumon* and *Genetta genetta*: the seminiferous tubules are scattered with an invisible lumen. The extra-tubular space is occupied by interstitial tissue (Fig. 2).



**Fig. 2:** The structural aspect of the seminiferous tube in *Felis catus*, *Herpestes ichneumon*, *Canis lupus familiaris* and *Genetta genetta* at Gx4 magnification. Stained with Hematoxylin-Eosin (HE), Congo Red (CR), Masson's Trichrome (MT), Scal bar: 200µm. ST: Seminiferous Tube, L: Lumen, IT: Interstitial Tissue.

At Gx10 magnification; in *Felis catus* and *Canis lupus familiaris*: We observed a network of ducts that collect the products of the seminiferous epithelium. The seminiferous tubules consist of a central lumen sometimes

occupied by spermatozoa, it is bordered by a seminiferous epithelium, the seminiferous tubules are fused, and between the tubules, we found spaces occupied by interstitial tissue (Fig. 3).



**Fig. 3:** The structural aspect of the seminiferous tube in *Felis catus*, *Herpestes ichneumon*, *Canis lupus familiaris* and *Genetta genetta* at Gx10 magnification. Stained with Hematoxylin-Eosin (HE), Congo Red (CR), Masson's Trichrome (MT), Scal bar: 100µm. Ept: Epithelium, L: Lumen, IT: Interstitial Tissue, F: Flagella.

The morphometric study showed that the surface area ( $\mu\text{m}^2$ ) of these tubes in cats and dogs was  $57704.5 \pm 1464.2$ ;  $33482,8 \pm 818,5$  respectively and the lumen of these tubes is  $16251 \pm 871.6$ ;  $5513 \pm 247.3$  respectively (Fig. 4: A, B).


We noted that the seminiferous tubules are large in both cats and dogs and are fused in an anarchic manner. A visible lumen poor of spermatozoa, which is large in the cat. The statistical

study showed that there is a highly significant difference between these two species studied in favor of *Felis catus*, the difference is 72.34%;  $p=0.0000$  for the surface of the tubes and 66.07%;  $p=0.0000$  for the lumen of these tubes.

In *Herpestes ichneumon* and *Genetta genetta*: the surface area ( $\mu\text{m}^2$ ) of these tubes is  $21728.4 \pm 524.3$ ;  $20191.1 \pm 1335.4$  respectively, the lumen is  $364.7 \pm 13.7$ ;  $9366.3 \pm 1054.8$

respectively (Fig. 4: A, B). An invisible lumen full of spermatozoa, which is large in *Genetta genetta*. The presence of a

large envelope surrounding the seminiferous tubules was also noticed.




**Fig. 4:** Tissue morphometry of the seminiferous tubules, *Canis lupus familiaris*, *Felis catus*, *Genetta genetta*, *Herpestes ichneumon*, A: Surface of the seminiferous tubule; B: Surface of lumen.

The surface area ( $\mu\text{m}^2$ ) of the envelope is  $54530.9 \pm 1262.8$  in *Genetta genetta* and  $40548.4801 \pm 1007.11657$  in *Herpestes ichneumon*, there is no significant difference between the two species studied since the difference is only 7.6%;  $p=0.2831$ . Whereas, for the surface area of the lumen of these tubes, the difference is statistically highly significant, -96.1%;  $p=0.000000$  and -25.9031%,  $p=0.0000$ .

In *Felis catus* and *Canis lupus familiaris*, the seminiferous epithelium delimiting a central lumen shows that the epithelium consists of a thin peripheral

tunica fibrosa and several layers of cells. The most mature cells are located close to the lumen while the least mature cells are close to the extern tunica. They correspond to four generations of cells that have started their evolution. The size and shape of the cells vary from the periphery to the lumen according to the size and appearance of the nucleus. The tissue occupies a small area and contains large cells either isolated or in clusters, closely related to the blood and lymphatic capillary networks and linked to the seminiferous tube (Fig. 5).




**Fig. 5:** Structural aspect of the seminiferous tube in *Felis catus*, *Herpestes ichneumon*, *Canis lupus familiaris* and *Genetta genetta* at Gx40 magnification. Stained with Hematoxylin-Eosin (HE), Congo Red (CR), Masson's Trichrome (MT), Scal bar: 50µm. LC: Leydig Cell, SPG: Spermatogonia, SPCI: Spermatocyte I, SPD: Spermatid, SPZ: Spermatozoa.

In *Herpestes ichneumon* and *Genetta genetta*: spermatozoa occupy the lumen. The seminiferous epithelium contains cells at different stages of maturation distributed along the epithelium in an unorganized manner, the interstitial tissue occupies a large area and contains large cells either isolated or in clusters (Fig. 5).

At the magnification G×100 (Fig. 6), in *Felis catus* and *Canis lupus familiaris*: The seminiferous epithelium is made up of cells that have reached different stages of differentiation, in particular, the characteristics of the cells, from the periphery to the lumen are successively observed as:

- Spermatogonia: Small and oval, with a large nucleus.
- Spermatocyte: Slightly larger than the above and circular, with a large round nucleus containing chromatin arranged in coarse clusters or fine filaments.
- Spermatid: Similar in size to the above, with a relatively smaller and denser nucleus and a homogeneous, finely granular cytoplasm, and their position is close to the light.
- Spermatozoa: Elongated, teardrop-shaped cells with a flagellum on the lumen side of the tube and containing an equally long, slightly curved nucleus of uniform coloration.



**Fig. 6:** The structural aspect of the seminiferous tube in *Felis catus*, *Herpestes ichneumon*, *Canis lupus familiaris* and *Genetta genetta* at Gx100 magnification. Stained with Hematoxylin-Eosin (HE), Congo Red (CR), Masson's Trichrome (MT), Scal bar: 10 $\mu$ m. LC: Leydig Cell, SPG: Spermatogonia▲, SPCI: Spermatocyte I, SPD: Spermatid, SPZ: Spermatozoa, BL: Basal Lamina.


#### The Histologic Aspect of the Epididymis:

At the magnification Gx10, in *Felis catus* and *Canis lupus familiaris*: The epithelium of the epididymis is of

the pseudostratified cylindrical type and the cells are provided with long stereocilia. The epididymis is surrounded by a circular layer of smooth muscle cells. The lumen is sometimes occupied

by spermatozoa. In *Genetta genetta*: The epithelium is pseudostratified cylindrical with the presence of stereocilia at the

apical pole, the extra-tubular space is occupied by smooth muscle cells (Fig. 7).




**Fig. 7:** The structural aspect of the seminiferous tube in *Felis catus*, *Herpestes ichneumon*, *Canis lupus familiaris* and *Genetta genetta* at Gx10 magnification. Stained with Hematoxylin-Eosin (HE), Congo Red (CR), Masson's Trichrome (MT), Scal bar: 100µm. SPZ: Spermatozoa, Ed: Epididymal duct, L: Lumen.

The surface area (µm<sup>2</sup>) of the epididymis in cats and dogs is 7356.7 ± 545.8; 39628.6 ± 1995.2 respectively, and that of the lumen is 8138.56 ± 332.3; 14227.42 ± 973.2 respectively (Fig. 8: A, B) so there is a highly significant statistical difference between these two species in favor of *Felis catus* and *Canis lupus familiaris*, the difference is 438.7%; p=0.000000 for the surface area of the tubes and 74.8%; p=0.000000 for epididymal lumen in favor of the dog.

In *Herpestes ichneumon* and *Genetta genetta*: The surface area (µm<sup>2</sup>)

of the epididymis is high in the mongoose (116379.7 ± 2800.4), compared to the genet (34.7 ± 1995.2), the same is true for the lumen, which is visible and sometimes full of spermatozoa (Fig. 8: A, B). Thus, the statistical study shows a highly significant difference between these two species in favor of *Herpestes ichneumon*. The difference is 482.6%; p=0.0000 for the surface of the tubes and 1165.5%; p=0.0000 for the epididymis lumen.




**Fig. 8:** Epididymis surface (A) and epididymis lumen (B), of the 4 species studied, *Felis catus*, *Canis lupus familiaris*, *Genetta genetta*, *Herpestes ichneumon*.

In all four species studied, the epididymal canal wall consists of an epithelium of constant height, which is not scalloped and rests on a muscular connective tissue layer via a basement membrane. The circular lumen contains numerous spermatozoa. We also observed cells resting on the basal lamina in the deep part of the epithelium, and prismatic principal cells with stereocilia

at the apical pole, and a chorion containing circular smooth muscle fibers (Fig. 9).


We noticed that the principal cells had long stereocilia in *Canis lupus familiaris* with a full sperm lumen with a higher epithelial height and principal cells resting on the basal lamina up to the apical pole and basal cells resting on the basal lamina.



**Fig. 9:** The structural aspect of the epididymis in *Felis catus*, *Canis lupus familiaris* and *Genetta genetta* at Gx40 magnification. Stained with Hematoxylin-Eosin (HE), Congo Red (CR), Masson's Trichrome (MT), Scal bar: 50µm. SPZ: Spermatozoa, Pc: Principal cell, Mc: Muscle cells, L: Lumen, Mv: Microvilli, EPT: Epithelium, Bc: Basal cell, Ept: Epithelium.

In *Felis catus* and *Canis lupus familiaris*: The heights ( $\mu\text{m}$ ) of the epithelium and the supra-nuclear zone are respectively  $33.9 \pm 0.6$ ;  $40.28 \pm 1.8$  and  $19.3 \pm 0.5$ ;  $28.6 \pm 0.8$  in dogs and cats (Fig. 10: A, B), so it can be seen that

the dog cells are higher and the differences are 18.78%;  $p=0.002241$  for epithelial height and 48.12%;  $p=0.000000$  for the height of the supra-nucleus.



**Fig. 10:** The height of the epididymis epithelium (A) and the height of the epididymis nucleus (B), of the 3 species studied, *Felis catus*, *Canis lupus familiaris*, *Genetta genetta*.

## DISCUSSION

Testicular cryopreservation enables the maintenance of reproductive potential, the creation of germplasm banks and the transport of genetic material between different regions. This biotechnology represents the only possibility of preserving the fertility of prepubertal animals that have already died or that need to undergo gonadotoxic treatments. Despite advances in the use of cryopreserved testicular fragments, protocols that can be used in the clinical routine of dogs and cats have not yet been established. Due to the great importance of the topic, the objective of this review is to provide an overview of the subject, approaching the main works on testicular cryopreservation in dogs and cats (Silva, 2022).

The need for promotional projects for the preservation of endangered species in Algeria was the main source of motivation to carry out this study. To get to know these species well, we've chosen to lead a comparative

histological study between the testicles and epididymis of wild animals (genet and mongoose) and domestic those of domestic ones (dog and cat). This study showed comparable histological characteristics overall. Some differences were observed and described. From an anatomical point of view, the histology of the testis and epididymis, is comparable to that found in the four species, but concerning the size of the gonads of these different animals studied, we noticed that the dog's testicle comes first, then the mongoose, and lastly the genet after the cat. This means that there is a conformity between the weight of the testis and the body weight of the animal.

Our results differ from the results of França and Godinho (2003), who found that the average weight of the testes of the cat is 1.2 g and that there is no significant correlation between the weight of the testes and the body weight ( $r = 0.36$ ), on the other hand, they found out that there is a similarity between the two testes ( $p > 0.05$ ). In fact, there is a

positive correlation between body weight, testicular weight and plasma testosterone concentration (Berger *et al.*, 1982). Comparing the histological aspects of the four testes of these species studied, we found that the seminiferous tubules of *Félis silvestris catus* and *Canis lupus familiaris* are fused together with a larger size, colored green with Masson's Trichrome, this space is reduced. Whereas, in the mongoose and genet, the seminiferous tubules are scattered, dispersed and separated from each other by an interstitial tissue in the form of a network, the latter taking up a large space, which is full of Leydig cells the tubes are small in relation to the first ones. The seminiferous tubules are surrounded by envelopes located between the seminiferous tubule and the interstitial tissue. These testicular envelopes are larger in *Genetta genetta* by contribution to *Herpestes ichneumon*.

On the other hand, in *Genetta genetta* and *Herpestes ichneumon*, it is difficult to identify the type of germ cells, with a seminal tube lumen full of spermatozoa, which means that they are in the reproductive period. Garcia-Tomas *and al.* (2009a) and Garcia-Tomas *and al.* (2009b) reported that spermatogenesis varies according to species, environment and management, which should be taken into consideration in studies. The interstitial tissue of the testis located in the intertubular compartment is important for the nutrition of the cells of seminiferous tubules, transporting hormones and androgen production. The spaces between the seminiferous tubules of the testes are filled with connective tissue, blood and lymph vessels, and Leydig cells or interstitial cells, the main components of this compartment (Junqueira and Carneiro, 2013).

The distribution of germ cells in the seminiferous epithelium (spermatogonia, spermatocytes I, spermatocytes II, spermatids) is well organized in the seminiferous epithelium in *Felis catus* and even in *Canis lupus familiaris*.

Caldeira (2010) reports on the morpho-functional analysis of the testis and spermatogenesis process of the crab-eating fox (*Cerdocyon thous*, Linnaeus, 1766) that spermatogenesis is a synchronous and regular process of differentiation and cell division, whereby a spermatogonium gradually differentiates a highly specialized haploid cell, the sperm.

The testicular envelopes are larger in *Genetta genetta* compared to *Herpestes ichneumon*. The distribution of germinal cells in the seminiferous epithelium (spermatogonia, spermatocytes, spermatids) is well organized in *Felis silvestris catus* and *Canis lupus familiaris*. These results are similar to those of França and Godinho (2003) who observed well-organized germ cells at different stages of spermatogenesis in the cat's testis. Yasser *and al.* (2012) reported in chinchilla rabbits that the height of the epithelium of the seminiferous tubes decreases from the first to the 4th week and remains constant until the 7th week then increases again, about the onset of spermatogenesis.

On the other hand, in *Genetta genetta* and *Herpestes ichneumon*, it is difficult to identify the type of germ cells, the latter has an elongated nucleus, and the light becomes invisible because it is full of spermatozoa which means that they are in the breeding season. Our results are in agreement with that of according to Mehanna *and al.* (2016), in cats, interstitial cells and Leydig cells are more abundant compared with other species and substantially fill the space intertubular; they have a polyhedral shape with a large spherical nucleus and evident nucleolus; which resembles the Leydig cells present in the Pampas cat. While, The Sertoli cells are less frequent, distinguished by a more elongated shape and pyramidal cells with irregular contours and extending from the basement membrane to the tubule lumen and appear spherical to the oval nucleus, nucleolus with noticeably stained.

Silva *and al.* (2009) describe the Leydig cells in domestic cats as showing varied dimensions with polyhedral shape, vacuolated cytoplasm, clear nucleus, and nucleolus evident.

In canids, interstitial cells and Leydig cells are present in intertubular spaces, but to a lesser extent than observed in cats, along with the connective tissue; have a polyhedral shape with a large spherical nucleus and evident nucleolus (Diagone *et al.*, 2012). The latter is probably associated with cell death, which increased with age. According to histological analysis of the testes of hoary fox *Lycalopex vetulus* (Lund, 1842), the seminiferous tubules are formed by the columnar epithelium consisting of spermatogenesis cells and Sertoli cells, surrounded by a basement membrane, separated by interstitial tissue (Mehanna *et al.*, 2018).

However, the *Leopardus colocolo* male individuals have the testicle constituted by a capsule called the tunica albuginea, seminiferous tubules with stratified epithelium well developed with alternating presence and absence of light, evident Sertoli cells, and intertubular compartments or developed and interstitial vascular tissue, with Leydig cells (Mehanna *et al.*, 2016).

According to Bacha and Bacha (2003), combinations of spermatogenesis cells of the epithelium development occur inside a seminiferous tubule, in which these cellular stages occupy a portion of the tubule. The wall of the epididymal canal in all four species consists of the epithelium of constant height, not scalloped, resting via a basement membrane on a connective-muscular layer, with a circular lumen containing numerous spermatozoa. We also observed cells resting on the basal lamina in the deep part of the Epithelium, and main prismatic cells presenting at the apical pole of long stereocilia and a chorion containing circular smooth muscle fibers. Knowing that the epididymis in mammals is considered a very important segment in extra-testicular sperm via. In *Canis lupus*

*familiaris*, the height of the epithelium varies depending on the epididymal segment considered, the epithelium being higher in the region of the head and lower in that of the tail.

On the other hand, the size of the epididymis is narrow at the level of the head and wider at the level of the tail. An internal circular smooth muscle layer, increasing in thickness from the head to the tail, and an external longitudinal layer, visible from the body, surrounding the epithelium and the basal lamina.

The maturation and storage of sperm are its main functions. Further, the epididymis provides a "biochemical environment" in which, the sperm undergoes morphological and physiological changes, affecting its functional maturation passing through several regions of the epididymal (Schimming *et al.*, 2002). In this sense, these latter present morphological data on epididymal ducts of dogs near the hoary fox, where the epididymal duct is lined by pseudostratified columnar epithelium with a cell population consisting of principal cells, basal and apical, present in all regions.

Hoshino *and al.* (2002) showed, in a morphometric study of epididymal ducts of domestic cats, the same pattern of cell structures of the *Leopardus colocolo* epididymal epithelium. The epididymal duct showed pseudostratified columnar epithelium, standing on a delicate basement membrane and integrated by myoid cells. In the domestic cat (*Felis catus*) the terminal part of the epididymis continues in the vas deferens where the epithelial continues to be pseudo-stratified with small cell stereocilia (Diagone *et al.*, 2012). As well as the pampas cat (*Leopardus colocolo*), whose vas deferens showed a thick layer of smooth muscle, a rounded mucosa without longitudinal folds and pseudostratified columnar epithelium with the presence of stereocilia.

## CONCLUSION

This study attempts to provide more information on the morphological

aspects of the testes of the different selected species. It appears that the histological appearance and distribution of the seminiferous tubules of wild animals are different from those of domestic animals. Although the number of germ cells produced during spermatogenesis is affected by hormones, such as gonadotropins or androgens, these structural histological differences necessarily reflect functional differences in the male reproductive system which may also reflect differences in the rate and timing of reproduction, which are key factors in animal behavior.

It is essential to try to illustrate the contribution of the anatomy of the reproductive system in the knowledge of these species to be used. Similarly, it would be interesting to see the possibilities of using biotechnologies, especially artificial insemination to increase their multiplication. Our aim is that this research is the starting point for other more profound studies on a large population, taking into account the impact of the different influential factors.

#### **Acknowledgment**

The authors would like to acknowledge the staff of the Belezma national park in Batna and the veterinarians of the veterinary clinics in Dely- Brahim and Kouba in Algiers.

#### **Ethical Approval:**

The experiment was carried out according to the national regulations on animal welfare.

#### **REFERENCES**

- Bacha, W. J. and Bacha, L. M. (2003). Mâle reproductive system, In: Ibid. (Eds), *Atlas Colorido de Histologia Veterinária*. 2a ed, Roca, São Paulo, 335-336.
- Berger, M.; Jean-Faucher C. H.; De-Turckhein, M.; Veyssiere, G.; Blanc, M. R.; Poirier, J. C. and Jean, C. (1982). Testosterone, luteinizing hormone (LH) and follicle stimulating hormone (FSH) in plasma rabbit from birth to adulthood. Correlation with sexual and behavioural development. *Acta endocrinol (Copenh)*, 99 (3): 459-465.
- Caldeira, B. C.; Rêgo de Paula, T. A.; Pinto da Matta, S. L.; Balarini, M. K. and Araújo Campos, P. K. (2010). Morphometry of testis and seminiferous tubules of the adult crab-eating fox (*Cerdocyon thous*, Linnaeus, 1766). *Revista Ceres, Viçosa*, 57 (5).
- Derouiche, L.; Houssou, H.; Zatra, Z.; Boussaidane, B.; Sayahoui, H. and Ouennes H. (2023). Histomorphological comparisons in testicles between Barb horse, donkey and mule. *Acta Scientiarum. Animal Sciences*, v. 45, e58593.
- Diagone, K. V.; Feliciano, M. A.; Pacheco, M. R. and Vicente, W. R. R. (2012). Histology and morphometry of the testes of adult domestic cats (*Felis catus*). *Journal of Feline Medicine and Surgery*, 14(2) 124-130.
- França, L. R. and Godinho, C. L. (2003). Testis morphometry, seminiferous epithelium cycle length, and daily sperm production in domestic cats (*Felis catus*). *Biology of Reproduction*, 68: 1554-1561.
- Garcia-Tomas, M.; Sanchez, J. and Piles, M. (2009a). Post-natal sexual development of testis and epididymis in the rabbit: Variability and relationships among macroscopic and microscopic markers. *Animal Reproduction Science*, 110(3-4): 347-55.
- Garcia-Tomas, M.; Sanchez, J. and Piles, M. (2009b). Postnatal sexual development of testis and epididymis in the rabbit: Growth and maturity patterns of macroscopic and microscopic markers. *Theriogenology*, 71 (2) : 292-301.
- Hoshino, P.; Nakaghi, L. S. O.; Pacheco, M. R.; Lui, J. F. and Malheiros,

- E. B. (2002). Morphometry of the seminiferous tubules and epididymal ducts of dogs and cats from neutering surgery. *Biotemas*, 15 (1):97-110.
- International Union for Conservation of Nature (2010). Red list categories prepared by the IUCN species survival commission. IUCN, Gland, Switzerland and Cambridge, UK.
- Junqueira, L. C. and Carneiro, J. (2013). Basic Histology. 12<sup>th</sup> Ed Rio de Janeiro: Guanabara Koogan, 412-424.
- Khamas, W.; Al-Tikriti, M.; Albayati, M.; Tkalcic, S. and Eng, C. (2014). Histological Description of the Testis, Epididymis and Ductus Deferens of the Northern Great Grey Kangaroo (*Macropus giganteus giganteus*). *Journal of Cytology & Histology*, 5: 287.
- Martoja, P. M. and Martoja, R. (1967). Introduction to animal histology techniques. Eds Masson et cie, Paris, 345.
- Mehanna, M.; Ferreira, A. L. S.; Ferreira, A. and Paz, R. C. R. (2016). Morphology of the testes and epididymal ducts in the Pampas cat *Leopardus colocolo* (Molina, 1782). *Animal Morphophysiology. Pesquisa Veterinária Brasileira*, 36 no.10.
- Mehanna, M.; Ferreira, A. L. S.; Ferreira, A.; Paz, R. C. R. and Morgado, T. O. (2018). Histology of the testis and the epididymal ducts from hoary fox. *Lycalopex vetulus* (LUND, 1842). *Bioscience Journal Uberlândia*, 34, n 6, p. 1697-1705.
- Ollivier, M. (2017). Reconstructing and understanding the history of dog domestication through paleogenetics. *Archéozoologies*, 148: 50-55.
- Schimming, B. C.; Vicentini, C. A.; Tirapelli, L. F. and Tirapelli, D. P. C. (2002). Morphological examination of the epididymal duct in the dog. *Brazilian Journal of Veterinary Research and Animal Science*, 39 (2):61-65.
- Silva, C. A. O.; Perri, S. H. V.; Koivisto, M. B.; Silva, A. M.; Carvalho, R. G. and Monteiro, C. M. R. (2009). Aspectos histológicos e morfométricos dos testículos de gatos domésticos (*Felis catus*). *Pesquisa Veterinária Brasileira*, 29 (4): 321-316.
- Silva, L. D. M. d. (2022). Canine and Feline Testicular Preservation. *Animals (Basel)*, 12 (1): 124.
- Vilar, J. M.; Batista, M.; Carrillo, J. M.; Rubio, M.; Sopena, J. and Álamo, D. (2017). Histological, cytogenetic and endocrine evaluation in twenty-five unilateral cryptorchid horses. *Journal of Applied Animal Research*, 46(1), 441-444.
- Yasser, A. A.; Mahmoud, M. A. E. and Gamal, K. M. A. (2012). Histologica and histomorphometric changes of rabbit testis during postnatal development. *Research journal of Veterinary Sciences*, 5 (2): 42-50.