

## EFFECT OF ENERGY SOURCE ON NUTRIENTS UTILIZATION AND RUMEN MICROBIAL ACTIVITY OF OSSIMI SHEEP

B. M. Ahmed, S.S. Omar and M. S.A. Yassin

Department of Animal Production, Faculty of Agriculture, Menufiya University, Shebin El-Kom

(Received: Jan. 13, 2015)

**ABSTRACT:** This experiment was carried out in order to study the mode of action of the dietary energy source on (digestibility, N-balance and rumen activity). Three Ossimi rams aged 3 years were used in a 3 × 3 Latin Square design. The first treatment was the control ration (clover hay + concentrate feed mixture) without additive ( $R_0$ ); the second was the control ration plus 46g dry commercial fat (2.65% on DM basis,  $R_F$ ) while the third was the control ration plus 85g corn grain (4.88% on DM basis,  $R_C$ ).

Data reveal that dietary energy source significantly ( $P<0.01$ ) increased intakes of DM, OM, EE and NFE. Fat-supplementation showed the highest values EE intake while corn-supplementation showed the highest intakes for DM, OM, CP and NFE. Water intake followed the same pattern being more for energy-supplemented ration than the control group. Addition of fat increased ( $P<0.01$ ) the digestibility of DM, CP and EE. Corn-supplementation also increased the digestion coefficient of DM, OM, CP and NFE. Digestibility of CF was higher ( $P<0.01$ ) for the control group than the energy-supplemented groups. Dietary fat and corn grain supplementation improved nitrogen balance through the improvement of N digestibility. Values of ruminal pH before feeding were 7.43, 7.44 and 7.39 for the treatment groups  $R_0$ ,  $R_F$  and  $R_C$ , respectively. Differences were not significant. At 2-hr post feeding, pH declined with all treatment groups to reach the lowest values (being 6.55, 6.67 and 6.57 for the same respective groups). Significant ( $p<0.01$ ) differences among groups were found at all times post feeding being lower for the control and  $R_C$  ration than  $R_F$ . Rumen total VFA did not differ among the dietary treatments at 2hrs (post-feeding); however, it was significantly ( $P<0.01$ ) differed at 4 and 6h post feeding. Total VFA were 8.85, 8.44 and 9.03meq/dl of rumen liquor before feeding for  $R_0$ ,  $R_F$  and  $R_C$ , respectively. In general, VFA increased in the treated-groups to reach its peak at 4-hrs post feeding and declined thereafter. The highest values were reported for the corn-treated group being 9.03, 17.23 and 12.60meq/dl; at 0, 4 and 6h post-feeding, respectively; the respective values were 8.44, 15.39 and 9.73meq/dl for  $R_F$ . Before feeding,  $NH_3$ -N was 14.27, 13.06 and 12.45 mg/dl rumen liquor, differences were significant. Ammonia-N increased after feeding to reach the highest values for all dietary treatments at 2-hrs being 22.79, 21.15 and 20.60 mg/dl rumen liquor; and it decreased thereafter. Concentration of  $NH_3$ -N was significantly higher ( $P<0.01$ ) for the control group than both treated groups.

**Key words:** Dietary energy source, digestibility, microbial activity, Ossimi sheep.

### INTRODUCTION

The addition of fat to the diets of ruminant animals can help in covering the requirements of energy for high productive performance without causing metabolic disorders that often associated with large intakes of grain (Simas *et al.*, 1998). To increase energy density without the rumen acidosis and depression in milk fat due to the use of high starch and low fiber diets, attention has been directed to the inclusion of fats and feed stuffs with high concentration of fat and oils in ruminant

rations (Choi and Palmquist, 1999; Chan *et al.*, 1997; Simas *et al.*, 1997 and 1998; Zervas *et al.*, 1998; Casals *et al.*, 1999; Kowalski *et al.*, 1999; Offer *et al.*, 1999).

The published reports contain lots of conflicts about the effect of dietary fat on animal performance, digestion kinetics, rumen fermentation, blood constituents and carcass traits. Some reports indicate a negative effect on feed intake (Bendary *et al.*, 1994; Lough *et al.*, 1994; Talha, 1996). They reported that dry matter intake was

lower with animals fed fat-supplemented than those on un-supplemented rations. Others (Zinn, 1988 and 1989; El-Bedawy, 1989) found that fat supplementation did not affect feed intake. Digestibility of DM, OM and ADF was reported to decrease with adding fat (Hill and West, 1991); however, others (Bayourthe *et al.*, 1993; El-Bedawy *et al.*, 1994; El-Bedawy *et al.*, 1996; Talha, 1996) found an increase in digestibility of almost all nutrients. Bendary *et al.* (1994) found that digestibility of OM and NFE were not affected but CP and EE were significantly increased in supplemented-fat ration compared with the control. White *et al.* (1992) reported that supplemental fat did not affect average daily gain and feed efficiency. Although, Krehbiel *et al.* (1995) found that average daily gain linearly decreased ( $P < 0.01$ ) as dietary fat increased. Zinn and Plascencia (1996) found that addition of fat to the high forage diet increased ADG.

The present research was conducted to study the effect of supplementing rations with different dietary energy sources i.e., corn and dry fat on nutrient digestibility, N-balance and rumen fermentation of Ossimi sheep.

## **MATERIALS AND METHODS**

This experiment was carried out in the Farm of Animal Production Department of the Faculty of Agriculture, Menofiya University (Shebin El-Kom), in order to study the mode of action of the dietary energy source on (digestibility, N-balance and rumen activity). Three Ossimi rams aged 3 years with an average body weight of  $49 \pm 2$  kg were used in this study. Animals were randomly allocated to three treatments in  $3 \times 3$  Latin Square design. The first treatment was the control ration (clover hay + concentrate feed mixture) without additive ( $R_0$ ); the second was the control ration plus 46g dry commercial fat (2.65% on DM basis,  $R_F$ ) while the third was the control ration plus 85g corn grain (4.88% on DM basis,  $R_C$ ). Dietary energy of the second and third CFM was increased by 0.31 MCal/d (according to NRC, 2001). Lambs were fed at DM level of 3.5% body weight. The ratio of concentrate to roughage was 50: 50. The experimental concentrate mixtures are

presented in Table (1). Rams were housed individually in metabolic crates (160m x 0.53m) as described by Maynard *et al.* (1979) for separate collection of feces and urine. Rations and fresh water were presented once every day at 8.30h. Residuals were weighed daily every first week of the collection periods and subtracted from the offered amount to obtain the actual feed intake.

During the collection periods, feces were quantitatively collected at 8:00 a.m. before feeding; 10% from feces was withdrawn and dried to a constant weight in a forced air oven at 70°C for 24h. Dry fecal samples were kept ground in plastic pocket for later analysis. Urine was collected daily and a 10% aliquot was composited and refrigerated till analysis for nitrogen.

Rumen fluid samples were taken during the last two days of the collection periods. The samples were collected using the stomach tube attached to a vacuum pump, before feeding and then at 2, 4 and 6h after feeding. Rumen pH was measured immediately after collection using a digital pH meter (Sophisticated microprocessor, pH meter). Rumen fluid was strained through four layer of cheesecloth into plastic containers and kept at -20°C for later analysis. Half of the samples were acidified using concentrated ortho-phosphoric acid and 0.1N hydrochloric acid to determine the total volatile fatty acids (VFA). The second half of samples was alkaline using 0.1N NaOH to determine the concentration of rumen ammonia nitrogen.

The chemical analysis was carried out at the Laboratory of Nutrition, Department of Animal Production, Faculty of Agricultural, Minufiya University. The determination of dry matter (DM), organic matter (OM), crude protein (CP), crude fiber (CF), ether extract (EE), nitrogen-free extract (NFE) and ash in the feed, and feces were carried out according to AOAC (1990). Ammonia-nitrogen determination was carried out as soon as possible using the steam distillation method described by Ahmed (1976). Total volatile fatty acids (VFA) were measured according to AOAC (1990).

**Table 1: Composition of the formulated experimental concentrate mixtures .**

Ingredient	The experimental concentrate mixtures		
	Control ration R <sub>0</sub>	Fat- supplemented R <sub>F</sub>	Corn- supplemented R <sub>C</sub>
	%		
Yellow corn	44.6	42.6	48.9
Soybean meal	11.9	11.4	11
Cottonseed meal	19.8	18.9	18.1
wheat bran	19.8	18.9	18.1
Protected fat	-	4.3	-
NaCl	1	1	1
Limestone	2	2	2
Mineral and Vitamin premix	0.3	0.3	0.3
Sodium bicarbonate	0.3	0.3	0.3
Di-calcium phosphate	0.3	0.3	0.3
Total	100	100	100

Data of the present study were analyzed using GLM procedure of the SPSS program, version 13 (SPSS, 1997). Data were analyzed using the following linear model:  $Y_{ij} = \mu + A_i + e_{ij}$

where:  $Y_{ij}$  = the observation on the  $ij^{\text{th}}$  animal;  $\mu$  = overall mean;  $A_i$  = the effect of the  $i^{\text{th}}$  dietary treatments ( $i = 1, 2, 3$ );  $e_{ij}$  = a random error assumed to be independently randomly distributed with zero mean and variance  $\sigma^2_e$ , i.e. NID (0,  $\sigma^2_e$ ). All data measured as percentages were subjected to arc-sin transformation to approximate normal distribution before being analyzed.

**RESULTS AND DISCUSSION**

**1. Nutrient intakes of Ossimi rams as affected by dietary energy source:**

Data in Table (2) reveal that dietary energy source significantly ( $P < 0.01$ ) increased intakes of DM, OM, EE and NFE. Fat-supplementation showed the highest values for EE intake while corn-

supplementation showed the highest intakes for DM, OM, CP and NFE. Water intake followed the same pattern being more ( $P < 0.01$ ) for energy-supplemented ration than the control group.

Canale *et al.* (1990) reported that energy supplementation increased intake of DM and net energy. However, different kinds of fat-supplementation did not affect feed intake (West and Hill, 1990; Jenkins and Jenny, 1992; Schuff and Clark, 1992; Wu *et al.*, 1993; Salfer *et al.*, 1995; Cervantes *et al.*, 1996; Madison-Anderson *et al.*, 1997; Weiss and Wyatt, 2003). Others (Simas *et al.*, 1997; Patton, 2004) reported that DM intake tended to decrease when rations were supplemented with different fat sources. The conflict in the response of experimental animals in the above mentioned studies may have been due to the effect of fat sources used on the appetite. Mullins *et al.* (2010) found that DM Intake was increased by Holstein cows fed diets containing different levels of wet corn gluten feed.

Table 2: Nutrient intakes of Ossimi rams fed different energy sources.

Treatment*	Intake of						
	DM kg/d	OM kg/d	CP g/d	CF g/d	EE g/d	NFE g/d	Water l/d
R <sub>0</sub>	1.52 <sup>c</sup> ± 0.37	1.31 <sup>c</sup> ± 0.32	217.79 <sup>b</sup> ± 0.51	406.15 ± 1.27	49.36 <sup>c</sup> ± 0.12	641.14 <sup>c</sup> ± 1.33	5.23 <sup>b</sup> ± 0.11
R <sub>F</sub>	1.55 <sup>b</sup> ± 0.26	1.34 <sup>b</sup> ± 0.22	218.87 <sup>b</sup> ± 0.35	407.22 ± 0.87	79.53 <sup>a</sup> ± 0.08	636.04 <sup>b</sup> ± 0.91	6.19 <sup>a</sup> ± 0.10
R <sub>C</sub>	1.59 <sup>a</sup> ± 0.34	1.37 <sup>a</sup> ± 0.29	220.58 <sup>a</sup> ± 0.46	405.84 ± 1.13	52.22 <sup>b</sup> ± 0.10	691.72 <sup>a</sup> ± 1.19	6.11 <sup>a</sup> ± 0.11
Sig.	0.01	0.01	0.01	NS	0.01	0.01	0.01

(\*)R<sub>0</sub>, R<sub>F</sub> and R<sub>C</sub>, are control ration without or supplemented with 47g/d fat or 85g/d corn grain, respectively.

<sup>a,b</sup>Means with different superscripts within each column for each parameter are different (P<0.01). NS, not significant

## 2. Nutrient digestibility by Ossimi rams as affected by dietary energy source:

Table (3) presents the nutrient digestibility by Ossimi rams as affected by dietary energy source. Addition of fat increased (P<0.01) the digestibility of DM, CP and EE. Corn-supplementation also increased the digestion coefficient of DM, OM, CP and NFE. Digestibility of CF was higher (P<0.01) for the control group than the energy-supplemented groups.

Holter *et al.* (1992) reported that digestibility of ether extract was higher with fat supplemented diets. However, digestibility of CP was not affected by fat supplementation. Andrae *et al.* (2000) used sixty crossbred beef steers and found that digestibility of DM, OM, starch, and GE was greater (P<0.05) for the high-oil diet than the control diet, but lipid digestibility did not differ among treatments. Baraghit *et al.* (2003) reported that digestibility of DM, OM and CP was higher for fat-supplemented-followed by oil-supplemented-ration than the control one. El-Bedawy *et al.* (2004) found that 4% protected fat did not affect DM digestibility but increasing fat level to 8% decreased (P<0.05) DM digestibility by about 4 units. Digestibility of OM and CP was improved by incorporation of 4% dietary fat.

## 3. Nitrogen balance by Ossimi rams as affected by dietary energy source:

Results of nitrogen balance (Table 4) revealed that sheep fed all the experimental rations had almost similar nitrogen intake (NI) being 34.85, 35.02 and 35.29g/d. This could be attributed to that all the experimental ration were iso-nitrogenous.

The differences of fecal nitrogen (FN) between the control and the other groups were significant (P<0.01). Sheep on R<sub>F</sub> excreted significantly (P<0.01) less N in the feces (10.03g) than the other groups (11.24 and 10.89g for R<sub>0</sub> and R<sub>C</sub>, respectively). This resulted in better crude protein digestibility. Animals released almost equal amounts of N in the urine being 18.84, 18.25 and 18,21g for R<sub>0</sub>, R<sub>F</sub> and R<sub>C</sub> groups, respectively. The NB values were significantly (P<0.01) more in R<sub>F</sub> (6.74g) and R<sub>C</sub> (6.19g) than the control group (4.76g).

It is obvious that dietary fat and corn grain supplementation improved nitrogen balance through the improvement of N digestibility. El-Bedawy *et al.* (2004) reported that feeding 4% fat containing rations improved (P<0.05) nitrogen retention by 15% in comparison with control or 8% fat rations.

**Effect of energy source on nutrients utilization and rumen microbial.....**

**Table 3: Nutrient digestibility (%) by Ossimi rams fed different energy sources.**

Treatment*	DM	OM	CP	EE	NFE	CF
<b>R<sub>0</sub></b>	67.83 <sup>b</sup> ± 0.24	67.49 <sup>b</sup> ± 0.21	67.82 <sup>c</sup> ± 0.21	69.46 <sup>b</sup> ± 0.30	72.97 <sup>b</sup> ± 0.23	58.50 <sup>a</sup> ± 0.30
<b>R<sub>F</sub></b>	69.06 <sup>a</sup> ± 0.38	68.06 <sup>a<sup>b</sup></sup> ± 0.35	71.40 <sup>a</sup> ± 0.46	81.85 <sup>a</sup> ± 0.30	72.51 <sup>b</sup> ± 0.32	56.60 <sup>b</sup> ± 0.47
<b>R<sub>C</sub></b>	68.96 <sup>a</sup> ± 0.20	68.67 <sup>a</sup> ± 0.26	69.17 <sup>b</sup> ± 0.34	69.88 <sup>b</sup> ± 0.47	74.83 <sup>a</sup> ± 0.20	57.87 <sup>a</sup> ± 0.36
<b>Sig.</b>	0.01	0.05	0.01	0.01	0.01	0.01

(<sup>1</sup>)R<sub>0</sub>, R<sub>F</sub> and R<sub>C</sub>, are control ration without or supplemented with 47g/d fat or 85g/d corn grain, respectively. <sup>a,b</sup>Means with different superscripts within each column for each parameter are different (P<0.01). NS, not significant.

**Table 4: Nitrogen balance (g/d) by rams fed different energy sources.**

Treatment*	NI	FN	UN	NB
<b>R<sub>0</sub></b>	34.85 <sup>b</sup> ± 0.08	11.24 <sup>a</sup> ± 0.08	18.84 <sup>a</sup> ± 0.13	4.76 <sup>b</sup> ± 0.13
<b>R<sub>F</sub></b>	35.02 <sup>b</sup> ± 0.06	10.03 <sup>b</sup> ± 0.16	18.25 <sup>b</sup> ± 0.16	6.74 <sup>a</sup> ± 0.25
<b>R<sub>C</sub></b>	35.29 <sup>a</sup> ± 0.07	10.89 <sup>a</sup> ± 0.11	18.21 <sup>b</sup> ± 0.17	6.19 <sup>a</sup> ± 0.21
<b>Sig.</b>	0.01	0.01	0.01	0.01

(<sup>1</sup>)R<sub>0</sub>, R<sub>F</sub> and R<sub>C</sub>, are control ration without or supplemented with 47g/d fat or 85g/d corn grain, respectively. <sup>a,b</sup>Means with different superscripts within each column for each parameter are different (P<0.01). NS, not significant.


**4. Rumen activity by Ossimi rams as affected by dietary energy source:**

Data of sheep rumen fermentation (pH, VFA and NH<sub>3</sub>-N) as affected by dietary energy source are illustrated in Figures (1-3).

Values of pH before feeding were 7.43, 7.44 and 7.39 for the treatment groups R<sub>0</sub>, R<sub>F</sub> and R<sub>C</sub>, respectively. Differences were not significant. At 2-hr post feeding, pH declined with all treatment groups to reach the lowest values (being 6.55, 6.67 and 6.57 for the same respective groups). Significant (p<0.01) differences between groups were found at all times post feeding being lower

for the control and R<sub>C</sub> ration than R<sub>F</sub>. The lower pH was due to the higher fermentation ability of corn grain. Logically, pH values took the opposite trend of the total VFA.

Steele *et al.* (2012) reported that rumen pH for cattle fed high grain diet displayed lower rumen pH compared with cattle receiving the control diet. Zened *et al.* (2013) showed that rumen pH 5-h post feeding decreased from 6.4 to 5.7 when starch was added to the diet. Many studies (Ohajuruka *et al.*, 1991; Knapp and Grummer, 1991; Schauff and Clark, (1992) Madison-Anderson *et al.*, 1997; Abdelqader and Oba, 2012) reported that fat supplementation did not affect rumen pH.



**Fig. 1: Rumen pH rams fed different energy sources.**

Rumen total VFA (Fig 2) was not different among the dietary treatments at 2hrs (post-feeding); however, it was significantly ( $P < 0.05$ ) different at 0 time (before feeding) and ( $P < 0.01$ ) at 4 and 6hrs post feeding. In this respect Values were higher for the control group than the treated ones. Total VFA was 8.85, 8.44 and 9.03meq/dl of rumen liquor before feeding for  $R_0$ ,  $R_F$  and  $R_C$ , respectively. In general, VFA increased in the treated-groups to reach its peak at 4-hrs post feeding and decline thereafter. The higher values were reported for control and the corn-treated group. The values of corn-treated group being 9.03, 17.23 and 12.60meq/dl; at 0, 4 and 6h post-feeding, respectively; the respective values were 8.44, 15.39 and 9.73meq/dl for  $R_F$ .

The lowest VFA values reported for the dry fat-treated ration may have been due to a lower microbial activity in the rumen of sheep fed this ration than those received the control or corn-supplemented rations, perhaps because it contained less fermentable carbohydrate; it is worthy to mention that the control ration also contained more corn than the fat-supplemented ration (Table 1).


The published articles have some conflicts regarding the effect of dietary fat on rumen VFA concentration. Differences may have been due to different factors such as the form and/or level of fat used, the level of the dietary energy, the other ingredients in the diet, feeding frequency etc. Schauff and Clark (1992) reported that total VFA's concentrations were decreased when Ca-soaps was added to the diet. Madison-Anderson *et al.* (1997) reported that rumen total VFA's concentrations were greater for cows fed the control diet than those fed fat supplemented diet. Zened *et al.* (2013) reported that total VFA were significantly higher in the rumen of cows receiving high starch (HS) and high starch + sunflower oil (HS+OL) diets than in those of control CON and OL diet fed cows being 133 vs. 98 mM (on average), respectively. Sun and Oba (2014) reported that cows fed the dried distillers grains with soluble (DDGS) diet tended to have higher total VFA concentration in rumen fluid compared with cows fed the CON diet (107 vs. 116 mM;  $P = 0.06$ ).

Data of ammonia nitrogen concentration in the rumen of sheep as affected by the


**Effect of energy source on nutrients utilization and rumen microbial.....**

dietary energy source are presented in Fig (3). Before feeding, NH<sub>3</sub>-N was 14.27, 13.06 and 12.45 mg/dl rumen liquor, differences were significant. Ammonia-N increased after feeding to reach the highest values for all dietary treatments at 2-hrs being 22.79, 21.15 and 20.60 mg/dl rumen liquor; ammonia-N decreased thereafter.

Concentration of NH<sub>3</sub>-N was significantly higher (P<0.01) for the control group than the treated groups. Demeterova *et al.* (2002) investigated the influence of Ca-soaps on rumen fermentation. They reported that NH<sub>3</sub>-N decreased for cows fed diets contained Megalac.



**Fig. 2: Rumen totl VFA concentration (meq/dl) by rams fed different energy sources.**



**Fig. 3: Rumen ammonia-N concentration (mg/dl) by rams fed different energy sources.**

From the present study it could be recommended that sheep diets should be supplemented with energy source, either dry fat or corn in order to improve intake, digestibility, N balance as well as rumen fermentation.

## REFERENCES

- Abdelqader, M. M. and M. Oba (2012). Lactation performance of dairy cows fed increasing concentrations of wheat dried distillers grains with solubles. *J. Dairy Sci.* 95:3894–3904.
- Ahmed, B. M. (1976). The use of non-protein nitrogenous compounds in rabbit rations. M. Sc. Thesis. Tanta Univ.
- Andrae, J. G., C. W. Hunt, S. K. Duckett, L. R. Kennington, P. Feng, F. N. Owens and S. Soderlund (2000). Effect of high-oil corn on growth performance, diet digestibility, and energy content of finishing diets fed to beef cattle. *J Anim Sci.*, 78:2257.
- AOAC. (1990). *Association of Official Agricultural Chemists*. Official Method of Analysis. Washington D.C.
- Baraghit, G. A., El- Kholy, M. Nazley, S. S. Omar, B. M. Ahmed and Kh. Zedan (2003). I. Effect of dietary fat sources on digestibility, rumen fermentation and blood parameters of buffalo calves. *9<sup>th</sup> Conf. of Nutr. Hurghada, Egypt, 14-17 Oct. (2003)*.
- Bayourthe, C., R. Moncoulon and M. Veinay (1993). Effect of protein-Protected fat on ruminal and total nutrient digestibility of sheep diets. *J. Anim. Sci.*, 71:1026-1031.
- Bendary, M. M., I. A. Abou-Selim, M. R. M. Moustafa, A. M. Mahmoud and A. E. M. Khinzy (1994). Performance of fattening buffalo calves fed different levels of palm oil for two different periods. *Egyptian J. Anim. Prod.* (31). Suppl. Issue, Nov. 613-626.
- Canale, C. J., L. D. Muller, H. A. Mccahon, T. J. Whitsel, G. A. Varga and M. J. Lormore (1990). Dietary fat and ruminally protected amino acids for high producing dairy cows. *J. Dairy Sci.* 73: 135.
- Casals, R., G. Caja, X. Such, C. Torre and S. Calsamiglia (1999). Effects of calcium soaps and rumen undegradable protein on the milk production and composition of dairy ewes. *J. Dairy Res.* 66: 177.
- Cervantes, A., T. R. Smith and J. W. Young (1996). Effects of nicotin amide on milk composition and production in dairy cows fed supplemental fat. *J. Dairy Sci.*, 79:105.
- Chan, S. C., J. T. Huper, K. H. Chen, J. M. Simas and Z. Wu (1997). Effect of ruminally inert fat and evaporative cooling on dairy cows in hot environmental temperatures. *J. Dairy Sci.*, 80:1172.
- Choi, B. R. and D. L. Palmquist and Y. S. Son (1999). Effect of dietary fat on ruminal propionate and plasma insulin concentrations in lactating cows. *Korean J. Dairy Sci.*, 21:31.
- Demeterova, M., V. Vajida, P. Pastierik and A. Koteles (2002). The effect of protected fat and protein supplements on rumen metabolism, on some parameters of intermediary metabolism, and on the quality and production of milk in dairy cows. *Folia. Veterinaria*, 46:1.
- El-Bedawy, T. M. (1989). Fat in small ruminant nutrition, preliminary study.1- Effect of fat inclusion on intake digestibility and growth performance of goats and sheep fed high concentrate diets. *3<sup>rd</sup> Egyptian – British conference on animal, fish and poultry production, Alexandria, Egypt, 7-10 October*.
- El-Bedawy, T. M., H. M. El-Husseini, S. M. Allam and F. H. Shahin (1994). Effect of dietary fat and calcium supplements on in Vivo digestibility, rumen fermentation and some blood constituents of sheep. *Egyptian J. Anim. Prod.* Vol. 31 suppl. Issue, Nov. 59-73.
- El-Bedawy, T. M., M. A. I. Salem and E. A. Badr. (1996). Effect of dietary fat on growth performance and carcass characteristics of finishing bulls. *Egyptian J. Anim. Prod.* 33. Suppl. Issue, Nov. 103-111.

## **Effect of energy source on nutrients utilization and rumen microbial.....**

- El-Bedawy, T. M., Sawsan. M. Ahmed, M. A. I. Salem and H. A. A. Omer (2004). Effect of dietary protected fat and roughage level on digestion, rumen metabolism and plasma lipids of growing finishing lambs. Egyptian. J. Anim. Prod. 41. Supple Issue: 219- 236.
- Hill, G. M. and J. W. West. (1991). Rumen protected fat in Kline barley or corn diets for beef cattle: Digestibility, physiological, and feedlot responses. J. Anim. Sci. 69: 3376-3388.
- Holter, J. E., H. H. Hayes, W. E. Urban, Jr. and A. H. Duthie (1992). Energy balance and lactation response in Holstein cows supplemented with cottonseed with or without calcium soap. J. Dairy Sci., 75: 1480-1494.
- Jenkins, T. C. and B. F. Jenny (1992). Nutrient digestion and lactation performance of dairy cows fed combinations of prilled fat and canola oil. J. Dairy Sci., 75: 796-803.
- Knapp, D. M. and R. R. Grummer (1991). Response of lactating dairy cows to fat supplementation during heat stress. J. Dairy Sci., 74: 2573-2579.
- Kowalski, Z.M., P.M. Pisuleski and M. Sponghero (1999). Effect of calcium soaps of rapeseed fatty acids and protected methionine on milk yield and composition in dairy cows. J. Dairy Res., 66:475.
- Krehbiel, C. R., R. A. McCoy, R. A. Stock, T. J. Klopfenstein, D. H. Shain and R. P. Huffman (1995). Influence of grain type, tallow level and tallow feeding system on feedlot cattle performance. J. Anim. Sci. 73: 2916-2921.
- Lough, D. S., M. B. Solomon, T. S. Rumsey, S. Kalf and L. L. Slyter (1994). The effects of high-Forage diets with added palm oil on performance, plasma lipids, and carcass characteristics of ram lambs with initially high or low plasma cholesterol. J. Anim. Sci. 72: 330-336.
- Madison-Anderson, R. J., D. J. Schingoethe, M. J. Brouk, R. J. Bear and M. R. Lentsch. (1997). Response of lactating cows to supplemental unsaturated fat and niacin. J Dairy Sci., 80:1329.
- Maynard, L. A, J. K. Loosl, H. S. Hintz and R. G. Warner (1979). *Animal nutrition*. McGraw-Hill Book Co. Inc. NY.
- Mullins, C. R., K. N. Grigsby, D. E. Anderson, E. C. Titgemeyer and B. J. Bradford (2010). Effects of feeding increasing levels of wet corn gluten feed on production and ruminal fermentation in lactating dairy cows. J. Dairy Sci. 93:5329.
- NRC NRC, (2001). National Research Council. Nutrient requirements of domestic animals. Nutrient requirements of sheep. National Academy of science. Washington DC.
- Offer, N.W., M. Marsden, J. Dixon, B.K. Speake and F.E. Thacker (1999). Effect of dietary fat supplements on levels of n-3 poly-unsaturated fatty acids, trans acids and conjugated Linoleic acid in bovine milk. J. Anim. Sci. 69:613.
- Ohajuruka, O. A., W. U. Zhiguo and D. L. Palmquist (1991). Ruminal metabolism, fiber and protein digestion by lactating cows fed calcium soap or animal-vegetable fat. J. Dairy Sci. 74: 2601-2609.
- Patton, R. S., C. E. Sorenson and A. R. Hippen (2004). Effects of dietary glucogenic precursors and fat on feed intake and carbohydrate status of transition dairy cows. J. Dairy Sci. 87:2122-2129
- Salfer, J. A., J. G. Linn, D. E. Otterby and W. P. Hansen (1995). Early lactation response of Holstein cows fed a rumen inert fat prepartum, postpartum, or both. J. Dairy Sci., 78: 368.
- Schauff D. J. and J. H. Clark. (1992). Effects of feeding diets containing calcium salts of long-chain fatty acids to lactating dairy cows. J. Dairy Sci., 75: 2990-3002.
- Simas, J. M., J. T. Huber, C. B. Theurer, K. H. Chen, F.A.P. Santos and Z. Wu (1997). Influence of fat source and sorghum grain treatment on performance and digestibilities of high yielding dairy cows. J. Dairy Sci., 80: 2907-2912.
- Simas, J.M., J.T. Huber, C.B. Theurer, K.H. Chen, F.A.P. Santos and Z. Wu (1998). Influence of sorghum grain

- processing on performance and nutrient digestibilities in dairy cows fed varying concentration of fat. *J. Dairy Sci.*, 81: 1966.
- SPSS, (1997). (*Statistical Package for Social Science*) program version 13.0.
- Steele, M. A., O. Al-Zahal, M. E. Walpole and B. W. McBride (2012). Short communication: Grain-induced sub-acute-ruminal acidosis is associated with the differential expression of insulin-like growth factor-binding proteins in rumen papillae of lactating dairy cattle. *J. Dairy Sci.*, 95:6072.
- Sun, Y. and M. Oba (2014). Effects of feeding a high-fiber byproduct feedstuff as a substitute for barley grain on rumen fermentation and productivity of dairy cows in early lactation. *J. Dairy Sci.* 97:1594.
- Talha. M. H. (1996). Nutritional studies on green forage. *Ph. D. Thesis. Ain shams Univ. Cairo Egypt.*
- Weiss, W. P. and D. J. Wyatt (2003). Effect of dietary fat and vitamin E on  $\alpha$ -tocopherol in milk from dairy cows. *J. Dairy sci.*, 86: 3582-3591.
- West, J. W. and G. M. Hill (1990). Effect of a protected fat product on productivity of lactating Holstein and Jersey cows. *J. Dairy sci.*, 73: 3200-3207.
- White, L., D. Bunting, L. S. Sticker, F. G. Hembry and A. M. Saxton (1992). Influence of fish meal and supplemental fat on performance of finishing steers exposed to moderate or high ambient temperatures. *J. Anim. Sci.*, 70: 3286-3292.
- Wu, Z., J. T. Huber, F. T. Sleiman, J. M. Simas, K. H. Chen, S. C. Chan and C. Fontes (1993). Effect of three supplemental fat sources on lactation and digestion in dairy cows. *J. Dairy sci.*, 76: 3562-3570.
- Zened, A., F. Enjalbert, M. C. Nicot and A. Troegeler-Meynadier (2013). Starch plus sunflower oil addition to the diet of dry dairy cows results in a trans-11 to trans-10 shift of bio-hydrogenation. *J. Dairy Sci.* 96:451.
- Zervas, G., K. Fegeros, K. Koysotolis, C. Goulas and A. Mantzios (1998). Soy hulls as a replacement for maize in lactating dairy ewe diets with or without dietary fat supplements. *Anim. Feed Sci. and Tech.* 76: 65.
- Zinn, R. A. (1988). Comparative feeding value of supplemental fat in finishing diets for feed lot steers supplemented with and without monensin. *J. Anim. Sci.* 66: 213.
- Zinn, R. A. (1989). Influence of level and source of dietary fat on its comparative feeding value in finishing diets for steer: feedlot cattle growth and performance. *J. Anim. Sci.* 67: 1029-1037.
- Zinn, R. A. and A. Plascencia (1996). Effects of forage level on the comparative feeding value of supplemental fat in growing finishing diets for feedlot cattle. *J. Anim. Sci.* 74: 1194.

## **تأثير مصادر طاقة الغذاء على الاستفادة من الغذاء والنشاط الميكروبي بكرش الأغنام الأوسيمي**

**بركات محمد أحمد، سعيد سعيد عمر و محمود سمير عبد الموجود ياسين**

قسم الإنتاج الحيواني - كلية الزراعة - جامعة المنوفية

### **الملخص العربي**

أجريت التجربة بهدف دراسة تأثير إضافة مصادر طاقة مختلفة بعلائق الأغنام الأوسيمي على كفاءة الهضم والإتزان النيتروجيني وتخمرات الكرش. أستخدم ثلاثة كباش عمر ٣ سنوات في تصميم تجريبي ٣×٣ مربع لاتيني - المعاملة الأولى عليقة المقارنة مكونة من دريس البرسيم ومخلوط مركزات بينما في الثانية إضيف ٦جم دهن جاف إلى المخلوط المركز وفي الثالثة أزيدت كمية الذرة في مخلوط المركزات بمقدار ٨٥جم. أشارت النتائج إلى أن مصدر الطاقة الغذائية كان له تأثيرا إيجابيا على كميات الغذاء المستهلك وكان التأثير الأكبر لدهن الغذاء على المأكول منه بينما أدت إضافة الذرة إلى زيادة المأكول من المادة الجافة والعضوية والبروتين والكربوهيدرات. تحسنت معاملات الهضم نتيجة إضافة مصادر الطاقة عدا الألياف الخام فقد كانت معاملات هضمها أكبر في العليقة المقارنة. أدت إضافة الطاقة من مصادرها المختلفة إلى تحسن الاتزان النيتروجيني. تحسنت تخمرات الكرش في المعاملة المضاف لها الذرة عن إضافة الدهن. انخفضت قيمة الحموضة بالكرش بعد التغذية لتصل إلى الحد الأدنى عند ساعتين ثم بدأت في الارتفاع ثانية. التغذية على الذرة كمصدر إضافي للطاقة أدى إلى ارتفاع تركيز الأحماض الدهنية الطيارة عن المعاملة بالدهن - وقد تبعت حموضة الكرش الاتجاه المعاكس بينما لم تتأثر تركيزات أمونيا الكرش بنفس الدرجة.