



## A Spotlight on *Retama* spp., The Mediterranean Evergreen Stem-assimilating Xerophyte

Reham A. Youssef<sup>(1)</sup>, Wafaa M. Amer<sup>(1)#</sup>, Omran N. Ghaly<sup>(2)</sup>, Azza B. Hamed<sup>(1)</sup>

<sup>(1)</sup>Department of Botany and Microbiology, Faculty of Science, Cairo University, Giza 12613, Egypt; <sup>(2)</sup>The Herbarium, Desert Research Center (DRC), Cairo, Egypt.



**G**ENUS *Retama* L. (Leguminosae, Tribe Genisteae), is a Mediterranean genus, it naturalized in the new world recently. It includes four (*Retama dasycarpa*, *R. monosperma*, *R. raetam* and *R. sphaerocarpa*) xeromorphic, stem assimilating species. The associated species to the *Retama* spp. are variable and controlled by the spatial distribution and geographic location of each species. The pollen grain of *Retama* spp. is monad, trizonate, some *Retama* spp. flowers are lacking nectar, others are nectar-producing species and insect pollinated. *Retama* spp. utilize a combination of defense mechanisms, acclimatization and adaptation strategies including the stress response genes, physiological and microbiological adaptations to both drought and salinity stresses. Soil and endophytic microbes play an important role in improving establishment of *Retama* species under environmental stresses. Genus *Retama* is eutetraploid with homogeneous karyologic chromosomes, the genome size is an adaptive trait correlated to environmental situations.

This review covered the taxonomy of *Retama* species and their potentialities in arid lands particularly for their role in soil protection, stability, fertility, and desert rehabilitation. The phytochemistry of *Retama* species including flavonoids, essential oil, alkaloids, terpenes, steroids, fatty acids, polysaccharides, and mineral composition was elaborated. In addition to its multipurpose uses as fodder, ethnobotanical uses, and pharmacological activities its active compound possess anticancer, antioxidant, hypoglycemic, anti-inflammatory, analgesic, antileukemic, antiarrhythmic, antibacterial, antifungal activities, and others. This study aimed to highlight the potentialities of the C3 shrubby-leguminous species and its adaptive strategies to the arid-desert stresses, to offer interesting trends to improve the economic crops.

**Keywords.** Assimilating stem, Mediterranean species, *Retama dasycarpa*, *R. monosperma*, *R. raetam*, *R. sphaerocarpa*.

### Introduction

#### Taxonomic description

##### Genus *Retama* Raf.

Genus *Retama* belongs to family Fabaceae; which is one of the most prevalent families in the North Africa of the Mediterranean region (Samala et al., 2023). Unarmed, much branched shrubs (Jafri, 1980; Boulos, 1999) or small trees (Boulos, 1999). Branches erect or ascending, silky-hairy or glabrous, with slender, green (Zohary, 1972), grooved twigs (Zohary, 1972; Jafri, 1980). Leaves simple, small (Jafri, 1980; Boulos, 1999), exstipulate, (Zohary, 1972),

soon deciduous (Jafri, 1980; Boulos, 1999). Flowers white or yellow (Jafri, 1980). Racemes lateral (Boulos, 1999), few- to many flowered (Zohary, 1972; Jafri, 1980). Calyx spathe-like with hemispherical tube (Zohary, 1972) and 5 teeth (Jafri, 1980), ± bilabiate (Jafri, 1980). Corolla much longer than the calyx (Zohary, 1972; Jafri, 1980; Boulos, 1999). Claws of petals adnate to staminal tube. Standard plicate, oblong to obovate-orbicular; wings longer than keel (Zohary, 1972; Jafri, 1980). Stamens 10 Monadelphous (Jafri, 1980), all connate into a closed tube (Zohary, 1972; Boulos, 1999). Ovary sessile or short-stipitate, 2- to few ovules (Zohary,

#Corresponding author email: [wamer@sci.cu.edu.eg](mailto:wamer@sci.cu.edu.eg), <https://orcid.org/0000-0003-0126-6719>

Received 09/07/2023; Accepted 03/01/2024

DOI: 10.21608/ejbo.2024.222012.2402

Edited by: Prof. Dr. Francisco Huerta Martinez, University of Guadalajara, Mexico.

©2024 National Information and Documentation Center (NIDOC)

1972; Jafri, 1980). Style filiform (Zohary, 1972), glabrous. Stigma papillose (Zohary, 1972; Jafri, 1980). Pod indehiscent (Zohary, 1972; Jafri, 1980; Boulos, 1999) or dehiscent (Boulos, 1999), dilated, ovoid to globose (Jafri, 1980), 1- to 2-seeded (Zohary, 1972; Jafri, 1980; Boulos, 1999). Pericarp leathery or fleshy (Zohary, 1972). Seeds spherical (Boulos, 1999) or globular, yellow or brown (Jafri, 1980), radicle thick, less than half as long as cotyledons (Zohary, 1972), without caruncle (Jafri, 1980; Boulos, 1999).

#### *Genus Retama Raf. in Egypt*

Genus *Retama* Raf. in Egypt was subjected to taxonomic revision by Youssef et al. (2023), however, the close morphological similarity between the studied *Retama* taxa hinder their taxonomical identification. Youssef et al. (2023), study was able to resolve several questions concerning genus *Retama* in Egypt among them (1) What are the taxa representing this genus? (2) What are the distinguishing morphological traits? (3) What is the geographic distribution of each taxon? (4) What is molecular fingerprinting/ taxa? The study covered by field and herbarium work using 94 macro-morphological characters and the molecular fingerprinting using ISSR markers to clarify the addressed questions. This revision revealed the identification of two species: *Retama raetam* and *R. monosperma* this is consistent with the records given earlier by Boulos (2009); in addition to five forms (needs complete genome analysis for accurate taxonomic judgement) under *R. raetam* (Form 2, 4, 6, 7, and Form 8). And only Form 5 under *R. monosperma*. The conducted morphological cluster was confirmed with the molecular one, and significant differences were recorded between these taxa. The geographic distribution of the identified *Retama* forms indicates its localization in South Sinai, this is consistent with the postulation given earlier by Täckholm (1974). This study emphasizes that *Retama* spp. are important fodder species, which now susceptible to climatic change and requires urgent conservation effort.

#### *Nomenclature history of the Retama spp.*

Genus *Retama* is belonging to family Leguminosae and Tribe Genisteae. *Retama* species was first termed as *Genista raetam* by Peter Forsskål in 1775. The epithet was named based on the Arabic name “rætæm beham” that was transcribed by Forsskal. *Retama* was given

to this species in the year 1842 by Philip Barker-Webb and Sabin Berthelotthe. In Canary Islands this species named *R. rhodorhizoides*, this name later was known as *R. raetam* (Al-Sharari et al., 2020).

#### *Retama raetam (Forssk.) Webb*

*Common names:* Retem in Arabic, Weeping Broom or White Broom.

*Synonyms:* *Genista raetam* Forssk., *Lygos raetam* (Forssk.) Heywood and *Spartium raetam* (Forssk.) Spach.

*Intraspecific taxon:* According to IUCN, red list (Roskov et al., 2020; Al-Sharari et al., 2020), this species includes two subspecies namely: *R. raetam* subsp. *raetam* and *R. raetam* subsp. *gussonei* (Webb) Greuter (Al-Sharari et al., 2020). While (POWO, 2023) recorded three subspecies: *R. raetam* subsp. *bovei* (Spach) Talavera & P.E. Gibbs, subsp. *raetam* and subsp. *gussonei* (Webb) Greuter (Al-Sharari et al., 2020).

#### *Species description*

##### *Retama dasycarpa* Coss.

*Synonyms:* *Genista dasycarpa* Ball, *Lygos dasycarpa* (Coss.) Jäger

*Retama dasycarpa* is an endemic and native to Morocco particularly in the semi-arid cold bioclimates region of the Elevated Atlas Mountains (Lamrabet et al., 2023). Other references expand its range to include Spain beside Morocco (GBIF, 2020). It grows chiefly in the subtropical biome and in Africa grows in the Mediterranean-Sahara transition zone (POWO, 2023). To the date of issue there is a lack of sufficient morphological description covered this species, based on the herbarium specimens deposited in Kew website (POWO, 2023) and Flora online (WFO, 2023), the species description is as follows: numerous branched shrubs with assimilating stem. Branches ribbed, rush-like, flexible. Leaves are simple, alternate, and soon deciduous. Flowers in short inflorescence with 1-8 flowers. Fruit one seeded pod.

##### *R. monosperma* (L.) Boiss.

*Synonyms:* *Genista monosperma* (L.) Lam., *Lygos monosperma* (L.) Heywood, *Retama webbii* (Spach) Webb, *Spartium monospermum* L., *S. clusii* Spach (Muñoz Vallés et al., 2013).


*Retama monosperma* is shrub up to 4.5 m high and have numerous branching green stems (Quezel & Sant, 1962; Herrera, 2001; Muñoz Vallés et al., 2013). Branches are unarmed, juvenile branches silvery-sericeous and glabrescent with age (Herrera, 2001; Muñoz Vallés et al., 2013). Branches pendent (Tutin et al., 1968), ribbed, rush-like, silky-haired, flexible (Polunin, 1969). Leaves are simple, alternate, stipulate, very shortly pedunculated, sericeous to pubescent, deciduous. Leaf lamina is 4–8 mm x 0.7–1mm, lanceolate or oblanceolate to linear-lanceolate, pubescent allover (Tutin et al., 1968; Herrera, 2001; Muñoz Vallés et al., 2013), apex obtuse (Barker-Webb & Berthelot, 1836). Stipules narrow, acute, absent or caducous, inconspicuous (Barker-Webb & Berthelot, 1836). Inflorescences are lax axillary racemes, with 10–26 flowers (Tutin et al., 1968; Herrera, 2001; Muñoz Vallés et al., 2013). Bracts and bracteoles are 2.5–2mm x 0.7–1.7mm, sericeous, caducous. Flowers zygomorphic, 14-15mm (Quezel & Sant, 1962). Calyx length 3–4mm, campanulate-cylindrical, glabrous, reddish, nearly bilabiate caducous after anthesis (Tutin et al., 1968). Corolla clawed white, 9–13 mm long, papilionate, zygomorphic. Corolla white, papilionaceous, standard rhombic-ovate (Webb, 1853), shorter than the wing (Quezel & Sant, 1962) and the keel (Polunin, 1969), or wing as long as or shorter than the keel (Tutin et al., 1968), Keel about half shorter than the wings, lanceolate to ovate, recurved (Barker-Webb & Berthelot, 1836), apex cuspidate -acuminate (Tutin et al., 1968). Androecium monadelphous, 10 stamens united below with 4- short stamens with basifixed anthers, and 5- medium stamens with dorsifixed anthers, and one larger basifixed anthers (Herrera, 2001; Muñoz Vallés et al., 2013). Ovary not pedunculated, glabrous, with four to seven seminal rudiments (Herrera, 2001; Muñoz Vallés et al., 2013), ovate (Barker-Webb & Berthelot, 1836), unilocular, indehiscent, leathery, shiny, wrinkled or reticulate, exerted from calyx, short mucronate directed towards ventral suture style terete, stigma capitate, forward (Webb, 1853) or backward sloping (Barker-Webb & Berthelot, 1836). Pericarp is fleshy and sugary when it ripens. Fruit legume, 10-16mm (Tutin et al., 1968; Polunin, 1969), 10–22mm single seed (rarely 2–3), more or less globose (Herrera, 2001; Muñoz Vallés et al., 2013), with dilated ventral suture (Quezel & Sant, 1962). Flowering from January to June (Domínguez, 1987; Muñoz Vallés et al., 2013). Flowering from February to May (Jahandiez &

Maire, 1932; Polunin, 1969) and fruiting from late May to early September (Muñoz Vallés et al., 2013).

*R. raetam* (Forssk.) Webb

Synonyms: *Genista raetam* Forssk., *Lygos raetam* (Forssk.) Heywood, *Retama duriaei* (Spach) Webb, *R. raetam* subsp. *raetam*, *R. raetam* is a North African shrub up to 3.0m (Zohary, 1972; Benhouhou, 2005; Al-Sharari et al., 2020), 0.5-2.0m (Boulos, 1999), and may reach up to 6.0m (Al-Sharari et al., 2020), spartoid shrub with furrowed stem (Zohary, 1972; Boulos, 1999). Richly branching (Boulos, 1999), ascending spreading branches (Barker-Webb & Berthelot, 1836), mainly vertical and yellowish-green (Zohary, 1972). Young stems covered with silky-silvery sparse to dense trichomes (Boulos, 1999), become glabrous with age (Al-Sharari et al., 2020). Leaves 1.0-23.0 x 0.25- 3.0 (8.0) mm (Zohary, 1972; Jafri, 1980); soon deciduous; narrow elliptical (Al-Sharari et al., 2020), alternate (Ahmed et al., 2016; Al-Sharari et al., 2020), simple, subsessile (Ahmed et al., 2016), linear to lyrate or lanceolate (Barker-Webb & Berthelot, 1836), covered with dense trichomes; apex acute-obtuse; margin entire (Boulos, 1999). Inflorescences are dense racemes (Ahmed et al., 2016; Al-Sharari et al., 2020), raceme lax-dense 1.0 to 15- flowered (Jafri, 1980). Flower bisexual, in clusters of 3 – 15 flowers (Al-Sharari et al., 2020), flower 8 -10cm (Fig. 1 & Al-Sharari et al., 2020), 7-16mm, becoming creamy-colored on drying (Jafri, 1980), zygomorphic and short pedicellate (Ahmed et al., 2016). Flower pentamerous, 5-gamosepalous, caducous after anthesis, pale to dark purple occasionally with purple spots (Tutin et al., 1968). Corolla 5-papilionate, standard white or white with purplish tips or spots (Boulos, 1999; Ahmed et al., 2016) or with one purple main vein (Barker-Webb & Berthelot, 1836). Standard 7-15 x 5-11mm, equal or longer (Boulos, 1999), occasionally shorter than the wings, elliptical to orbicular (Zohary, 1972; Jafri, 1980) or ovate to oblong (Tutin et al., 1968), clawed and arched (Barker-Webb & Berthelot, 1836). Wings 7-15 x 1.5-4mm, acinaciform (Barker-Webb & Berthelot, 1836) or oblong with one main vein, obtuse apex. Keel 7-13 x 3-5mm, somewhat shorter than wings (Zohary, 1972), lanceolate (Barker-Webb & Berthelot, 1836) or oblong, apex obtuse or acute, densely hairy along occasionally with purple mid main vein (Zohary,

1972; Jafri, 1980). Stamens 10, monadelphous, connate below into a closed tube, 6-13mm, nearly equal in lengths, random in lengths or 1 long and the rest nearly equal (Boulos, 1999). Ovary superior, sessile, with one ovule with capitate stigma (Ahmed et al., 2016). Fruit indehiscent legume (Ahmed et al., 2016) or tardily dehiscent (Zohary, 1972), 10–15mm diameter (Al-Sharari et al., 2020), 3.0-10mm (Zohary, 1972; Jafri, 1980), green or black (Barker-Webb & Berthelot, 1836), ventral suture filiform to dilated, glabrous, elliptical-ovoid-globose (Zohary, 1972) or to oblong (Barker-Webb & Berthelot, 1836; Jafri, 1980), legume apex attenuate into a gradually or abruptly short erect or curved beak (Zohary, 1972). Fruit includes 1-2 kidney-shaped, yellow to brown seeds (Al-Sharari et al., 2020). Pericarp thin, smooth or wrinkled, leathery or fleshy horny (Täckholm, 1974), smooth, glabrous, yellow to orange, olive green to brown or black (Tutin



**Fig. 1.** Field image for *Retama* species in Egypt: A & B, *R. raetam* in Wadi Hagoul, Eastern desert; C & D, *R. monosperma* in sand dune at the western Mediterranean stripe, near Marsa Matruh. (Photograph by Dr. Omran Galy).

et al., 1968).  $2n=48$  (Zohary, 1972). Flowering from February up to May (Al-Sharari et al., 2020), from end of January to April (Zohary, 1972).

*R. raetam* subsp. *gussonei* (Webb) Greuter: Low shrub (Täckholm, 1974) (Fig. 1), diffusely branching, with silky-silvery indumentum and deflexed upper branches (Zohary, 1972). Flowers as in type, but the pod is dehiscent (Täckholm, 1974) or tardily dehiscent. Fruit short ellipsoidal or ovoid, with more or less abrupt, short, straight or recurved beak (Zohary, 1972). Fruit pericarp fleshy, shiny, wrinkled. Seed lemon to yellow, rarely green (Zohary, 1972).

*R. sphaerocarpa* (L.) Boiss.

Synonyms: *Boelia sphaerocarpa* (L.) Webb, *Lygos sphaerocarpa* (L.) Heywood, *Retama atlantica* Pomel, *R. lutea* Raf., *R. sphaerocarpa* f. *atlantica* (Pomel) Batt, *Spartium sphaerocarpum* L.

*R. sphaerocarpa* is a leguminous shrub with evergreen assimilating stems, up to 4.0m height (Haase et al., 1996; Prieto et al., 2010), to 5.0m in height (Rodríguez-Echeverría & Pérez-Fernández, 2003). It possesses a dimorphic root system, with root system grows up to 30 m deep, with long lateral roots and deep roots (Haase et al., 1996; Prieto et al., 2010). Flower yellow papilionaceous corolla and bilabiate calyx (Rodríguez-Riaño et al., 1999c). Androecium is 10 dimorphic anthers, monadelphous, the upper whorl with oblong to linear-oblong basifixed stamens, and the lower whorl with similar morphology but the apex is a triangular-acuminate and the anther is dorsifixed (López et al., 1999; Rodríguez-Riaño et al., 1999c). Ovary is glabrous with a linear-curved style terminated with tufted stigma. Fruit indehiscent legume, single seeded. Plant flowering in spring (Rodríguez-Riaño et al., 1999b, c).

#### Seed morphology in *Retama* spp.

Different colors were reported for *R. raetam* seed among them yellow-brown (Zohary, 1972; Jafri, 1980), black (Barker-Webb & Berthelot, 1836; Webb, 1853), brownish (Boulos, 1999) and brown-black (Täckholm, 1974). While seeds of the *R. monosperma* are reniform, with dark-green to black or smooth integument (Barker-Webb & Berthelot, 1836; Webb, 1853; Morsy et al., 2015) or reddish-brown or olive green (Boulos, 1999). The seed dimensions are 5.7–7.2mm x 4.2–5.7mm, and  $93.1 \pm 2.1$ mg seed mean weight (Morsy et al., 2015). The *R. raetam* subsp. *gussonei* seeds are sub-reniform with uniform feature and smooth integument (Ferrauto et al., 2015). Different colors were reported by earlier authors, among them lemon yellow to yellow color (Zohary, 1972; Täckholm, 1974), rarely green (Zohary, 1972), and varying in color from olive-green to golden-yellow, occasionally brown spotted (Ferrauto et al., 2015). Mean seed size is  $5.94 (\pm 0.72)$  mm x  $5.70 (\pm 0.67)$  mm with classical embryo with two cotyledons (Ferrauto et al., 2015).

*Seeds of Retama spp.: production, dispersion and germination*

*Retama monosperma* possesses one- rarely two seeds/fruit. The seed production/shrub varied from year to other, ranged from 335 - 2800 fruits/m<sup>2</sup> beneath canopy of the full mature shrubs of 6-7 years old (Muñoz Vallés et al., 2013). Seeds of this species lack the special mechanism for seed dispersal, seeds fall down under the shrub canopy and become available for dispersers or stay in the soil for the coming season (Muñoz Vallés et al., 2013). Wild rabbit (*Oryctolagus cuniculus*) is among the active important dispersers of *R. monosperma* seeds (Dellafiore et al., 2006; Muñoz Vallés et al., 2013). On the other hand, seeds consumed as goat fodder and this cause loss of seeds and no record about the bird disperser while, rain runoff sometimes act as seeds dispersers (Muñoz Vallés et al., 2013). The seedling pairing two fleshy cotyledons, while the juvenile leaves are alternate covered with trichomes from both sides even in mature shoots (Muñoz Vallés et al., 2013). *R. monosperma* stabilizes the coastal sand forming dunes of medium- high levels which limits the seedling establishment (Muñoz Vallés et al., 2013). Despite lacking vegetative reproduction in this species, the populations expand and become denser due to the effective dispersal mechanisms and high seed productivity (Fernández et al., 2010; Muñoz Vallés et al., 2013). A comparable effect has been reported in *R. raetam* forming nebkhas (accumulations of wind-borne sand) in Mediterranean coastal dunes around the shrub canopy (El-Bana et al., 2002; Muñoz Vallés et al. 2013). Abdenour et al. (2020) reported that *R. sphaerocarpa*, *R. raetam* and *R. monosperma* stabilize the dunes, leading to soil fixation and the rebuilding of the plant cover of semi- and arid ecosystems (Muñoz Vallés et al., 2013; Kheloufi et al., 2020).

The *R. monosperma* seeds germinate in autumn throughout the rainy season, the rate of germination increases significantly after animals feeds on seeds, where the wild rabbits rise up this percentage to 24% (Dellafiore et al., 2006; Muñoz Vallés et al., 2013). Seed viability is very high (90.0–98.8%), despite this its germination rates were ranged from 5 to 13% (Dellafiore et al., 2006; Dellafiore, 2007; Muñoz Vallés et al., 2013). The germination percentage in *Retama* sp. is caused by the seed coat impermeability (Kigel, 1995; Ferrauto et al., 2015).

The earlier reports about the germination of *R. raetam* ssp. *gussonei* (Gutterman, 1993; Izhaki & Ne'eman, 1997; Ferrauto et al., 2015; Seglie et al., 2012) showed very low of non-uniform germination owing to the physical dormancy, due to the seed coat impermeability to water (Kigel, 1995; Ferrauto et al., 2015). To improve the germination rate, various treatments have been applied to defeat dormancy. From these methods the acid scarification is the most common and effective method through which the seeds immersed in sulphuric acid for 3h at 25°C. The achieved maximum germination rate was 67.5% ± 22.17 with germination rate (T50 = 5.6 ± 1.2 & MGT = 8.6 ± 0.8 days). However, the manual scarification followed by boiling reduces the integument resistance, the method fails to affect the *Retama* seed dormancy (Youssef, 2009; Ferrauto et al., 2015). Relevant results were achieved in case of *R. sphaerocarpa*, where its seeds dormancy can be overcomes by immersion several methods as boiling water, conc. sulphuric acid (0min - 240min) and seed coat abrasion by sandpaper (Kildisheva, 2019; Kheloufi et al., 2020). The best of all these pretreatments was immersion in sulphuric acid for 120 min at temperature 25°C (± 2°C), this resulting 86% seed germination (Kheloufi et al., 2020).

Biological treatment can improve seed germination as a non-conventional method (Steponkus, 1982; Ouf, 1994). Microorganisms may promote germination, probably by enhancing the seed coat softening (Guttridge et al., 1984; Ouf, 1994). Referring to the fungal hydrolytic enzymes produced during biodegradation of plant materials (Ouf, 1994). Bathing of *R. raetam* seeds in the filtrate of the fungal growth media as *Penicillium capsulatum*, *P. spinulosum* and *Sporotrichum pulverulentum* could stimulate the seed germination (Ouf, 1994).

*Factors influencing seed germination of Retama spp.*

**Salinity:** Seeds of *R. raetam* are sensitive to salinity, it tolerates from 34 to 68 mM NaCl concentrations. The germination rate dropped to 3.33% on 238 and 272mM of NaCl concentrations with 272 mM threshold of tolerance (Mehdadi et al., 2017). *R. raetam* subsp. *bovei* seeds were also negatively correlated to the medium salinity. It can tolerate salinity up to 15g/L. Where germination reached up to 81% in distilled water and significantly decreased with increase in NaCl

concentrations, similar behavior was reported with the increase in osmotic potential (Mechergui et al., 2017). The germination inhibition cannot recover after the toxicity action caused by the high  $\text{Na}^+$  and  $\text{Cl}^-$  ions concentrations, even after the removal of the higher concentrations. Contradictory behavior was noticed, and germination inhibition was reversible in case of germination inhibition related to osmotic potential (Mehdadi et al., 2017). In *R. raetam* subsp. *bovei* the seed weight plays a major factor in stimulating germination under salinity stress and the germination inhibition has notable impact on the seeds with small size (Mechergui et al., 2017).

**Temperature:** The germination of *R. raetam* seeds was optimum (70%) at the temperature range from 20°C - 25°C, while germination was impossible at the higher (35°C - 40°C) or lower (0°C, 5°C) temperature ranges (Mehdadi et al., 2017). While, in *R. monosperma* the optimal germination temperature was 20°C, and germination was impossible at the same temperatures range as in *R. raetam* (Bouredja et al., 2011; Mehdadi et al., 2017).

#### *Distribution of genus Retama spp.*

Genus *Retama* L. is mainly with Mediterranean distribution, the western part is its diversity center, particularly in NW Africa (Lems, 1960; Maire, 1987; Greuter et al., 1989; Chiapella et al., 2009). *Retama* spp. belongs to the Mediterranean endemic species, where 60% of the endemic species of this region possess narrow geographic location (Bedair et al., 2024).

*R. dasycarpa* Cosson is an endemic species to the Great Atlas and Morocco (Maire, 1987; Chiapella et al., 2009).

*R. monosperma* distributed in the Algeria and Morocco; SW Iberian Peninsula, var. *webbii* (Spach) Maire grows along the Atlantic coast of Morocco (Maire, 1987; Talavera, 1999b; Chiapella et al., 2009). It is endemic to the NW Morocco (Talavera, 1999b; Zunzunegui et al., 2017) and SW Iberian Peninsula on coastal sandy dunes and borders of saltmarshes (Muñoz Vallés et al., 2013; Zunzunegui et al., 2017). And expanded to SW Spain and in areas with Mediterranean climate in California, USA (Randall, 1997; Zunzunegui et al., 2017), and S. Australia (Randall, 2007; Zunzunegui et al., 2017).

*R. raetam* (Forssk.) Webb extends eastward to Sinai Peninsula, NW Saudi Arabia from Al-Jouf to Tabuk, and Palestine (Al-Tubuly et al., 2011; POWO, 2023; Al-Sharari et al., 2020). Recently, it naturalized in new places, including Queensland, Greece, S & W Australia (POWO, 2023; Al-Sharari et al., 2020). While *Retama raetam* (Forsskål) Webb subsp. *raetam*, is a Saharo-Arabian element, spreads in N. Africa, Arabian Peninsula, Israel, Syria and Lebanon (Chiapella et al., 2009), along the Mediterranean coast and arid deserts of Egypt (El-Bahri et al., 1999; Mittler et al., 2001; Barakat et al., 2013), and extends up to Sicilia (Al-Sharari et al., 2020).

*R. sphaerocarpa* (L.) Boiss. is distributed in the Iberian Peninsula, Morocco, Algeria and Tunisia (Chiapella et al., 2009), native to NW Africa and Iberian Peninsula (Tutin et al., 1968; Rodríguez-Echeverría & Pérez-Fernández, 2003); and C & S Spain (Rodríguez-Echeverría & Pérez-Fernández, 2003).

#### *Soil type affecting the distribution of Retama species*

The type of soil plays an important role in distribution of *R. monosperma* (Gómez González et al., 2004; Muñoz Vallés et al., 2013). It is a drought tolerant species and can modify the habitat of the coastal dunes and plant community, it can grow in the geographic areas possess Mediterranean climate (Muñoz Vallés et al., 2013). Generally, it grows in siliceous coastal soil with grain size 0.2–0.3mm and sandy soil with low organic and water content (Muñoz Vallés et al., 2013). It grows in wide pH 5.8 to 9.0 range (García Novo & Merino, 1997; Muñoz Vallés et al., 2011, 2013).

#### *Cytogenetics of Retama spp.*

Genus *Retama* is eutetraploid with homogeneous karyologic chromosomes. The basis of serologic and morphologic characters showed a greater similarity to Genista group (Polhill, 1976; Bisby, 1981; Cristofolini & Feoli Chiapella, 1984; Feoli Chiapella & Prodan, 1989; Chiapella et al., 2009). The cytogenetics study carried out by Benmiloud-Mahieddine et al. (2011) on 33 population from the three *Retama* spp. in Algeria namely *R. sphaerocarpa*, *R. monosperma* and *R. raetam*, using flow cytometry and molecular cytogenetics for karyotype analysis revealed that the three species have the similar chromosome number ( $2n=48$ ). Also, (Chiapella

et al., 2009) study four genera of Genisteae and revealed similar chromosome numbers ( $x = 12$  &  $2n=48$ ; with some B chromosomes up to 6 in some species). Where, *R. monosperma*,  $2n= 48 + 4B$ , *R. sphaerocarpa*  $2n = 48 + 2B$  and *R. dasycarpa* was  $2n = 48 + 3B$  (Chiapella et al., 2009). The numbers  $2n = 24$  &  $52$  are only reported respectively for *R. raetam* subsp. *raetam* and *R. sphaerocarpa* (Chiapella et al., 2009), while *R. monosperma* (L.) Boiss. was of  $2n = 24$  &  $48$  (Muñoz Vallés et al., 2013). According to Gallego-Martin et al. (1988) and Benmiloud-Mahieddine et al. (2011), *Retama* species has two ancestral basic chromosome numbers ( $x=6$  and  $x=8$ ) with the possible polyploidy (secondary or series) of aneuploids and euploids (Gallego-Martin et al., 1988; Benmiloud-Mahieddine et al., 2011).

Genome size is among the useful traits applied for evolution and systematics studies (Godelle et al., 1993; Cerbah et al., 1999a & b, 2001; Ohri, 1998; Benmiloud-Mahieddine et al., 2011). Flow cytometry is a rapid and accurate technique used to study the genome size to assess the inter- & intraspecific variations (Biradar & Rayburn, 1993; Doležel & Bartoš, 2005; Benmiloud-Mahieddine et al., 2011). The study carried out by Benmiloud-Mahieddine et al. (2011) revealed that the chromosome structure and genome size distinguished *R. sphaerocarpa* from *R. monosperma* and *R. raetam*.

The genome size for *Retama monosperma*, *R. raetam* and *R. sphaerocarpa*, was 1.76 - 1.89 pg, 1.88 - 1.93 pg and 1.80 pg - 1.97 pg; respectively. The genome size showed significant intraspecific difference connected to the spatial distribution of the studied populations, and also approved by rDNA organization in chromosome analysis (Benmiloud-Mahieddine et al., 2011). The genome size is an adaptive trait correlated to the environmental situations (Bennett, 1987; Benmiloud-Mahieddine et al., 2011). Benmiloud-Mahieddine et al. (2011) reported that the *R. raetam* population collected from the lowest temperatures at the high plateau possess the highest nuclear DNA content ( $2C= 1.97$  pg). While the smallest genome size ( $2C=1.80$  pg) was specific for the population located in the Algerian mountains, desert at 552 and 456 m. The two Agrobacterium strains, described by Benmiloud-Mahieddine et al. (2011) were co-localized on the satellite chromosome pair distinguished *R. raetam* and *R. monosperma* from *R. sphaerocarpa*.

#### *Anatomy of the assimilating branch in Retama raetam*

*R. raetam* is a xeromorphic species with soon deciduous leaves, and the assimilating branches appeared with longitudinal grooved in which the anomocytic stomata are located with reticulate cell wall (Ahmed et al., 2016). The deeply placed stomata have cuticular crest from both sides enclosing very small respiratory cavity; the number of stomata per unit area in this plant is impossible to estimate (Evenari, 1938). The transverse section in assimilating branch showed the presence of very thick epidermal cells (20-25u). The assimilation parenchyma was characterized by a few intercellular spaces. The conducting system is an advantageous in its construction, its tracheids are small number wide lumen (Evenari, 1938).

#### *Pollen morphology of Retama raetam*

The pollen grain of *R. raetam* is monad, trizono-colporate, isoplar, triangular-obtuse in polar view and rectangular-obtuse in equatorial view. The grain size  $> 20\mu\text{m}$ , while the exine sculpture is supra-reticulate (Sekina & Moore, 1995).

*Pollen morphology of R. raetam subsp. gussonei*: As seen in Fig. 2, the pollen of *R. raetam* subsp. *gussonei* is monad, trizono-colporate (sometimes trizono-colporoidate), radially symmetric, and its size ranges from small - medium ( $P= 23-33\mu\text{m}$ ). Subcircular in polar view, while, in equatorial view it is subtriangular-elliptic. The polar axis ranges from 23 - 33  $\mu\text{m}$  and the equatorial diameter 17 - 27  $\mu\text{m}$ ; the pollen aperture is ora and colpi. A longitudinal parallel edge supporting the colpus, and often constricted at the equator (De Leonardis & Zizza, 1994; Ferrauto et al., 2015). Exine tectate columellate, with same thickness nexina and sexina. Tectum ornamentation is




Fig. 2. Pollen grains of *R. raetam* ssp. *gussonei*. using SEM: A, Equatorial view; B, Polar views; C, Magnified exine. Scale bar: A-C = 2mm. (after Ferrauto et al., 2015).

supra-reticulate under light microscope, while appeared, supramicroreticulate-perforate under the scanning electron microscope (Ghirardelli et al., 1997; Rizzi Longo et al., 2006; Ferrauto et al., 2015). The mesocolpium showed fossulate-perforate ornamentation with perforation more than one (Pardo et al., 2000; Ferrauto et al., 2015), while the aperture membrane regulate ornamentation (Ferrauto et al., 2015).

#### Pollination Biology

*Retama monosperma*: Nevertheless, the monadelphous flowers are normally lacking nectar, *R. monosperma* is an exception and nectar producing species. The flowers during pollination were visited mostly by *Hymenoptera* (Herrera, 1988), and others species from other groups like *Licaenidae* and *Syrphydae* (Muñoz Vallés et al., 2013). This species is of low biomass of pollen grain production (mean 92 dmm<sup>3</sup> / flower), where this value reached 624.3 ± 923.3 dmm<sup>3</sup>/ flower in the whole *Retama* tribe (Rodríguez-Riaño et al., 1999a; Muñoz Vallés et al., 2013).

*R. sphaerocarpa*: Rodríguez-Riaño et al. (1999c), studied the reproductive biology of *R. sphaerocarpa* (L.) Boiss. The results revealed that this species is a nectar producing species, the nectar gland is an extra-staminal observed between the staminal column and calyx its nectar

production is maximum at early morning. Where the pollinators are primarily *Apis mellifera* and minor beetle (*Heliotaurus* sp.) which made 14 visits/2 min to the flowers of the same or neighbor individuals. Honeybees can pollinate *R. sphaerocarpa*, flowers in 4-9 seconds. The genetic self-incompatibility (SI) is the rejection of male gametophyte by ovule when it carries the same alleles of the same plant. The interaction between pollen and pistil in the studied species revealed the presence of prezygotic SI- of gametophytic type. The hand-self-pollination produced very few fruits, indicating the presence of postzygotic rejection mechanism. On the other hand, the hand-cross pollination increased the fruit formation if compared to the control reflecting the pollen grains were the limiting factor in reproduction process. Rodríguez-Riaño et al. (1999c), reported that the low nectar production could be linked with the small flower-size. The less nectar production is linked with the ability of pollen grains to farther distances than the pollen produced from rich nectar producing plants (Zimmermann, 1988; Rodríguez-Riaño et al., 1999c).

*Species associated to R. raetam and R. monosperma*

*Species associated to R. monosperma in Spain*: *R. monosperma* colonized SW Spain and dominating the psammophyte vegetation was co-dominated by *Anisantha rigida*, *Artemisia*

TABLE 1. Species associated to *R. monosperma* and *R. raetam* in Egypt (Based on field collection carried out by Wafaa Amer)

Species associated to <i>Retama monosperma</i> (L.) Boiss		
<b>Family: Amaranthaceae</b>	<b>Family: Boraginaceae</b>	<b>Family: Plumbaginaceae</b>
• <i>Anabasis oropedioides</i> Maire	• <i>Echium angustifolium</i> Mill.	• <i>Limonium tubiflorum</i> (Delile) Kuntze
• <i>Atriplex halimus</i> L	<b>Family: Brassicaceae</b>	<b>Family: Poaceae</b>
<b>Family: Apiaceae</b>	• <i>Moricandia nitens</i> (Viv.) E.A.Durand & Barratte	• <i>Aegilops kotschyi</i> Boiss.
• <i>Deverra tortuosa</i> (Desf.) DC.	<b>Family: Fabaceae</b>	• <i>Hordeum murinum</i> L.
• <i>Eryngium creticum</i> Lam.	• <i>Acacia tortilis</i> subsp. <i>raddiana</i> (Savi) Brenan	• <i>Lygeum spartum</i> Loeff. ex L.
<b>Family: Aspragraceae</b>	• <i>Prosopis farcta</i> (Banks & Sol.) J.F.Macbr.	<b>Family: Solanaceae</b>
• <i>Asparagus horridus</i> L.	<b>Family: Lamiaceae</b>	• <i>Lycium europaeum</i> L.
<b>Family: Asteraceae</b>	• <i>Marrubium alysson</i> L.	<b>Family: Thymelaeaceae</b>
• <i>Achillea tenuifolia</i> Lam.	• <i>Salvia lanigera</i> Poir.	• <i>Thymelaea hirsuta</i> (L.) Endl.
• <i>Atractylis carduus</i> (Forssk.) C. Chr.	<b>Family: Nitrariaceae</b>	<b>Family: Xanthorrhoeaceae</b>
• <i>Calendula arvensis</i> L.	• <i>Peganum harmala</i> L.	• <i>Asphodelus aestivus</i> Brot.
• <i>Carthamus mareoticus</i> Delile	<b>Family: Plantaginaceae</b>	<b>Family: Zygophyllaceae</b>
• <i>Centaurea alexandrina</i> Delile	• <i>Plantago crypsoides</i> Boiss.	• <i>Fagonia indica</i> Burm.f.
• <i>Centaurea glomerata</i> Vahl		• <i>Tetraena alba</i> (L.f.) Beier & Thulin
• <i>Echinops spinosissimus</i> Turra		

TABLE 1. Cont.

Species associated to <i>Retama raetam</i> (Forssk.) Webb.		
<b>Family: Aizoaceae</b>	• <i>Echium angustifolium</i> Mill.	<b>Family: Poaceae</b>
• <i>Mesembryanthemum crystallinum</i> L.	• <i>Echium angustifolium</i> subsp. <i>sericeum</i> (Vahl) Klotz.	• <i>Aegilops kotschy</i> Boiss.
• <i>Mesembryanthemum nodiflorum</i> L.	<b>Family: Brassicaceae</b>	• <i>Ammophila arenaria</i> (L.) Link
<b>Family: Amaranthaceae</b>	• <i>Cakile maritima</i> Scop.	• <i>Elymus farctus</i> (Viv.) Runemark ex Melderis
• <i>Aerva javanica</i> (Burm.f.) Juss. ex Schult.	• <i>Diploaxis acris</i> (Forssk.) Boiss.	• <i>Hordeum marinum</i> Huds.
• <i>Anabasis setifera</i> Moq.	<b>Family: Capparaceae</b>	• <i>Lygeum spartum</i> Loefl. ex L.
• <i>Arthrocnemum macrostachyum</i> (Moric.) K.Koch	• <i>Capparis aegyptia</i> Lam.	• <i>Panicum turgidum</i> Forssk.
• <i>Atriplex halimus</i> L.	<b>Family: Caryophyllaceae</b>	<b>Family: Polygonaceae</b>
• <i>Haloxylon salicornicum</i> (Moq.) Bunge ex Boiss.	• <i>Gymnocarpus decandrus</i> Forssk.	• <i>Rumex vesicarius</i> L.
<b>Family: Apiaceae</b>	• <i>Gypsophila capillaris</i> (Forssk.) C.Chr.	<b>Family: Posidonaceae</b>
• <i>Deverra tortuosa</i> (Desf.) DC.	<b>Family: Cistaceae</b>	• <i>Posidonia oceanica</i> (L.) Delile
• <i>Eryngium creticum</i> Lam.	• <i>Helianthemum lippii</i> (L.) Dum.Cours.	<b>Family: Resedaceae</b>
<b>Family: Apocynaceae</b>	<b>Family: Cleomaceae</b>	• <i>Ochradenus baccatus</i> Delile
• <i>Calotropis procera</i> (Aiton) W.T. Aiton	• <i>Cleome amblyocarpa</i> Barratte & Murb.	• <i>Reseda decursiva</i> Forssk.
• <i>Leptadenia pyrotechnica</i> (Forssk.) Decne.	• <i>Farsetia aegyptia</i> Turra	<b>Family: Rubiaceae</b>
<b>Family: Asteraceae</b>	• <i>Zilla spinosa</i> (L.) Prantl	• <i>Crucianella maritima</i> L.
• <i>Achillea tenuifolia</i> Lam.	<b>Family: Euphorbiaceae</b>	<b>Family: Scrophulariaceae</b>
• <i>Atractylis carduus</i> (Forssk.) C.Chr.	• <i>Euphorbia paralias</i> L.	• <i>Verbascum letourneuxii</i> Asch. & Schweinf.
• <i>Centaurea alexandrina</i> Delile	<b>Family: Fabaceae</b>	<b>Family: Solanaceae</b>
• <i>Centaurea calcitrapa</i> L.	• <i>Acacia tortilis</i> subsp. <i>raddiana</i> (Savi) Brenan	• <i>Hyoscyamus muticus</i> L.
• <i>Centaurea glomerata</i> Vahl	• <i>Astragalus spinosus</i> (Forssk.) Muschl.	• <i>Lycium europaeum</i> L.
• <i>Echinops spinosissimus</i> Turra	• <i>Crotalaria aegyptiaca</i> Benth.	• <i>Lycium shawii</i> Roem. & Schult.
• <i>Iphiona mucronata</i> (Forssk.) Asch. & Schweinf.	• <i>Lotus arabicus</i> L.	<b>Family: Tamaricaceae</b>
• <i>Launaea nudicaulis</i> (L.) Hook.f.	• <i>Lotus polyphyllus</i> Clarke	• <i>Tamarix senegalensis</i> DC.
• <i>Launaea spinosa</i> (Forssk.) Sch.Bip. ex Kuntze	<b>Family: Geraniaceae</b>	<b>Family: Thymelaeaceae</b>
• <i>Otanthus maritimus</i> (L.) Hoffmanns. & Link	• <i>Erodium oxyrhinchum</i> subsp. <i>bryoniiifolium</i> (Boiss.) Schön.-Tem.	• <i>Thymelaea hirsuta</i> (L.) Endl.
• <i>Pulicaria undulata</i> (L.) C.A.Mey.	<b>Family: Lamiaceae</b>	<b>Family: Xanthorrhoeaceae</b>
• <i>Senecio glaucus</i> subsp. <i>coronopifolius</i> (Maire) C.Alexander	• <i>Lavandula coronopifolia</i> Poir.	• <i>Asphodelus tenuifolius</i> Cav.
<b>Family: Boraginaceae</b>	• <i>Ononis vaginalis</i> M.Vahl	<b>Family: Zygophyllaceae</b>
• <i>Anchusa humilis</i> (Desf.) I. M. Johnst.	• <i>Trigonella stellata</i> Forssk.	• <i>Fagonia arabica</i> L.
• <i>Echiochilon fruticosum</i> Desf.	<b>Family: Plantaginaceae</b>	• <i>Fagonia cretica</i> L.
	• <i>Plantago coronopus</i> L.	• <i>Fagonia mollis</i> Delile
	• <i>Plantago crypsoides</i> Boiss.	• <i>Tetraena alba</i> (L.f.) Beier & Thulin
	• <i>Plantago notata</i> Lag.	• <i>Tetraena coccinea</i> (L.) Beier & Thulin
	<b>Family: Plumbaginaceae</b>	• <i>Tetraena simplex</i> (L.) Beier & Thulin
	• <i>Limonium tubiflorum</i> (Delile) Kuntze	• <i>Zygophyllum decumbens</i> Delile

*campestris Maritima*, *Carduus meoanthus*, *Crucianella maritima*, *Erodium cicutarium*, *Euphorbia terracina*, *Hedypnois arenaria*, *Helichrysum italicum Picardii*, *Malcolmia littorea*, *Medicago littoralis*, *Paronychia argentea*, *Silene nicaeensis*, *Sonchus tenerrimus*, *Stellaria pallida*, *Thymus carnosus*, *Urtica membranacea*, and *Vulpia alopecuros* (Table 1. Muñoz Vallés et al., 2013).

#### *Adaptation to the environmental stresses*

##### *Climatic factors*

Climatic factors are the main player out of all the environmental factors that restraint not only the plant growth but extends to include development, densities, distribution and the vegetation abundance all over the globe (Cramer et al., 2001; Booth & Grime, 2003; Scholze et al., 2006; Amer et al., 2021). Amer et al. (2021) studied the genetic variations induced by the climate change (mainly air temperature and rainfall) over the period 1975–2018 on representative Mediterranean species in the western Mediterranean stripe in Egypt. The *R. raetam* and *Vicia monantha* were representing the shrubby and annual Mediterranean species; respectively. The genetic variations were carried out using ISSR-technique, the study proved remarkable genome variability in *R. raetam* than the annual *Vicia monantha* with 77% and 91%; respectively. The study announces for the species vulnerability to climate change inducing the unidirectional genetic change in *R. raetam* shrubs; and serious conservation plan is requested (Amer et al. 2021).

##### *Drought adaptation of R. raetam*

The species utilizes a combination of numerous adaptation strategies beside its active defense mechanisms. In addition to, a combination of the stress response genes to tolerate the stressful restrictions that prevail in the desert environment in different growth seasons (Merquiol et al., 2002). It is a drought deciduous shrub where its xeromorphic adaptations include falling of leaves at the beginning of the drought season to reduce evaporation (Wickens, 1998). *R. raetam* uses the dormancy among its acclimatization strategy to resist long periods of water scarcity (< 30mm rain/year; Merquiol et al., 2002; Pnueli et al., 2002).

During dormancy *R. raetam* is not exposed to severe dehydration its relative water content is about 35% in the dormant tissue accordingly it was clustered as drought-tolerant species (Pnueli et al.,

2002). Notable decrease in overall metabolism such as apparent disappearance of cellular-proteins and suppression of photosynthesis was observed. During rainfall the dry plants recovered and rapidly gather the 'disappeared proteins, and escapes dormancy (Mittler et al., 2001; Pnueli et al., 2002).

##### *Adaptation of R. monosperma to salinity*

*R. monosperma* is a leafless leguminous shrub, nitrogen fixing (Talavera, 1999b; Zunzunegui et al., 2017). It grows on semi-stabilized and stabilized sandy dunes in marsh borders and in the coast vicinity (Muñoz Vallés et al., 2013; Zunzunegui et al., 2017). The woody perennial species like *R. monosperma* are long lived, its life span can extend to 55–80 years with main stem extends inside the soil to promote its establishment (Muñoz Vallés et al., 2005 & 2013). The response of *R. monosperma* to salinity was assessed by Zunzunegui et al. (2017), in a greenhouse experiment. Where the juvenile *R. monosperma* plants were watered with NaCl saline solution in concentrations ranges from 5.0 to 600mM. The retrieved data showed that the proline content in the assimilating stems ranged from 2  $\mu\text{mol g}^{-1}$  DM in 5mM saline to 37 $\mu\text{mol g}^{-1}$  DM in the 600mM saline, accordingly the proline content was 18-times higher owing to salinity stress (Zunzunegui et al., 2017). Sodium ions were accumulated in roots more than the assimilating stems in all NaCl concentrations, and the root and nodules dry weight increased, and the stem decreased significantly with the salinity increase (Zunzunegui et al., 2017).

*Positive interactions of R. monosperma: R. monosperma* significantly enriching the ecosystem (Jacobsen, 2000; Muñoz Vallés et al., 2011 & 2013). It facilitates the establishment and survival of other associate species. It is able to ameliorate the extreme-temperatures under the canopy during the coldest and hottest seasons, conserve the relative humidity and effectively increase the soil organic matter that enriching the plant biomass beneath the canopy with 442% (Muñoz Vallés et al., 2011 & 2013). *R. monosperma* canopies increased the enrolment of the ruderal, nitrophilous and/or weed species, which are not recorded earlier to be specific to the dune environment. It grows below the *R. monosperma* canopy compete each other due to scarcity of water, light and other resources (Muñoz Vallés et al., 2013).

*Grazing and predation of R. monosperma:* light and irregular grazing by wild rabbits has been monitored in the *R. monosperma* grow in the maritime sand dune in the Iberian Peninsula grazing on the stems (Muñoz Vallés et al., 2013). Insects (e.g., *coleopterous borers*, family Bruchidae) can damage the *R. monosperma* seeds (Muñoz Vallés et al., 2013).

*Response of R. monosperma to water stress:* It survived after subjected to the sea water invasion of the coastal dune system (Zunzunegui et al., 2017). However, it cannot survive under long-term flooding, it is shedding its leaves above-ground biomass, when exposed to this environment, fortunately, it can resprouting in the following year (Muñoz Vallés et al., 2013).

*Invasion of R. monosperma:* *R. monosperma* can easily expands in the lack of other competitor woody species (Muñoz Vallés et al., 2013), its spreading rate was up to 80m/y (Muñoz Vallés et al., 2005). In older *R. monosperma* populations that lasts more than 50 years the large individuals were dominating, and the juvenile individuals are rarer (Muñoz Vallés et al., 2013). This species showed invasive character and can modify the environment, its fast expansion was reported by Randall (1997), Zunzunegui et al. (2017) in many areas among them the sandy coasts with Mediterranean climate in California, USA and S. Australia. Fortunately, the species' salinity tolerance is a limiting factor controlling its invasion to the coastal systems. Congruent reports were addressed by Wilson & Sykes (1999), Zinnert et al. (2012), Zunzunegui et al. (2017); Zunzunegui et al. (2017) confirming that salinity is the main player affecting the spatial distribution of the plant species in the coastal environments.

*Microbes and drought stress:* Haase et al. (1996) and Selami et al. (2014), reported that *Retama* species root systems were developed to more than 20 m deep, accordingly it increased soil fertility and stability. Its root is able to fix nitrogen by the rhizobium bacteria forming the bacterial nodules developed on lateral roots (Valladares et al., 2002; Selami et al., 2014). The drought stress is a limiting factor to the species establishment and survival (Prieto et al., 2010). Under such habitat the soil microbes can play an important role in plant establishment (Prieto et al., 2010). However, *R. raetam* and *R. monosperma* are nitrogen fixing species, no detailed investigation carried out to

clarify the role of the soil microbes on the plant growth and establishment. On the other hand, the effect of indigenous *Bacillus thuringiensis*, and *Glomus intraradices* strain (isolated from the dry Mediterranean soil, Ph 7.2) with *G. intraradices* strain (isolate BEG 123; non-indigenous, and not adapted to drought), while the *R. sphaerocarpa* under drought stress the native drought-adapted fungus increased its physiological capabilities to colonize this area (Prieto et al., 2010). *R. sphaerocarpa* colonized by *G. intraradices* and *B. thuringiensis*, increased its root length by 140% and shoot length after 30 days similar to that developed after 150 days from sowing, compared to the non-inoculated *Retama* plants. While the inoculation of *R. sphaerocarpa* plants with drought-adapted bacterium increased the root growth by 201%, with higher relative water uptake (Prieto et al., 2010). The co-inoculation of indigenous microorganisms reduced 42% from the water requests to produce one milligram of shoot biomass. Rhizosphere bacterium, singly or in association with an arbuscular mycorrhizal (AM) fungus are efficient in increasing plant water uptake, under drought environments, and can be used as alternative strategy for the best colonization of *R. sphaerocarpa* under environmental limitations (Prieto et al., 2010).

#### *Symbiosis of endophytic bacteria to Retama species*

The symbiotic endophytes of genus *Retama* were grouped under four bacterial genera (*Agrobacterium*, *Bradyrhizobium*, *Rhizobium* and *Sinorhizobium*) and genus *Glomus* from fungi (Valladares et al., 2002; Caravaca et al., 2003; Mahdhi et al., 2008; Farida et al., 2009; Muñoz Vallés et al., 2013).

*Bradyrhizobium* is the principal genus of the nitrogen-fixing endophytic bacteria to *Retama* species (Rodríguez-Echeverría et al., 2003; Farida et al., 2009; Guerrouj et al., 2013; Lamin et al., 2019). Eleven haplotypes belonged to the genus *Bradyrhizobium* were identified from *Retama sphaerocarpa* and *R. raetam* inhabiting seven sites from different ecological-climatic areas in NE Algeria. One rhizobial haplotype from the identified populations was common across these sites, the rest haplotypes showed significant biogeographical difference (Farida et al., 2009).

Mahdhi et al. (2008) and Lamin et al. (2019), reported that *Sinorhizobium* sp., *Sinorhizobium*

*kostiense* and *S. meliloti* (genus *Bradyrhizobium*) were isolated from *Retama raetam* growing in Tunisian arid area (Lamin et al., 2019). Twelve bacterial isolates were isolated from *R. raetam* root-nodules, the species growing in Tunisian arid regions. These isolates were grouped under *Agrobacterium*, *Rhizobium* and *Sinorhizobium* (Mahdhi et al., 2008). Zakhia et al. (2006) identified five genera, *Bo sea*, *Microbacterium*, *Ochrobactrum*, *Paracraurococcus*, and *Starkeya* from the *R. raetam* inhabited the infra-arid areas in Tunisia, all these strains do not succeed to nodulate their host of origin (Mahdhi et al., 2008).

*Bradyrhizobium retamae* was isolated from *R. monosperma*, inhabiting a sub-humid region in N. Morocco (Guerrouj et al., 2013; Lamin et al., 2019), this species can nodulate also *R. sphaerocarpa* (Lamin et al., 2019). Fast growing isolates (*Sinorhizobium* and *Rhizobium* genera) were isolated from the Spanish *R. sphaerocarpa* similar, isolates were isolated from the bacterial nodules in *R. raetam* growing in arid zone in Tunisia (Mahdhi et al., 2008; Farida et al., 2009).

The inoculated *R. sphaerocarpa* seedlings with rhizobia isolated from *Retama* sp. were growing in Spain under two different fertilization levels. The inoculated seedlings growing in good fertilization showed higher photosynthetic rates and taller individuals than the non-inoculated seedling at same fertilization level (Valladares et al., 2002). The heavy metal tolerant *Ensifer aridi* were isolated from *R. monosperma* nodule growing in lead-rich mine soils and improve the plant resilience to this environment (Lamin et al., 2019).

#### *Genetic diversity in R. raetam related to environmental conditions*

Abdellaoui et al. (2014) studied the genetic diversity among and within the *R. raetam* populations, the C3 stem assimilating desert legume from different habitats in S. Tunisia using RAPD molecular marker. The results revealed that the main variations (68%) were noticed within populations and there were significant variations among populations ( $\Phi_{PT} = 0.316$ ,  $P < 0.001$ ). The gene differentiation coefficient was found in range from 0.490 - 4.609 and 1.42 between pair-wise and among populations; respectively. This study highlights the adaptive strategy of this important species in S. Tunisia (Abdellaoui et al., 2014).

#### *Phytochemistry of Retama species*

The phytochemistry of the *Retama* species have been subjected to detailed analysis, several chemical constituents were reported among them, the alkaloids (Abdel Halim et al., 1997; Edziri et al., 2010), essential oils (Maghrani et al., 2005a; Merghoub et al., 2009; El Hamdani & Fdil, 2015), flavonoids (Morales et al., 1971; El-Shazly et al., 1996; Kassem et al., 2000; Maghrani et al., 2005b; Edziri et al., 2010; El Hamdani & Fdil, 2015), and quinolizidine alkaloids (Eddouks et al., 2007; Cheriti et al., 2009; El Hamdani & Fdil, 2015).

**Flavonoids:** The flavonoids identified in *R. raetam* seeds were apigenin, chrysoeriol 7-*O*-glucoside, daidzein 7,4'-dimethyl ether, daidzein, kaempferol, kaempferol-7-*O*-glucoside, naringenin, and quercetin. In addition to orientin detected in leaves (Kassem et al., 2000; Edziri et al., 2010). Seven flavonoids were identified from the *R. raetam* and *R. sphaerocarpa* in Algeria as: four isoflavones (daidzein, daidzin, genistein and genistin), two flavones (apigenin and luteolin), and the rutin flavonol (Hammouche-Mokrane et al., 2017). Touati et al. (2015) and Leon-Gonzalez et al. (2018), reported several isoflavones identified from *R. monosperma*, *R. raetam* and *R. sphaerocarpa*, as genistein, genistin, daidzin, and daidzein. In addition to biochanin A, puerarin and 6'-methoxypseudobaptigenin isoflavones were also reported (Abdalla & Saleh, 1983; López-Lázaro et al., 1998; Djeddi et al., 2013; Leon-Gonzalez et al., 2018), Derrone and 5 hydroxyderrone, are among the identified isoflavones from *R. raetam* stem (Xu et al., 2015; León-González et al., 2018), While Quinic acid (phenolic compound) is among the main constituents of seeds and stems of *R. sphaerocarpa* (Touati et al., 2017; León-González et al., 2018).

*R. raetam* contains phenolic compounds (represented as Isoflavonoids) in considerable amount (23.9mg GAE·g<sup>-1</sup>) and 27.2µg·g<sup>-1</sup> FW carotenoids (Saada et al., 2018). Flavonoids were varying with the plant organ, seeds containing such as quercetin, daidzein, naringenin, vicenin-2, apigenin, kaempferol, and kaempferol-7-*O*-glucoside, while leaves containing chrysoeriol 7-*O*-glucoside, daidzein 7,4'-dimethyl ether, and orientin (Xu et al., 2015). Aerial parts containing 5,4'-dihydroxy-(3",4"-dihydro-3",4"-dihydroxy)-2",2"-dimethylpyrano-(5",6":7,8)-flavone and luteolin 4'-*O*-neohesperidoside (Saada et al., 2018).

**Essential oil:** the analysis of the essential oil from the *R. raetam* flowers from Libya revealed that the 85.60% of the total oil content comprises Oxygenated monoterpenes, monoterpenes, oxygenated sesquiterpenes and other compounds, 62.0%, 10.6%, 1.6%, and 11.0%; respectively. The oxygenated monoterpene fraction containing 50.9%  $\beta$ -linalool comprising the main constituent followed by 6.6% 2-decen-1-ol, 2.3% cis-linalool oxide, 1.6% geraniol format and 0.6% ethyl linalool. One compound from the oxygenated sesquiterpenoid represented 1.6% (nerolidol acetate), and three monoterpenes namely limonene, terpinolene, and cis- $\beta$ -ocimene with concentrations 7.4%, 1.7% and 1.5%; respectively. While *R. raetam* oil is dominated by 50.9%  $\beta$ -linalool, 7.4% limonene 6.6% 2-decen-1-ol and 3% oleic acid (Awen et al., 2011). The main essential oil extracted from *R. raetam* flowers is nonanal alcohol; in addition to octanal, dodecanal and undecanal aldehydes (Touati et al., 2015; León-González et al., 2018). Edziri et al. (2010), León-González et al. (2018) reported that the flowers of *R. raetam* are rich of other essential oils ( $\alpha$ -humulene,  $\beta$ -linalool and nonanal). Touati et al. (2015), León-González et al. (2018) reported the identification of triterpene  $\beta$ -amyrin (0.06%), and steroids from *R. raetam* stem (Belayachi et al., 2014). While the stems of *R. monosperma*, *R. sphaerocarpa* and *R. raetam* showed the presence of pinitol with percentages 2.3%, 1.9% and 1.8%; respectively (González-Mauraza et al., 2016; León-González et al., 2018). The norterpenoids is the pleasant aroma present in a significant amount in *R. monosperma* flower (León-González et al., 2018).

**Alkaloids:** in Algeria, ten alkaloids were identified from the *R. etama r aetam* and *R. sphaerocarpa* six of them were tetracyclic quinolizidine ( $\alpha$ -isolupanine, 5,6-dehidrolupanine, dehydroretamine, lupanine, retamine and sparteine); two bipiperidyl (ammodendrine and N-formyl-ammodendrine) and two tricyclic quinolizidine (anagryne and N-methylcytisine) (Hammouche-Mokrane et al., 2017). El-Shazly et al. (1996) identified from the quinolizidine alkaloids 28 compounds and one dipiperidine alkaloids (ammodendrine) in all plant parts of *R. monosperma*, *R. sphaerocarpa* and *R. raetam*, but they were present in lower concentrations in *R. raetam*. Also, Quinolizidine alkaloids and 31 bipiperidine alkaloids were identified from different parts of the same three *Retama* species

(El-Shazly et al., 1996; León-González et al., 2018).

Alkaloids were more analogous in these three species of Mediterranean distribution than its profile between the organs of the same species. The major alkaloids detected in stem were retamine and sparteine; while fruits and flowers were dominated by cytisine, lupanine, N-methylcytisine and retamine (El-Shazly et al., 1996); and seeds are the storage organ of cytisine alkaloid (El-Shazly et al., 1996; Sadik et al., 2020). Sadik et al. (2020), reported the presence of three alkaloids in *R. monosperma* seeds in descending percentages as 77.60% cytisine, 13% N-methylcytisine and 9.40% dehydro-cytisine. While the stem contains ten alkaloids: Ammodendrine, Anagryne, Cytisine, Dehydrosparteine, 5,6-Dehydrolupanine, 11,12-Dehydrolupanine, Isolupanine, N-methylcytisine, 17-Oxosparteine and Sparteine.

**Lipid and fatty acids:** Chemical identification analysis of *R. monosperma* hexane extract indicated that  $\alpha$ -linolenic acid, campesterol, stigmasterol and sitosterol were the major components (Belayachi et al., 2014). The lipids component in seeds and branches/leaves of *R. monosperma* from western Morocco were 5% and 0.3 %; respectively (El Hamdani & Fdil, 2015). Fatty acids of *R. monosperma* seeds and stems revealed the identification of five unsaturated fatty acids and eleven saturated (El-Hamdani & Fdil, 2015; León-González et al., 2018). The fatty acids in seeds and branches/leaves of *R. monosperma* from western Morocco were dominated by palmitic acid, followed by stearic acid. Oleic acid is the dominant uni-unsaturated fatty acid in seeds. The major polyunsaturated acids in seeds and branches/leaves were linoleic and linolenic acid, the earlier concentrated in seeds and the later was major in branches/leaves (El Hamdani & Fdil, 2015).

Touati et al. (2015) and León-González et al. (2018), reported the identification of 14 fatty acids from *R. sphaerocarpa* stems and seeds where the unsaturated (14%) were dominant over the saturated fatty acids (2.3% w/w). From stems and seeds of *R. monosperma* and *R. sphaerocarpa*, León-González et al. (2018) and Touati et al. (2015) identified campesterol, b-sitosterol and stigmasterol. At the vegetative stage *R. raetam* has high percentage of polyunsaturated fatty

acids (66.49%), and notable amount of vitamin C (645.6mg·100g<sup>-1</sup> FW) and the proline content (25.4μmol·g<sup>-1</sup> DW) (Saada et al., 2018).

*Mineral composition:* the mineral analysis of *R. monosperma* seeds and branches/leaves from western Morocco were revealed that the highest mineral constituents were Al, Ca, Fe, K, Mg, Na, P and Zn, the concentrations are mainly dependent and affected by soil nature and rock composition (El Hamdani & Fdil, 2015). *R. sphaerocarpa* shrub is rich in its mineral composition, it contains considerable amounts of Ca (8.62g.kg<sup>-1</sup>), Mg (3.11g.kg<sup>-1</sup>), Na (2.21g.kg<sup>-1</sup>) and (P 2.24g.kg<sup>-1</sup>); while zinc and Mn in leaves were 0.04g.kg<sup>-1</sup> and 0.03g.kg<sup>-1</sup>, respectively (Zamberlin et al., 2012; Abdenour et al., 2020). Based on their chemical composition, the leaves of *R. sphaerocarpa* showed high fiber (NDF= 598 g.kg<sup>-1</sup> DM and ADF= 432g.kg<sup>-1</sup> DM), lignin contents (ADL =178g.kg<sup>-1</sup> DM) and the crude protein (CP=137g.kg<sup>-1</sup>) (Abdenour et al., 2020).

*Polysaccharides:* two galactomannans polysaccharides were identified from *R. raetam* seeds from Libyan (Ishurd et al., 2004; León-González et al., 2018).

#### *Economic uses of Retama*

*Retama raetam* is a leguminous shrub, that plays a significant role in arid lands, particularly in soil protection against overheating and direct irradiance (Moro et al., 1997; López-Pintor et al., 2000; Barakat et al., 2013). It stabilizes soil against water and wind erosions, and offers a valuable fodder for sheep, camels and goats, this species possesses significant nutritional value for the livestock (Laudadio, 2009; Barakat et al., 2013; Al-Sharari et al., 2020). It increases soil fertility through its nitrogen fixation and enhances the soil fertility (Dart, 1998; Barakat et al., 2013); Its wood is a fuel source (Cheriti et al., 2009; Barakat et al., 2013). *Retama* tree was used by local inhabitants in Sahara as adsorbent used to remove copper ions from aqueous polluted solutions (Cheriti et al., 2009). It also applied for remediation of heavy metal contaminated soil and mining sites (Lamin et al., 2019; Al-Sharari et al., 2020). From an environmental point of view, *Retama* is one of the most important species used for rehabilitation of the degraded soil and soil under desertification stress (Caravaca et al., 2003; Al-Sharari et al., 2020).

#### *Ethnobotanical uses of the Retama spp.*

*Retama raetam* is used in folk medicine in all the countries of its geographic range (Table 2 & Leon-Gonzalez et al., 2018; Al-Sharari et al., 2020). Leave powder used for wound healing or antiseptic for wounds or for treatment of skin irritation (Awen et al., 2011; Al-Sharari et al., 2020). *Retama* sp. was traditionally used to cure renal disorders (González-Tejero et al., 2008; Al-Sharari et al., 2020); it possesses significant diuretic activity (Kassem et al., 2000; Al-Sharari et al., 2020). Moreover, *R. raetam* is used for treatment of many diseases, among them the jaundice, sore throat, inflammation, joint pains, fever and microbial infections (Edziri et al., 2012; Djeddi et al., 2013; Al-Sharari et al., 2020). In Tunisia, it is used as medication for snake bites (El Hamrouni, 2001; Al-Sharari et al., 2020) and in some renal diseases (Edziri et al., 2010; Al-Sharari et al., 2020). Bedouins used *Retama* in treatment of arthralgia, backache, infertility and for inducing abortions (Bailey & Dannin, 1981; Al-Sharari et al., 2020). In Morocco and Saudi Arabia, it used in hypertension and diabetes treatment (Nur-e-Alam et al., 2019; Al-Sharari et al., 2020). Bedouins still used *R. raetam* to prepare slow combustion coals (Schmid et al., 2006). *R. raetam* roots are treated diarrhea, while leaves are applied to help eye troubles and aching joints back pain (Said et al., 2002; Barakat et al., 2013).

In Tunisia folk medicine, *R. raetam* is used for treatment of several diseases among them: renal diseases due to its significant diuretic activity (Caceres et al., 1987; Edziri et al., 2010), hypertension (Archer & Pyke, 1991; Izhaki & Neeman, 1997; Edziri et al., 2010). The flower oil showed antibacterial, antifungal for yeasts and antioxidant with IC50 = 0.800mg/mL (Edziri et al., 2010). Edziri et al. (2010) attributed the antioxidant activity of this flower-oil to the presence of monoterpenes in high concentrations, this natural oil is a potential preservative in pharmaceutical and/or food products.

#### *Pharmacological activities of Retama spp.*

*Retama raetam:* the *R. raetam* extract of the vegetative parts showed remarkable antioxidant activity (Saada et al., 2018; Al-Sharari et al., 2020); anti-inflammatory (González-Mauraza et al., 2016; Al-Sharari et al., 2020); hypoglycemic (Maghrani et al., 2005a; Hayet et al., 2008; Al-Sharari et al., 2020) and antibacterial activities (Hammouche-Mokrane et al., 2017; Al-Sharari et al., 2020); in

**TABLE 2. Ethnopharmacological applications of *Retama* spp. species, country, used plant part in each case, preparation, and route of administration**

Country/ province	Used plant part	Preparation	Traditional use	Route of administration	References
<b><i>R. dasycarpa</i> Coss.</b>					
Morocco (Atlas Mountains)	Seeds		Nephrological disease, urological	Oral	Teixidor-Toneu et al. (2016)
<b><i>R. monosperma</i> (L.) Boiss</b>					
Algeria	Cladodes	Decoction	Hydrophobia (rabies) prevention	Oral	Helmstädter (2016)
Morocco	Cladodes	Decoction	Purgative and Vermifuge	Rectal washings	Bellakhdar (1997)
Morocco	Cladodes	Grounded and blending with honey	Emetic	Oral	Bellakhdar (1997)
<b><i>R. raetam</i> (Forsk.) Webb.</b>					
Algeriac, (M'Sila)	Cladodes	Decoction	Eczema	External use	Boudjelal et al. (2013)
Algeria (Ouanougha)	Cladodes	NS	Skin disease recovery, diarrhea, inflamed eyes, fever		Rebbas et al. (2012)
Algeria (Ouargla)	Cladodes	NS	Rheumatism, Skin wounds, Scorpion sting	NS	Ould El Hadj et al. (2003)
Algeria (Ouargla)	Seeds, fruits	Decoction, Infusion	Diabetes	Oral	Telli et al. (2016)
Algeria	Cladodes	Grounded and blending with olive oil	Back pain, skin wounds	External use	Rebbas et al. (2012)
Algeria	Cladodes	Infusion	Treat stomachache	Oral	Rebbas et al. (2012)
Algeria	NS	NS	To treat rabies	NS	Louaar et al. (2005)
Israel	Cladodes	Decoction	Joint aches, skin bruise and back pain	Bath	El-Mokasabi (2014)
Jordan	Cladodes	Decoction	Burns and fractures	Poultice	Hudaib et al. (2008)
Lebanon	Cladodes	Decoction	Joint aches	Bath	El-Beyrouthy et al. (2008)
Libya (Al-Jabal Al-Akhder)	NS	NS	Sinusitis, diabetes	NS	Said et al. (2002)
Middle East	Flowers, Cladodes	Decoction	Women infertility, syphilis	External use	Yaniv & Dudai (2014)
Morocco (Marrakech)	Cladodes	Decoction	Scabies and Antipruritic	Liniments	Bellakhdar et al. (1991); Bellakhdar (1997)
Morocco (Sahara)	Roots	Decoction	Abortive	Vaginal washings	Bellakhdar (1997)
Morocco (Sahara)	Roots	NS	Diphtheria	NS	Mouhajir (2002)
Morocco (Taounate, Tata)	Flowers, Cladodes	Decoction	Skin disease	External use	Bellakhdar et al. (1991); El-Hilaly et al. (2003); Abouri et al. (2012)
Morocco (Tata)	Cladodes	Cataplasm	Wounds healing, scorpion bite	External use	Abouri et al. (2012)
Morocco (Tata)	Cladodes	Infusion	Rheumatism	Oral	Abouri et al. (2012)

TABLE 1. Cont.

Country/ province	Used plant part	Preparation	Traditional use	Route of administration	References
Morocco (Tissint)	Cladodes	Powdered	Local wound treatment antiseptic and also sedative, Circumcision healing Skin ulcers and wounds, vulnerary	Cataplasm	Bellakhdar et al. (1991); Bellakhdar (1997)
Morocco	Cladodes	Grounded and blending with honey	Emetic	Oral	Bellakhdar (1997)
Morocco	Cladodes	Decoction	Purgative and Vermifuge	Rectal washings	Bellakhdar et al. (1991); Bellakhdar (1997)
Morocco	Flowers, Cladodes	Infusion	Abortive	Oral	Abouri et al. (2012); Bellakhdar (1997)
Palestine	Seeds, cladodes	NS	Treat stomachache, analgesic	Oral	Ali-Shtayeh et al. (1998)
Palestine	Seeds, cladodes	NS	Anti-inflammatory, sore throat treatment and treat inflamed eyes, paralysis and infertility, antirheumatic	Poultice	Ali-Shtayeh et al. (1998)
Tunisia	Cladodes	NS	Scabies	Poultice	Viegi & Ghedira (2014)
Yemen	Cladodes	Infusion	Jaundice, hepatitis	Internal use	Hehmeyer & Schönig (2012)
<b><i>R. sphaerocarpa</i> Boiss</b>					
Morocco (Errachidia)	Root	Decoction	Diabetes	Internal use	Tahraoui et al. (2007)
Morocco (Sahara)	Roots	NS	Diphtheria	NS	Mouhajir (2002)
Morocco	Cladodes	Decoction	Purgative and Vermifuge	Rectal washings	Bellakhdar (1997)
Morocco	Cladodes	Grounded and blending with honey	Emetic	Oral	Bellakhdar (1997)
Spain	Flowers, Cladodes	Decoction	Rheumatism, Diabetes, warts Healing,	External and oral use	Benítez Cruz (2007)
Spain	Flowers	Crushed in water	Skin wound healing	Poultice	Benítez Cruz (2007)
Spain	Cladodes	No preparation	Luxation	Topic	Benítez et al. (2010)
Spain	Flowers	Cataplasm	Pain, contusion	Topic	Benítez et al. (2010)
Spain	Cladodes	Infusion, Decoction	Fever	Oral	Benítez Cruz (2007); Benitez et al. (2010)
Spain	Flowers	Infusion	Liver disease	Oral	Benítez Cruz (2007); Benitez et al. (2010)
Spain	Fruit	Fresh ingested	Diarrhea	Oral	Benitez et al. (2010)
Spain	Cladodes	Crushed by vinegar, salt or ash	Joint aches	Poultice	Benítez Cruz (2007)

NS: not specified.

addition to its pharmacological effect in nervous system (Al-Tubuly et al., 2011; Al-Sharari et al., 2020). Essential oils from flowers indicated antifungal, antibacterial (Edziri et al., 2010; Al-Sharari et al., 2020), antimicrobial and antiseptic activities (Awen et al., 2011; Al-Sharari et al., 2020). Flavonoids  $\alpha$ -glucosidase showed inhibitory effect (Ghani et al., 2019; Al-Sharari et al., 2020); antidiabetic activities (Algandaby et al., 2010; Al-Sharari et al., 2020) and minimal nephrotoxic toxicity (Algandaby, 2015; Al-Sharari et al., 2020).

Retamine and sparteine are among the 30 alkaloids identified from *R. raetam* stem; both may be lethal to experimental animals if given in large doses by oral or intravenous mode. Retamine and sparteine are block autonomic ganglia and cardiotoxic, while the sparteine causing paralyzing respiratory muscles leading to respiratory failure (Schmid et al., 2006). The isoflavones genistein, 6-hydroxygenistein, biochanin A, 3'-O-methylrobofol, pratensein, luteolin, the flavones 6-hydroxyapigenin and the flavonol kaempferol, as well as the p-coumaric acid (phenolic compound) isolated from *R. raetam* have reduce significantly the pain at dose of 1 mg/kg due to its analgesic activity. 3'-O-methylrobofol and biochanin A are the most active compounds 86.19% and 75.23%; respectively (Djeddi et al., 2013).

The aqueous extracts of all organs of the *R. raetam* were showed antioxidant activities, as it utilized very low free radical capturing activity if compared to the butylated hydroxytoluene (BHT) and possess lower hydrogen peroxide blocking activity compared to the gallic acid (Djeddi et al., 2013). Also, some of the isoflavonoids separated from the methanol extract of *R. raetam* showed analgesic activity, is almost equivalent to aspirin in the case of 86.9% from 3'-O-methylrobofol (Djeddi et al., 2013).

Abed & Benmrabet (1981) and Maghrani et al. (2005b), reported that the active principles such as saponins, flavonoids, and organic acids in the aqueous *R. raetam* extract causing a significant diuretic effect (Maghrani et al., 2005b), it observed as increase on diuresis from the second to the fourth hours if administered intravenous as 5 mg/kg/h in normal rats. Such compounds are acting as angiotensin converting enzyme (ACE) inhibitors that act separately or synergistically to generate this diuretic effect (Actis-Goretta et al.,

2003; Maghrani et al., 2005b).

*R. monosperma* and others: *R. monosperma* is a potential source of anticancer compounds, it exhibits a significant cytotoxic effect on two cervical cancer cell lines (Merghoub et al., 2009; Muñoz Vallés et al., 2013). *R. monosperma* tissues accumulate cytotoxic alkaloids in stems and fruits (Salatino & Gottlieb, 1980; El-Shazly et al., 1996; Muñoz Vallés et al., 2013). Among these alkaloids the anagryne and ammodendrine that cause congenital diseases in pregnant animals (Keeler, 1969; Muñoz Vallés et al., 2013). The ethyl acetate extract of *R. monosperma* seeds is a potential natural antioxidant source, it showed 197.95mg ascorbic acid equivalent (AAE)/g dry extract as total antioxidant capacity (TAA). While the butanol extract of the flower showed the lowest value it was 26mg (AAE)/g dry extract (Belmokhtar & Harch, 2014). The antioxidant activity of *R. monosperma* was attributed to the antioxidant activity of the genistein isoflavone which comprises its major flavonoid (González-Mauraza et al., 2013; Belmokhtar & Harch, 2014). Genistein is an efficient antioxidant as well as it increases the activity of several antioxidant enzymes among them: catalase, glutathione reductase, glutathione peroxidase, superoxide and dismutase (Qiuyin & Huachen, 1996; Trieu et al., 1999; Belmokhtar & Harch, 2014). The antioxidant activity of genistein isoflavone has potential applications for cancer prevention, and treatment of cardiovascular diseases and other human diseases (Belmokhtar & Harch, 2014). In Tunisia, the *R. raetam* containing lower content of the total phenols and flavonoids if compared to *R. monosperma* (Hayet et al., 2008; Belmokhtar & Harch, 2014). On the other hands, the hexane extract of *R. monosperma*, showed a significant cytotoxic activity against the human T-lymphocyte cells (Jurkat cells), due to its bioactive constituent namely: campesterol,  $\alpha$ -linolenic acid, sitosterol and stigmaterol (Belayachi et al., 2014).

Quinolizidine alkaloids were isolated from *R. raetam*, *R. sphaerocarpa* and *R. monosperma* (El-Shazly et al., 1996). The earlier reports outlined several toxicological and pharmacological activities of these alkaloids among them antipyretic, antiarrhythmic, depressant, diuretic, hallucinogenic, hypoglycemic, hypotensive, oxytocic, respiratory stimulant and uterotonic properties (Kingham & Balandrin, 1984; Wink, 1993; El-Shazly et al., 1996). Quinolizidine

alkaloids showed an important provision for future use to protect crops from microbial infection and herbivores feeding (El-Shazly et al., 1996). Touati et al. 1996; Fdil et al., 2012; Belmokhtar & Harch, 2014; Belmokhtar & Harch, 2014), claimed that *R. monosperma* contained high proportions of alkaloids. The pharmacological investigations showed that these compounds showed anti-inflammatory activity (González-Mauraza et al., 2013), anti-leukemic activity (Belayachi et al., 2014) and in vitro inhibition of the cervical cancer cell lines (Merghoub et al., 2011; Benbacher et al., 2012; Belmokhtar & Harch, 2014). The antidiabetic activity of *R. monosperma* may be attributed to its fatty acids composition as palmitic acid, oleic acid, linolenic acid and linoleic, the pharmaceutical indication of the latter is effective as antileukemic (El Hamdani & Fdil, 2015).

#### *Antimicrobial activity of the Retama spp.*

The ethyl acetate fraction of *R. raetam* is efficient antibacterial (MICs of 128–256mg ml<sup>-1</sup>) for the *Staphylococcus aureus* the methicillin-resistant Gram-positive bacteria (Hayet et al., 2007; Awen et al., 2011). As well as for the Gram-negative bacteria with inhibition zone 8-11mm and 11-14mm for the *Staphylococcus aureus*, *Escherichia coli* and *Bacillus subtilis*; respectively (Ahmed et al., 2016). While, the methanol-water (50:50) fraction of *R. sphaerocarpa* stems and the polyphenol-rich fraction revealed a significant antibacterial activity against *Pseudomonas aeruginosa* and *Staphylococcus aureus* (Touati et al., 2017; León-González et al., 2018). The essential oil of *R. raetam* is a potential antimicrobial mediator for drugs and traditional plant-based preparations (Awen et al., 2011) The derrone and licoflavone C flavonoids isolated from *R. raetam* by ethyl acetate showed antibacterial activity against *Pseudomonas aeruginosa* and *Escherichia coli* (Edziri et al., 2012; León-González et al., 2018). The aqueous extract of *Retama raetam* and *R. sphaerocarpa* showed the antibacterial activities against the methicillin-resistant *Staphylococcus aureus* (MRSA) and *S. aureus*, where the aqueous extract of the later was more active in MIC 125µg/mL than that of *R. raetam* and bactericidal to these *Staphylococcus* strains (Hammouche-Mokrane et al., 2017). *Retama* flavonoids induce antimicrobial activity where the extract of the vegetative parts showed considerable antibacterial capacities versus human pathogens namely, *Bacillus cereus* and *Vibrio vulnificus* (Saada et al., 2018).

#### *Antifungal activity of the Retama spp.*

The seed alkaloids of *R. monosperma* have been as anticorrosive extract (Dart, 1998; El Hamdani & Fdil, 2015). While extracts of its aerial parts (stems and leaves) were found to be efficient antifungal at (500µg/mL) concentration against *Candida albicans* (16.66- and 18.66-mm inhibition zone; respectively) and *Candida tropicalis* (14.33- and 20.66-mm inhibition zone; respectively). The leaf alkaloids were more effective on *Candida tropicalis* than stems alkaloids. The antifungal of the leaves and stems alkaloids at the same concentration extends to cover *Aspergillus niger* (8.66- and 9.33-mm inhibition zone; respectively). In contrast, *A. niger* and *Candida* species were resistant to the seed alkaloids of *R. monosperma* (El Hamdani & Fdil, 2015). The antifungal activity of the stems and leaves alkaloids may be attributed to the presence of anagryne, ammodendrine and sparteine. On the other hand, the high concentration of cytisine and its related derivatives in the seed alkaloid extracts inactivate its antifungal activity (El Hamdani & Fdil, 2015). The flavonoids (derrone and licoflavone C) isolated from *R. raetam* by ethyl acetate showed significant antifungal activity against *Candida* species (Edziri et al., 2012; León-González et al., 2018).

#### Conclusion

This review gathered a huge information in different disciplines about the four multipurpose Mediterranean *Retama* species. However, this genus still needs further effort concerning its genetic diversity, genetic response to environmental stresses in Arid-Mediterranean region and chromosome number as well as inter and intra-specific karyotyping this interm of research. In addition to its urgent need for a regional conservation plan.

*Acknowledgements:* The authors acknowledge the staff of Cairo University Herbarium (CAI), for the academic facilities given to the authors during this work.

*Conflict of interest:* None.

*Authorship contribution:* Reham A. Youssef: Data collection and prepared the manuscript. Wafaa M. Amer: supervised the whole work, and refined the manuscript. Omran N. Ghaly: helped in manuscript preparation and the field work.

Azza B. Hamed: Helped in writing and revision of the manuscript and the section of references.

*Ethics approval:* Not applicable.

### References

- Abdalla, M.F., Saleh, N.A.M. (1983) Flavonoids of *Retama raetam*. *Journal of Natural Products*, **46**(5), 755-756.
- Abdel Halim, O.B., Abdel Fattah, H., Halim, A.F., Murakoshi, I. (1997) Comparative chemical and biological studies of the alkaloidal content of *Lygos* species and varieties growing in Egypt. *Acta Pharmaceutica Hungarica*, **67**(6), 241-247.
- Abdellaoui, R., Yahyaoui, F., Neffati, M. (2014) Population Structure and Genetic Diversity of a Medicinal Plant Species. *Pakistan Journal of Biological Sciences*, **17**(2), 182-189.
- Abdenour, K., Mounia, M.L., Mohamed, D., Mourad, T., Abdallah, C., Charaf, D. (2020) Nutritional characteristics and seed germination improvement of the forage shrub *Retama sphaerocarpa* (L.) boiss. *Journal of Agricultural Science*, **69**(3-4), 53-60.
- Abed, L., Benmrabet, K. (1981) Intérêt de l'apport en potassium et sodium des infusions de plantes médicinales. *Plantes Médicinales et Phytothérapie*, **15**(1), 92-98.
- Abouri, M., El Mousadik, A., Msanda, F., Boubaker, H., Saadi, B., Cherifi, K. (2012) An ethnobotanical survey of medicinal plants used in the Tata Province, Morocco. *International Journal of Medicinal Plants Research*, **1**(7), 99-123.
- Actis-Goretta, L., Ottaviani, J.I., Keen, C.L., Fraga, C.G. (2003) Inhibition of angiotensin converting enzyme (ACE) activity by flavan-3-ols and procyanidins. *FEBS Letters*, **555**(3), 597-600.
- Ahmed, M.S., Abdel-Hameed, U.K., Saeed, M.N. (2016) Morphological characterization and biological activity of some ethno-medicinal plants of Sinai-Egypt. *The Egyptian Journal of Experimental Biology (Botany)*, **12**(2), 133-143.
- Algandaby, M.M. (2015) Assessment of acute and subacute toxic effects of the Saudi folk herb *Retama raetam* in rats. *Journal of the Chinese Medical Association*, **78**(12), 691-701.
- Algandaby, M., Alghamdi, H.A., Ashour, O.M., Abdel-Naim, A.B., Ghareib, S.A., Abdel-Sattar, E.A., Hajar, A.S. (2010) Mechanisms of the antihyperglycemic activity of *Retama raetam* in streptozotocin-induced diabetic rats. *Food and Chemical Toxicology*, **48**(8-9), 2448-2453.
- Ali-Shtayeh, M.S., Yaghmour, R.M., Faidi, Y.R., Salem, K., Al-Nuri, M.A. (1998) Antimicrobial activity of 20 plants used in folkloric medicine in the Palestinian area. *Journal of Ethnopharmacology*, **60**(3), 265-271.
- Al-Sharari, N., Bakhawain, A.S., Elfeel, A.A. (2020) Profiling and importance of underutilized neglected species of hyper arid climate of Saudi Arabia (*Retama raetam* - Retem): A review. *Life Science Journal*, **17**(7), 35-42.
- Al-Tubuly, R., Auzi, A.A., Al-Etri-Endi, A.A., Nahar, L., Sarker, S.D. (2011) Effects of *Retama raetam* (Forssk.) Webb & Berthel. (Fabaceae) on the central nervous system in experimental animals. *Archives of Biological Sciences*, **63**(4), 1015-1021.
- Amer, W.M., Elshayeb, N.F., Hegazy, A.K., Abbas, M.S., Soliman, A.S. (2021) Long-term species diversity and climate change: An intimate relationship over the last ten decades: case study in Egypt. In: "Handbook of Climate Change Management", W. Leal Filho et al. (Eds.), pp. 1-24. Springer Nature, Switzerland.
- Archer, S., Pyke, D.A. (1991) Plant animal interactions affecting establishment and persistence on revegetated rangelands. *Journal of Range Management*, **44**(6), 558-565.
- Awen, B.Z.S., Unnithan, C.R., Ravi, S., Kermagy, A., Sasikumar, J.M., Khrbash, A.S., Ekreem, W.L. (2011) Essential oils of *Retama raetam* from Libya: Chemical composition and antimicrobial activity. *Natural product research*, **25**(9), 927-933.
- Bailey, C., Danin, A. (1981) Bedouin plant utilization in Sinai and the Negev. *Economic Botany*, **35**(2), 145-162.
- Barakat, N.A., Laudadio, V., Cazzato, E., Tufarelli, V. (2013) Potential contribution of *Retama raetam* (Forssk.) Webb & Berthel as a forage shrub in Sinai, Egypt. *Arid Land Research and Management*, **27**(3), 257-271.

- Barker-Webb, M.M.P., Berthelot, S. (1836) "*Histoire Naturelle des Iles Canaries*". Saint-Andre-Des-Art, Paris.
- Bedair, H., Shaltout, K., Al-Sodany, Y., Halmy, M. (2024) Phytosociological Survey of Mediterranean Endemic Plants in Egypt. *Egyptian Journal of Botany*, 64(1), 359-374. doi: 10.21608/ejbo.2023.230811.2458.
- Belayachi, L., Aceves-Luquero, C., Merghoub, N., Bakri, Y., Fernández de Mattos, S., Amzazi, S., Villalonga, P. (2014) *Retama monosperma* n-hexane extract induces cell cycle arrest and extrinsic pathway-dependent apoptosis in Jurkat cells. *BMC Complementary and Alternative Medicine*, 14(38), 1-12.
- Bellakhdar, J. (1997) Contribution à l'étude de la pharmacopée traditionnelle au Maroc: la situation actuelle, les produits, les sources du savoir. Enquête ethnopharmacologique de terrain réalisée de 1969 à 1992. *Ph.D. Dissertation*, University of Metz, France.
- Bellakhdar, J., Claisse, R., Fleurentin, J., Younos, C. (1991) Repertory of standard herbal drugs in the Moroccan pharmacopoea. *Journal of Ethnopharmacology*, 35(2), 123-143.
- Belmokhtar, Z., Harche, M.K. (2014) *In vitro* antioxidant activity of *Retama monosperma* (L.) Boiss. *Natural Product Research*, 28(24), 2324-2329.
- Benbacer, L., Merghoub, N., El Btaouri, H., Gmouh, S., Attaleb, M., et al. (2012) Antiproliferative effect and induction of apoptosis by *Inula viscosa* L. and *Retama monosperma* L. extracts in human cervical cancer cells. In: "*Topics on Cervical Cancer with an Advocacy for Prevention*", R. Rajamanickam (Eds.), pp. 267-283. InTech, Croatia.
- Benhouhou, S. (2005) "*A guide to Medicinal Plants in North Africa*". IUCN Centre for Mediterranean Cooperation, Malaga, Spain.
- Benítez Cruz, G. (2007) "*El uso de las plantas a través de la cultura tradicional Lojeña*". Fundación Ibn al-Jatib de Estudios de Cooperación Cultural, Granada.
- Benítez, G., González-Tejero, M.R., Molero-Mesa, J. (2010) Pharmaceutical ethnobotany in the western part of Granada province (southern Spain): ethnopharmacological synthesis. *Journal of Ethnopharmacology*, 129(1), 87-105.
- Benmiloud-Mahieddine, R., Abirached-Darmency, M., Brown, S.C., Kaid-Harche, M., Siljak-Yakovlev, S. (2011) Genome size and cytogenetic characterization of three Algerian *Retama* species. *Tree genetics and Genomes*, 7(5), 987-998.
- Bennett, M.D. (1987) Variation in genomic form in plants and its ecological implications. *New Phytologist*, 106, 177-200.
- Biradar, D.P., Rayburn, A.L. (1993) Heterosis and nuclear DNA content in maize. *Heredity*, 71(3), 300-304.
- Bisby, F.A. (1981) Genisteae (Adans.) Benth. In: "*Advances in Legume Systematics*", R.M. Polhill, P.H. Raven (Eds.), pp. 409-425, Royal Botanic Gardens, Kew.
- Booth, R.E., Grime, J.P. (2003) Effects of genetic impoverishment on plant community diversity. *Journal of Ecology*, 91(5), 721-730.
- Boudjelal, A., Henchiri, C., Sari, M., Sarri, D., Hendel, N., Benkhaled, A., Ruberto, G. (2013) Herbalists and wild medicinal plants in M'Sila (North Algeria): an ethnopharmacology survey. *Journal of Ethnopharmacology*, 148(2), 395-402.
- Boulos, L. (1999) "*Flora of Egypt (Azollaceae-Oxalidaceae)*". Al Hadara Publishing, Cairo. 419p.
- Boulos, L. (2009) "*Flora of Egypt Checklist Revised Annotated Edition*". Al-Hadara Publishing, Cairo. 410p.
- Bouredja, N., Mehdadi, Z., Bendimered, F.Z., Cherifi, K. (2011) Effets de quelques prétraitements physico-chimiques sur la levée de l'inhibition tégumentaire des graines de *Retama monosperma* Boiss. Et recherches des conditions thermiques optimales de germination. *Acta Botanica Gallica*, 158(4), 633-643.
- Caceres, A., Giron, L.M., Martinez, A.M. (1987) Diuretic activity of plants used for the treatment of urinary ailments in Guatemala. *Journal of Ethnopharmacology*, 19(3), 233-245.

- Caravaca, F., Alguacil, M.M., Figueroa, D., Barea, J.M., Roldán, A. (2003) Re-establishment of *Retama sphaerocarpa* as a target species for reclamation of soil physical and biological properties in a semi-arid Mediterranean area. *Forest Ecology and Management*, **182**(1-3), 49-58.
- Cerbah, M., Kevei, Z., Siljak-Yakovlev, S., Kondorosi, E., Kondorosi, A., Trinh, T.H. (1999a) FISH chromosome mapping allowing karyotype analysis in *Medicago trunculata* lines Jemalong J5 and R108-1. *Molecular Plant-microbe Interactions*, **12**(11), 947-950.
- Cerbah, M., Coulaud, J., Brown, S., Siljak-Yakovlev, S. (1999b) Evolutionary DNA variation in the genus *Hypochaeris*. *Heredity*, **82**(3), 261-266.
- Cerbah, M., Mortreau, E., Brown, S., Siljak-Yakovlev, S., Bertrand, H., Lambert, C. (2001) Genome size variation and species relationships in the genus *Hydrangea*. *Theoretical and Applied Genetics*, **103**(1), 45-51.
- Cheriti, A., Talhi, M.F., Belboukhari, N., Taleb, S., Roussel, C. (2009) Removal of copper from aqueous solution by *Retama raetam* Forssk. growing in Algerian Sahara. *Desalin. Water Treatment*, **10**(1-3), 317-320.
- Chiapella, L.F., Velari, T.C., Kosovel, V., Pellizzari, L. (2009) Karyological notes on some genera of *Genisteae* from the Mediterranean region. *Bocconea*, **23**, 187.
- Cramer, W., Bondeau, A., Woodward, F.I., Prentice, I.C., Betts, R.A., Brovkin, V., et al. (2001) Global response of terrestrial ecosystem structure and function to CO<sub>2</sub> and climate change: results from six dynamic global vegetation models. *Global Change Biology*, **7**(4), 357-373.
- Cristofolini, G., Feoli Chiapella, L. (1984) Origin and diversification of *Genisteae* (Fabaceae): a serosystematic purview. *Webbia*, **38**(1), 105-122.
- Dart, P. (1998). Nitrogen fixation by tropical trees and shrubs. In *Biological Nitrogen Fixation for the 21<sup>st</sup> Century. Proceedings of the 11<sup>th</sup> International Congress on Nitrogen Fixation*, Institut Pasteur, Paris, France. doi: 10.1007/978-94-011-5159-7\_427
- De Leonardis, W., Zizza, A. (1994) Palynological study of endemic taxa from Sicily and central-southern Italy. *Acta Botanica Malacitana*, **19**, 217-229.
- Dellafiore, C.M. (2007) Ecología del Conejo Silvestre (*Oryctolagus cuniculus*) en un Sistema Dunar Costero. *Ph.D. Dissertation*, University of Seville, Spain.
- Dellafiore, C.M. Muñoz Vallés, S., Gallego Fernández, J.B. (2006) Rabbits (*Oryctolagus cuniculus*) as dispersers of *Retama monosperma* (L.) Boiss. seeds in a coastal dune system. *Ecoscience*, **13**(1), 5-10.
- Djeddi, S., Karioti, A., Yannakopoulou, E., Papadopoulos, K., Chatter, R., Skaltsa, H. (2013) Analgesic and antioxidant activities of Algerian *Retama raetam* (Forssk.) Webb & Berthel extracts. *Records of Natural Products*, **7**(3), 169-176.
- Doležel, J., Bartoš, J. (2005) Plant DNA flow cytometry and estimation of nuclear genome size. *Annals of Botany*, **95**(1), 99-110.
- Domínguez, E. (1987) *Retama* Raf. In: "*Flora Vasculare de Andalucía Occidental*", B. Valdés, S. Talavera, E. Fernández Galiano, (Eds.), pp. 162-162. Barcelona, Andalucía.
- Eddouks, M., Maghrani, M., Louedec, L., Haloui, M., Michel, J.B. (2007) Antihypertensive activity of the aqueous extract of *Retama raetam* Forssk. leaves in spontaneously hypertensive rats. *Journal of Herbal Pharmacotherapy*, **7**(2), 65-77.
- Edziri, H., Maštouri, M., Chéraif, I., Aouni, M. (2010) Chemical composition and antibacterial, antifungal and antioxidant activities of the flower oil of *Retama raetam* (Forssk.) Webb from Tunisia. *Natural Product Research*, **24**(9), 789-796.
- Edziri, H. Maštouri, M., Mahjoub, M.A., Mighri, Z., Mahjoub, A., Verschaeve, L. (2012) Antibacterial, antifungal and cytotoxic activities of two flavonoids from *Retama raetam* flowers. *Molecules*, **17**(6), 7284-7293.
- El Hamdani, N., Fdil, R. (2015) Evaluation of fatty acids profile and mineral content of *Retama monosperma* (L.) Boiss. of Morocco. *Journal of Materials and Environmental Science*, **6**(2), 538-45.
- El Hamrouni, A. (2001) Partie relative à la flore et la végétation. In: "*Conservation des zones humides littorales et des écosystèmes côtiers du Cap-Bon*",

- L. Baccar, M. Moussa, C. Ben Hamza, (Eds.), pp. 1-33. Med Wet et Coast, Tunisia.
- El-Bahri, L., Djegham, M., Bellil, H. (1999) *Retama raetam*: A poisonous plant of North Africa. *Veterinary and Human Toxicology*, **41**(1), 33-35.
- El-Bana, M.I., Nijs, I., Kockelbergh, F. (2002) Microenvironmental and vegetational heterogeneity induced by phytogenic nebkhas in an arid coastal ecosystem. *Plant and Soil*, **247**, 283-293.
- El-Beyrouthy, M., Arnold, N., Deleis-Dusollier, A., Dupont, F. (2008) Plants used as remedies antirheumatic and antineuralgic in the traditional medicine of Lebanon. *Journal of Ethnopharmacology*, **120**(2), 315-334.
- El-Hilaly, J., Hmammouchi, M., Lyoussi, B. (2003) Ethnobotanical studies and economic evaluation of medicinal plants in Taounate province (Northern Morocco). *Journal of Ethnopharmacology*, **86**(2-3), 149-158.
- El-Mokasabi, F.M. (2014) Floristic composition and traditional uses of plant species at Wadi Alkuf, Al-Jabal Al-Akhder, Libya. *American-Eurasian Journal of Agricultural & Environmental Sciences*, **14**(8), 685-697.
- El-Shazly, A., Ateya, A.M., Witte, L., Wink, M. (1996) Quinolizidine alkaloid profiles of *Retama raetam*, *R. sphaerocarpa* and *R. monosperma*. *Zeitschrift für Naturforschung C*, **51**(5-6), 301-308.
- Evenari, M. (1938) The physiological anatomy of the transpiratory organs and the conducting systems of certain plants typical of the wilderness of Judaea. *Botanical Journal of the Linnean Society*, **51**(340), 389-407.
- Farida, B., Géraldine, D., Abdelghani, B., Djellali, B., Said, B., Gisèle, L. (2009) *Retama* species growing in different ecological-climatic areas of northeastern Algeria have a narrow range of rhizobia that form a novel phylogenetic clade within the Bradyrhizobium genus. *Systematic and Applied Microbiology*, **32**(4), 245-255.
- Fdil, R., El Hamdani, N., El Kihel, A., Sraidi, K. (2012) Distribution of alkaloids in the aerial parts of *Retama monosperma* (L.) Boiss. in Morocco. *Annales de Toxicologie Analytique*, **24**(3), 139-143.
- Feoli Chiappella, L., Prodan, M. (1989) "Contributo sierologico alla sistematica di *Gonocytisus Spach* (Genisteae-Fabaceae)". *Giornale Botanico Italiano*.
- Fernández, J.B.G., Muñoz Vallés, S., Dellafiore, C.M. (2010) Expansión de un arbuŝto nativo-invasivo en dunas costeras: causas consecuencias ecológicas. *Revista Chagual*, **8**, 41-48.
- Ferrauto, G., Guglielmo, A., Lantieri, A., Pavone, P., Salmeri, C. (2015) Pollen morphology and seed germination studies on *Retama raetam* ssp. *gussonei*, endemic subspecies from Sicily. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology*, **149**(2), 251-259.
- Gallego-Martin, F., Sandez Anta, M.A., Navarro Andrés, F. (1988) Acerca de la cariológica de algunas genisteas del centro-occidente español. *Lazaroa*, **9**, 55-60.
- García Novo, F., Merino, J. (1997) Pattern and process in the dune system of the Doñana National Park, Southwestern Spain. In: "Ecosystems of the World", E. van der Maarel (Ed.), pp. 453-468, Elsevier, Amsterdam, the Netherlands.
- GBIF. org (2020) GBIF home page. Website <https://www.gbif.org>. [13 January 2020].
- Ghani, U., Nur-e-Alam, Yousef, M., Ul-Haq, Z., Noman, O.M., Al-Rehaily, A.J. (2019) Natural flavonoid  $\alpha$ -glucosidase inhibitors from *Retama Raetam*: Enzyme inhibition and molecular docking reveal important interactions with the enzyme active site. *Bioorganic Chemistry*, **87**, 736-742.
- Ghirardelli, L.A., Feoli Chiappella, L., Rizzi Longo, L. (1997) Preliminary study on the exina ultrastructure of *Cytisus* s.l. (Genisteae, Fabaceae). *Allionia*, **35**, 197-202.
- Godelle, B., Cartier, D., Marie, D., Brown, S.C., Siljak-Yakovlev, S. (1993) Heterochromatin study demonstrating the non-linearity of fluorometry useful for calculating genomic base composition. *The Journal of the International Society for Analytical Cytology*, **14**(6), 618-626.
- Gómez González, S., Cavieres, L.A., Teneb, E.A., Arroyo, J. (2004) Biogeographical analysis of species of the tribe Cystaseae (Fabaceae) in the Iberian Peninsula and Balearic Islands. *Journal of*

- Biogeography*, **31**(10), 1659-1671.
- González-Mauraza, H., Martín-Cordero, C., Alarcón-de-la-Lastra, C., Rosillo, M.A., León-González, A.J., Sánchez-Hidalgo, M. (2013) Anti-inflammatory effects of *Retama monosperma* in acute ulcerative colitis in rats. *Journal of Physiology and Biochemistry*, **70**, 163-172.
- González-Mauraza, N.H., León-González, A.J., Espartero, J.L., Gallego-Fernández, J.B., et al. (2016) Isolation and quantification of pinitol, a bioactive cyclitol, in *Retama* spp. *Natural Product Communications*, **11**(3), 405-406.
- González-Tejero, M.R., Casares-Porcel, M., Sánchez-Rojas, C.P., Ramiro-Gutiérrez, J.M., Molero-Mesa, J., et al. (2008) Medicinal plants in the Mediterranean area: synthesis of the results of the project Rubia. *Journal of Ethnopharmacology*, **116**(2), 341-357.
- Greuter, W., Burdet, H. M., Long, G. (1989) "*Med-checklist*". Conservatoire et jardin botaniques de la ville de Genève, Genève.
- Guerrouj, K., Ruíz-Díez, B., Chahboune, R., Ramírez-Bahena, M.H., Abdelmoumen, H., et al. (2013) Definition of a novel symbiovar (sv. *retamae*) within *Bradyrhizobium retamae* sp. nov., nodulating *Retama sphaerocarpa* and *Retama monosperma*. *Systematic and Applied Microbiology*, **36**(4), 218-223.
- Gutterman, Y. (1993) "*Seed Germination in Desert Plants*". Springer-Verlag, Berlin. doi: 10.1007/978-3-642-75698-6
- Guttridge, C.G., Woodly, S.E., Hunter, T. (1984) Accelerating strawberry seed germination by fungal infection. *Annals of Botany*, **54**(2), 223-230.
- Haase, P., Pugnaire, F.I., Fernández, E.M., Puigdefábregas, J., Clark, S.C., Incoll, L.D. (1996) An investigation of rooting depth of the semiarid shrub *Retama sphaerocarpa* (L.) Boiss. by labelling ground water with a chemical tracer. *Journal of Hydrology*, **177**(1-2), 23-31.
- Hammouche-Mokrane, N., León-González, A.J., Navarro, I., Boulila, F., Benallaoua, S., Martín-Cordero, C. (2017) Phytochemical profile and antibacterial activity of *Retama raetam* and *R. sphaerocarpa* cladodes from Algeria. *Natural Product Communications*, **12**(12). doi: 10.1177/1934578x1701201211
- Hayet, E., Samia, A., Patrick, G., Ali, M.M., Maha, M., Laurent, G., Mighri, Z., Mahjoub, L. (2007) Antimicrobial and cytotoxic activity of *Marrubium alysson* and *Retama raetam* grown in Tunisia. *Pakistan Journal of Biological Sciences*, **10**(10), 1759-1762.
- Hayet, E., Maha, M., Samia, A., Mata, M., Gros, P., Raida, H., et al. (2008) Antimicrobial, antioxidant, and antiviral activities of *Retama raetam* (Forssk.) Webb flowers growing in Tunisia. *World Journal of Microbiology and Biotechnology*, **24**(12), 2933-2940.
- Helmeyer, I., Schönig, H. (2012) "*Herbal Medicine in Yemen: Traditional Knowledge and Practice, and their Value for Today's World*". Brill, Leiden. doi: 10.1163/9789004232075
- Helmstädter, A. (2016) Ethnopharmacology in the work of Melville William Hilton-Simpson (1881–1938). historical analysis and current research opportunities. *Die Pharmazie-An International Journal of Pharmaceutical Sciences*, **71**(6), 352-360.
- Herrera, J. (1988) Pollination relationships in southern Spanish Mediterranean shrublands. *The Journal of Ecology*, **76**, 274-287.
- Herrera, J. (2001) The variability of organs differentially involved in pollination, and correlations of traits in *Genisteae* (Leguminosae: Papilionoideae). *Annals of Botany*, **88**(6), 1027-1037.
- Hudaib, M., Mohammad, M., Buštanji, Y., Tayyem, R., Yousef, M., et al. (2008) Ethnopharmacological survey of medicinal plants in Jordan, Mujib Nature Reserve and surrounding area. *Journal of Ethnopharmacology*, **120**(1), 63-71.
- Ishurd, O., Kermagi, A., Zgheel, Flefla, F., Elmabruk, M., Yalin, W., et al. (2004) Structural aspects of water-soluble galactomannans isolated from the seeds of *Retama raetam*. *Carbohydrate Polymers*, **58**(1), 41-44.
- Izhaki, I., Neeman, G. (1997) Hares (*Lepus* spp.) as seed dispersers of *Retama raetam* (Fabaceae) in a sandy landscape. *Journal of Arid Environments*, **37**(2), 343-354.

- Jacobsen, E. (2000) *Retama monosperma*. In: "Invasive Plants of California's Wildlands. Berkeley", C.M. Bossard, J.M. Randall, M.C. Hoshovny (Eds.), pp. 266-268. University of California Press, USA.
- Jafri, S.M.H. (1980) Fabaceae. In: "Flora of Libya", S.M.H. Jafri, A. El-Gadi (Eds.), pp. 31-33. Tripoli University, Libya.
- Jahandiez, E., Maire, D.R. (1932) "Catalogue des Plantes du Maroc". Tome Deuxième, Alger Imprimerie Minerva, En Vente a Paris.
- Kassem, M., Mosharrafa, S.A., Saleh, N.A.M., Abdel-Wahab, S.M. (2000) Two new flavonoids from *Retama raetam*. *Fitoterapia*, **71**(6), 649-654.
- Keeler, R.F. (1969) Toxic and teratogenic alkaloids of western range plants. *Journal of Agricultural and Food Chemistry*, **17**(3), 473-482.
- Kigel, J. (1995) Seed germination in arid and semiarid regions. In: "Seed Development and Germination", J. Kigel, G. Galili (Eds.), pp. 645-699. Marcel Dekker, New York.
- Kildisheva, O.A. (2019) Improving the outcomes of seed-based restoration in cold and hot deserts: an investigation into seed dormancy, germination, and seed enhancement. *Ph.D. Dissertation*, University of Western Australia, Perth.
- Kinghorn, A.D., Balandrin, M.F. (1984) Quinolizidine alkaloids of the Leguminosae: Structural types, analysis, chemotaxonomy, and biological activities. In: "Alkaloids: Chemical and Biological Perspectives", W.S. Pelletier (Eds.), pp. 105-148. Wiley, New York.
- Lamin, H., Alami, S., Bouhnik, O., ElFaik, S., Abdelmoumen, H., Bedmar, E. J., Missbah-El Idrissi, M. (2019) Nodulation of *Retama monosperma* by *Ensifer aridi* in an abandoned lead mine soils in eastern morocco. *Frontiers in Microbiology*, **10**, 1456. doi: 0.3389/fmicb.2019.01456
- Lamrabet, M., Chaddad, Z., Bouhnik, O., Alami, S., Kaddouri, K., Bennis, M., et al. (2023) Different species of *Bradyrhizobium* from symbiovars genistearum and retamae nodulate the endemic *Retama dasycarpa* in the High Atlas Mountains. *FEMS Microbiology Ecology*, **99**(5), fiad038. doi: 10.1093/femsec/fiad038.
- Laudadio, V., Tufarelli, V., Dario, M., Hammadi, M., Seddik, M.M., Lacalandra, G.M., Dario, C. (2009) Chemical and nutritional characteristics of halophytes plants used by camels in Southern Tunisia. *Tropical Animal Health and Production*, **41**, 209-215.
- Lems, K. (1960) "Floristic Botany of the Canary Islands: a compilation of the geographic distribution, dispersal types, life forms and leaf types of the species of vascular plants". Institut Botanique de l'Université de Montréal, Montréal.
- Leon-Gonzalez, A.J., Navarro, A.J.I., Acero, N., Mingarro, D.M., Cordero, C.M. (2018) Genus *Retama*: A review on traditional uses, phytochemistry, and pharmacological activities. *Phytochemistry Reviews*, **17**, 701-731.
- López, J., Rodríguez-Riaño, T., Ortega-Olivencia, A., Devesa, J.A., Ruiz, T. (1999) Pollination mechanisms and pollen-ovule ratios in some Genisteae from SW Europe. *Plant Systematics and Evolution*, **216**, 23-47.
- López-Lázaro, M., Martín-Cordero, C., Iglesias-Guerra, F., González, M. A. (1998) An isoflavone glucoside from *Retama sphaerocarpa* Boissier. *Phytochemistry Reviews*, **48**(2), 401-402.
- López-Pintor, A., Espigares, T., Rey-Benayas, J.M., Gómez-Sal, A. (2000) Effect of simulated parent-created micro environmental conditions on germination of *Retama sphaerocarpa* L. Boiss seeds. *Journal of Mediterranean Ecology*, **1**, 219-226.
- Louaar, S., Akkal, S., Boussetla, A., Medjroubi, K., Djarri, L., Seguin, E. (2005) Phytochemical study of *Retama sphaerocarpa*. *Chemistry of Natural Compounds*, **41**, 107-108.
- Maghrani, M., Michel, J.B., Eddouks, M. (2005a) Hypoglycaemic activity of *Retama raetam* in rats. *Phytotherapy Research*, **19**(2), 125-128.
- Maghrani, M., Zeggwagh, N.A., Haloui, M., Eddouks, M. (2005b) Acute diuretic effect of aqueous extract of *Retama raetam* in normal rats. *Journal of Ethnopharmacology*, **99**(1), 31-35.
- Mahdhi, M., Nzoue, A., de Lajudie, P., Mars, M. (2008) Characterization of root-nodulating bacteria on *Retama raetam* in arid Tunisian soils. *Progress in Natural Science*, **18**(1), 43-50.

- Maire, R. (1987) "Flore de l'Afrique du Nord". Le Chevalier, Paris.
- Mechergui, K., Mahmoudi, H., Khouja, M.L., Jaouadi, W. (2017) Factors influencing seed germination of the pastoral plant *Retama raetam* subsp. *bovei* (Fabaceae): interactive effects of fruit morphology, salinity, and osmotic stress. *Biologija*, **63**(2), 134-151.
- Mehdadi, Z., Bendimered, F.Z., Dadach, M., Aisset, A. (2017) Effects of temperature and salinity on the seeds germination of *Retama raetam* (Forssk.) Webb. scarified with sulfuric acid. *Journal of Fundamental and Applied Sciences*, **9**(3), 1284-1299.
- Merghoub, N., Benbacer, L., Amzazi, S., Morjani, H., El-Mzibri, M. (2009) Cytotoxic effect of some Moroccan medicinal plant extracts on human cervical cell lines. *Journal of Medicinal Plants Research*, **3**(12), 1045-1050.
- Merghoub, N., Benbacer, L., El Btaouri, H., Ait Benhassou, H., Terryn, C., et al. (2011) *In vitro* antiproliferative effect and induction of apoptosis by *Retama monosperma* L. extract in human cervical cancer cells. *Cellular and Molecular Biology*, **57**(2), 1581-1591.
- Merquiol, E., Pnueli, L., Cohen, M., Simovitch, M., Rachmilevitch, S., et al. (2002) Seasonal and diurnal variations in gene expression in the desert legume *Retama raetam*. *Plant, Cell and Environment*, **25**(12), 1627-1638.
- Mittler, R., Merquiol, E., Hallak-Herr, E., Rachmilevitch, S., Kaplan, A., Cohen, M. (2001) Living under a 'dormant' canopy: a molecular acclimation mechanism of the desert plant *Retama raetam*. *The Plant Journal*, **25**(4), 407-416.
- Morales Mendez, A., Gonzalez Gonzalez, A., Diaz Rodriguez, F. (1971) "Alkaloids from the Bark of *Retama monosperma*". Revista de la Facultad de Farmacia, Universidad Central de Venezuela, Los Andes.
- Moro, M.J., Pugnaire, F.I., Haase, P., Puigdefàbregas, J. (1997) Mechanism of interaction between *Retama sphaerocarpa* and its under storey layer in a semi-arid environment. *Ecography*, **20**, 175-184.
- Morsy, A.A., Khatab, H., El Sherbiny, E.A., Eldemirdash, J. (2015) Floristic Diversity and Vegetation Analysis of Wadi Sudr, South-West Sinai Peninsula. *Täeckholmia*, **35**(1), 99-119.
- Mouhajir, F. (2002) Medicinal plants used by Berber and Arab peoples of Morocco: ethnopharmacology and phytochemistry. *Ph.D. Dissertation*, University of British Columbia, Canada.
- Muñoz Vallés, S., Gallego Fernández, J.B., Dellafiore, C. (2005) Spatial and temporal pattern of *Retama monosperma* spread in the spit of El Rompido (Gulf of Cadiz, SW Spain). In: "Abstract Book of the European IALE Congress, Landscape Ecology in the Mediterranean, Inside and Outside Approaches", R.G.H. Bunce, R.H.G. Jongman (Eds.), pp. 101. Faro, Portugal.
- Muñoz Vallés, S., Gallego Fernández, J.B., Dellafiore, C., Cambrollé, J. (2011) Effects on soil, microclimate and vegetation of the native-invasive *Retama monosperma* (L.) in coastal dunes. *Plant Ecology*, **212**(2), 169-179.
- Muñoz Vallés, S., Gallego Fernández, J.B., Cambrollé, J. (2013) The biological flora of coastal dunes and wetlands: *Retama monosperma* (L.) Boiss. *Journal of Coastal Research*, **29**(5), 1101-1110.
- Nur-e-Alam, M., Yousaf, M., Parveen, I., Hafizur, R.M., Ghani, U., et al. (2019) New flavonoids from the Saudi Arabian plant *Retama raetam* which stimulates secretion of insulin and inhibits  $\alpha$ -glucosidase. *Organic & Biomolecular Chemistry*, **17**(5), 1266-1276.
- Ohri, D. (1998) Genome size variation and plant systematics. *Annals of Botany*, **82**(A), 75-83.
- Ouf, S.A. (1994) Colonization of *Retama raetam* seeds by fungi and their significance in seed germination. *Mycoscience*, **35**, 53-57.
- Ould El Hadj, M., Hadj-Mahammed, M., Zabeirou, H. (2003) Place des plantes spontanées dans la médecine traditionnelle de la région de Ouargla (Sahara septentrional est). *Courrier du Savoir Scientifique et Technique*, **3**, 47-51.
- Pardo, C., Tahiri, H., Cubas, P., El Alaoui-Faris, F.E.E. (2000) Pollen morphology in *Cytisus* (Papilionoideae, Leguminosae) from Morocco and the Iberian Peninsula. *Grana*, **39**(4), 159-168.
- Pnueli, L., Hallak-Herr, E., Rozenberg, M., Cohen,

- M., Goloubinoff, P., Kaplan, A., Mittler, R. (2002) Molecular and biochemical mechanisms associated with dormancy and drought tolerance in the desert legume *Retama raetam*. *The Plant Journal*, **31**(3), 319-330.
- Polhill, R.M. (1976) *Genisteae* (Adans.) Benth. and related tribes (Leguminosae). *Botanical Systematics*, **1**, 143-368.
- Polunin, O. (1969) "*Flowers of Europe*". Oxford University press, New York.
- POWO (2023) Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet. Website <http://www.plantsoftheworldonline.org/> [Accessed 30 May 2023].
- Prieto, I., Kikvidze, Z., Pugnaire, F.I. (2010) Hydraulic lift: soil processes and transpiration in the Mediterranean leguminous shrub *Retama sphaerocarpa* (L.) Boiss. *Plant Soil*, **329**, 447-456.
- Qiuyin, C., Huachen, W. (1996) Effect of dietary genistein on antioxidant enzyme activities in SENCAR mice. *Nutrition*, **25**(1), 1-7.
- Quezel, P., Santa, S. (1962) "*Nouvelle flore de l'Algérie et des régions désertiques méridionales*". Editions du Centre National de la Recherche Scientifique, France.
- Randall, J.M. (1997) Weed Alert! New invasive weeds in California. *proceedings of the California Exotic Pest Plant Council*, California, USA.
- Randall, R.P. (2007) "*The Introduced Flora of Australia and Its Weed Status*". CRC for Australian Weed Management, Adelaide.
- Rebbas, K., Bounar, R., Gharzouli R., Ramdani, M., Djellouli, Y., Alatou, D. (2012) Plantes d'intérêt médicinale et écologique dans la région d'Ouanougha (M'sila, Algérie). *Phytothérapie*, **10**, 131-142.
- Rizzi Longo, L., Ghirardelli, L.A., Feoli Chiapella, L. (2006) Pollen morphology and taxonomy of *Genista* sect. *Cephalospartum* Spach emend. P. Gibbs (*Genisteae*, Fabaceae). *Flora Mediterranea*, **16**, 169-191.
- Rodríguez-Echeverría, S., Pérez-Fernández, M.A. (2003) Soil fertility and herb facilitation mediated by *Retama sphaerocarpa*. *Journal of Vegetation Science*, **14**(6), 807-814.
- Rodríguez-Echeverría, S., Pérez-Fernández, M.A., Vlaar, S., Finnan, T. (2003) Analysis of the legume-rhizobia symbiosis in shrubs from central western Spain. *Journal of Applied Microbiology*, **95**(6), 1367-1374.
- Rodríguez-Riaño, T., Ortega-Olivencia, A., Devesa, J.A. (1999a) "*Biología Floral en Fabaceae*". Editorial CSIC- CSIC press, Madrid.
- Rodríguez-Riaño, T., Ortega-Olivencia, A., Devesa, J.A. (1999b) Reproductive phenology in three *Genisteae* (Fabaceae) shrub species of the W Mediterranean region. *Nordic Journal of Botany*, **19**(3), 345-354.
- Rodríguez-Riaño, T., Ortega-Olivencia, A., Devesa, J. A. (1999c). Reproductive biology in two *Genisteae* (Papilionoideae) endemic of the western Mediterranean region: *Cytisus striatus* and *Retama sphaerocarpa*. *Canadian Journal of Botany*, **77**(6), 809-820.
- Roskov, Y., Zarucchi, J., Novoselova, M., Bisby, F. (Eds.) (2020) ILDIS: ILDIS World Database of Legumes (version 12, May 2014). In: "*Naturalis, Leiden. the Netherlands. Species 2000 & ITIS Catalogue of Life*". Roskov, Y., Ower, G., Orrell, T., Nicolson, D., Bailly, N., Kirk, P.M., Bourgoin, T., DeWalt, R.E., Decock, W., Nieukerken, E. & van Penev, L. (Eds.). 2020-04-16 Beta Digital resource at [www.catalogueoflife.org/col](http://www.catalogueoflife.org/col).
- Saada, M., Falleh, H., Catarino, M.D., Cardoso, S.M., Ksouri, R. (2018) Plant growth modulates metabolites and biological activities in *Retama raetam* (Forssk.) Webb. *Molecules*, **23**(9), 2177. doi: 10.3390/molecules23092177
- Sadik, K., Hamdani, N.E., Hachim, M.E., Byadi, S., Bahadur, I., Aboulmouhajir, A. (2020) Towards a theoretical understanding of alkaloid-extract Cytisine derivatives of *Retama monosperma* (L.) Boiss. Seeds, as eco-friendly inhibitor for carbon steel corrosion in acidic 1M HCl solution. *Journal of Theoretical and Computational Chemistry*, **19**(05), 2050013. doi: 10.1142/s0219633620500133
- Said, O., Khalil, K., Fulder, S., Azaizeh, H. (2002) Ethnopharmacological survey of medicinal herbs in *Egypt. J. Bot.* **64**, No. 2 (2024)

- Israel, the Golan Heights and the West Bank region. *Journal of Ethnopharmacology*, **83**(3), 251-265.
- Salama, F., El-Tayeh, N., Zaher, A., El- Naggar, S., Gaafar, A. (2023) Phytosociological Studies on the Associated Species of *Balanites aegyptiaca* In The Eastern and Western Egyptian Deserts. *Egyptian Journal of Botany*, **63**(3), 1005-1029. doi: 10.21608/ejbo.2023.205410.2308.
- Salatino, A., Gottlieb, O.R. (1980) Quinolizidine alkaloids as systematic markers of the Papilionoideae. *Biochem. Systematics and Ecology*, **8**(2), 133-147.
- Schmid, T., Turner, D., Oberbaum, M., Finkelstein, Y., Bass, R., Kleid, D. (2006) Respiratory failure in a neonate after folk treatment with broom bush (*Retama raetam*) extract. *Pediatric Emergency Care*, **22**(2), 124-126.
- Scholze, M., Knorr, W., Arnell, N.W., Prentice, I.C. (2006) "A Climate-change Risk Analysis for World Ecosystems". Carnegie Institution of Washington, Stanford, CA. 103(35), 13116-13120. doi: 10.1073/pnas.0601816103
- Seglie, L., Scariot, V., Larcher, F., Devecchi, M., Chiavazza, P.M. (2012) *In vitro* seed germination and seedling propagation in *Campanula* spp. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology*, **146**(1), 15-23.
- Sekina, M.A., Moore, P.D. (1995) Morphological studies of the pollen grains of the semi-arid region of Egypt. *Flora*, **190**(2), 115-133.
- Selami, N., Auriac, M.C., Catrice, O., Capela, D., Kaid-Harche, M., Timmers, T. (2014) Morphology and anatomy of root nodules of *Retama monosperma* (L.) Boiss. *Plant Soil*, **379**, 109-119.
- Steponkus, P.L. (1982) Responses to extreme temperatures. Cellular and subcellular bases. In: "Encyclopedia of Plant Physiology 12A Physiological Plant Ecology", pp. 372-401, O. Lang, C.B. Osmond, P.S. Nobel (Eds.), Springer, Berlin. doi: 0.1007/978-3-642-68090-8\_13
- Täckholm, V. (1974) "Student's flora of Egypt". Cairo University Press, Cairo. 888p.
- Tahraoui, A., El-Hilaly, J., Israili, Z.H., Lyoussi, B. (2007) Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in south-eastern Morocco (*Errachidia province*). *Journal of Ethnopharmacology*, **110**(1), 105-117.
- Talavera, S. (1999a) *Echinopartum* (Spach) Fourr. In: "Flora Iberica", S. Talavera, C. Aedo, S. Castroviejo, C. Romero Zarco, L. Sáez, F. J. Salgueiro, M. Velayos (Eds.), pp. 119-127. Madrid.
- Talavera, S. (1999b) *Retama* Raf. In: "Flora Iberica", S. Talavera, C. Aedo, S. Castroviejo, C. Romero Zarco, L. Sáez, F. J. Salgueiro, M. Velayos (Eds.), pp. 137-141. Madrid.
- Teixidor-Toneu, I., Martin, G.J., Ouhammou, A. (2016) An ethnomedicinal survey of a Tashelhit-speaking community in the High Atlas, Morocco. *Journal of Ethnopharmacology*, **188**, 96-110.
- Telli, A., Esnault, M.A., Khelil, A.O. (2016) An ethnopharmacological survey of plants used in traditional diabetes treatment in south-eastern Algeria (Ouargla province). *Journal of Arid Environments*, **127**, 82-92.
- Touati, D., Allain, P., Pellecuer, J., Fkih-tetouani, S., Agoumi, A. (1996) Alkaloids from *Retama monosperma* ssp. *Eumonosperma*. *Fitoterapia*, **67**(1), 49-52.
- Touati, R., Santos, S.A., Rocha, S.M., Belhamel, K., Silvestre, A.J. (2015) *Retama sphaerocarpa*: An unexploited and rich source of alkaloids, unsaturated fatty acids and other valuable phytochemicals. *Industrial Crops and Products*, **69**, 238-243.
- Touati, R., Santos, S.A., Rocha, S.M. Belhamel, K., Silvestre, A.J. (2017) Phenolic composition and biological prospecting of grains and stems of *Retama sphaerocarpa*. *Industrial Crops and Products*, **95**, 244-255.
- Trieu, V.N., Dong, Y., Zheng, Y., Uckun, F.M. (1999) In vivo antioxidant activity of genistein in a murine model of singlet oxygen-induced cerebral stroke. *Radiation Research*, **152**(5), 508-516.
- Tutin, T.G., Heywood, V.H., Burges, N.A., Moore, D.M., Valentine, D.H., Walters, S.M., Webb, D.A. (1968) "Flora Europaea (Rosaceae-Umbelliferae)". Cambridge University Press, Cambridge.
- Valladares, F., Villar-Salvador, P., Domínguez, S., Fernández-Pascual, M., Peñuelas, J.L., Pugnaire, I.

- F.I. (2002) Enhancing the early performance of the leguminous shrub *Retama sphaerocarpa* (L.) Boiss.: Fertilisation versus *Rhizobium* inoculation. *Plant Soil*, **240**, 253-262.
- Viegi, L., Ghedira, K. (2014) Preliminary study of plants used in ethnoveterinary medicine in Tunisia and in Italy. *African Journal of Traditional, Complementary and Alternative Medicines*, **11**(3), 189-199.
- Webb, P.B. (1853) "*Plantarum Rabiorum*". Visctor Masson, Bibliopola, Parisis.
- WFO (2023) World flora online. Published on the internet. Website <http://www.worldfloraonline.org>. [ Accessed 30 May 2023].
- Wickens, G.E. (1998) "*Ecophysiology of Economic Plants in Arid and Semi-arid Lands*". Springer, United Kingdom.
- Wilson, J.B., Sykes, M.T. (1999) Is zonation of coastal sand dunes determined primarily by sand burial or by salt spray? A test in New Zealand dunes. *Ecology Letters*, **2**, 233-236.
- Wink M. (1993) Allelochemical properties or the raison d' être of alkaloids. In: "*The Alkaloids*", G.A.C. G.A.C. Ordell, (Eds.), pp. 1-118. Academic Press, San Diego. doi: 10.1016/s0099-9598(08)60134-0
- Xu, W. H., Al-Rehaily, A. J., Yousaf, M., Ahmad, M.S., Khan, S.I., Khan, I.A. (2015) Two new flavonoids from *Retama raetam*. *Helvetica Chimica Acta*, **98**(4), 561-568.
- Yaniv, Z., Dudai, N. (2014) "*Medicinal and Aromatic Plants of the Middle-east*". Springer, Berlin. 342p.
- Youssef, A.M. (2009) Seed germination of some desert plants from Egypt. *Journal of Applied Sciences Research*, **5**(2), 144-150.
- Youssef, R.A., Amer, W.M., Azza, B.H. (2023) Genus *Retama* Raf. 1838 (Fabales, fabaceae): Taxonomic revision in Egypt supported by molecular finger printing. *Bulletin of The Iraq Natural History Museum*, **17**(93), 435-458.
- Zakhia, F., Jeder, H., Willems, A., Gillis, M., Dreyfus, B., De Lajudie, P. (2006) Diverse bacteria associated with root nodules of spontaneous legumes in Tunisia and first report for nifH-like gene within the genera *Microbacterium* and *Starkeya*. *Microbial ecology*, **51**, 375-393.
- Zamberlin, Š., Antunac, N., Havranek, J., Samaržija, D. (2012) Mineral elements in milk and dairy products. *Mljekarstvo: Časopis za Unaprjeđenje Proizvodnje i Prerade Mlijeka*, **62**(2), 111-125.
- Zimmermann, M. (1988) Nectar production, flowering phenology, and strategies for pollination. In: "*Plant reproductive ecology. "Patterns and Strategies"*", J. Lovett Doust, L. Lovett Doust (Eds.). pp. 157-178, Oxford University Press, New York.
- Zinnert, J.C., Nelson, J.D., Hoffman, A.M. (2012) Effects of salinity on physiological responses and the photochemical reflectance index in two co-occurring coastal shrubs. *Plant Soil*, **354**, 45-55.
- Zohary, M. (1972) "*Flora Palaestina (Platanaceae-Umbelliferae)*", pp. 47-48. The Israel academy of Sciences and Humanities, Jerusalem.
- Zunzunegui, M., Esquivias, M.P., Fernández-González, P., Valera-Burgos, J., Diaz Barradas, M. C., Gallego-Fernández, J.B. (2017) Morphophysiological response of *Retama monosperma* to extreme salinity levels. *Ecohydrology*, **10**(7), e1871.

## ضوء على أنواع الرتم الصحراوية بمنطقة البحر المتوسط المستديمة الخضرة وذات السيقان التمثيلية

ريهام يوسف (1)، ووفاء عامر (1)، وعمران غالي (2)، وعزة حامد (1)

(1) معشبة جامعة القاهرة، قسم النبات والميكروبيولوجي، كلية العلوم، جامعة القاهرة، الجيزة، مصر،

(2) معشبة مركز بحوث الصحراء- القاهرة - مصر.

جنس الرتم (الفصيلة البقولية) هو جنس من منطقة البحر الأبيض المتوسط، وقد تم توطينه في العالم الجديد مؤخرًا. وهذا الجنس يشمل أربعة أنواع من الرتم هي ديزيكاربا مونوسيرما والريثاما والسفيروكاربا. وكلها أنواع صحراوية ذات سيقان تمثيلية. أما الأنواع المصاحبة لها فدائمًا ما تتغير بتغير الموقع الجغرافي للنوع قيد الدراسة. أما حبوب اللقاح في جنس الريثاما فهو فردي وثلاثي الشقوق وبعض الأزهار تفتقر إلى الرحيق، والبعض الآخر منتج للرحيق وتلقح بالحشرات. ويتميز جنس الريثاما بأنه يمتلك العديد من آليات المقاومة والدفاع واستراتيجيات التأقلم والتكيف مع ضغوط الجفاف والملوحة بما في ذلك جينات الاستجابة للضغط والتكيف الفسيولوجي والميكروبيولوجي. وتلعب ميكروبات التربة والميكروبات الداخلية دورًا مهمًا في تحسين تأقلم أنواع الريثاما تحت الضغوط البيئية. وبنس الريثاما ذات صبغيات رباعية التضاعف ويوجد بها ظاهرة الكروموسومات الكاريولوجية المتجانسة. وحجم الجينوم مرتبط بطبيعة البيئية.

غطت هذه المراجعة تصنيف أنواع الريثاما وأهميتها في الأراضي الفاحلة وخاصة دورها في حماية التربة واستدامته وخصوبتها وإعادة تأهيل الصحراء. كما تم حصر المركبات الكيميائية في أنواع الريثاما بما في ذلك مركبات الفلافونويد، والزيوت العطرية، والقلويدات، والتربين، والمنشطات، والأحماض الدهنية، والسكريات، والتركييب المعدني. بالإضافة إلى استخداماته المتعددة الأغراض كعلف واستخدامات نباتية وأنشطة دوائية، وتتميز الريثاما بوجود مركبات مضادة للسرطان ومضادات الأكسدة وخافضه لسكر الدم ومضاد للالتهابات ومسكن ومضاده لسرطان الدم ومضاده للبكتيريا والفطريات وغيرها.