

Evaluation of Gluten-Free Flatbread Substituted with Some Cereals and Legumes

^{*1}Hayat, H. Abd-Elsattar, ² Hanan, M.A. EL-Ghandour & ³Nahed, L. Zaki

¹Crops Technology Research Department, Food Technology Research Institute, Agricultural Research Center, Giza, Egypt.

²Regional Center for Food and Feed, Agricultural Research Center, Giza, Egypt.

³Experimental Kitchen Research unit, Food Technology Research Institute, Agricultural Research Center, Giza, Egypt.

Original Article

ABSTRACT

Article information

Received 09/01/2025 Revised 22/01/2025 Accepted 26/01/2025 Published 05/02/2025 Available online 01/03/2025

Keywords

Celiac disease, protein quality, legumes and cereals. albino rats There is a growing consumer demand for functional gluten-free foods, as products containing gluten can trigger celiac disease in genetically susceptible individuals. This research aimed to develop a gluten-free flatbread with enhanced nutritional quality and functional properties. The flatbread was prepared using legume flour (lentil or chickpea) in combination with rice and corn flour. Sensory evaluation, chemical composition, and physical properties of various formulations were analyzed. The results showed that gluten-free flatbread made from 100% lentil or 100% chickpea had higher protein content compared to other formulations. The addition of chickpeas and lentils significantly reduced hardness while increasing springiness and resilience. Flatbreads made from 100% chickpea or 100% lentil remained fresher than those made from corn flour by the end of the storage period. The highest sensory scores were attributed to combinations of (50% lentil/rice), (50% chickpea/ rice), (50% lentil/corn), and (50% chickpea/corn). The study also examined the effects of various legume/cereal blends on albino rats, focusing on Biological Value (BV), True Digestibility (TD), and Net Protein Utilization (NPU) of the flatbread formulations. The Biological Value (BV) of the recipes was notably high, with all exceeding 70%. The Protein Efficiency Ratio (PER) of the gluten-free flatbreads ranged between 2.37 and 3.0, surpassing the standard value of 2.5. The NPU values were significantly influenced by the proportion and quality of protein. In conclusion, it is possible to create gluten-free flatbreads with quality similar to conventional bread.

1. Introduction

Celiac disease is a chronic inflammatory, autoimmune-mediated disorder that primarily affects the duodenum (Caio et al., 2019). It is classified as a digestive disorder that damages the villi—tiny, hair-like projections in the small intestine responsible for nutrient absorption leading to an immune reaction against gluten (Barada et al., 2010). This condition results in a continuous intolerance to gluten, gliadin, and related prolamins found in wheat, rye, and barley. The disease is characterized by intestinal damage caused by an immune system defect (autoimmune response) that occurs in genetically predisposed individuals (Amin et al., 2002). A gluten-free diet requires the consumption of gluten-free cereals such as rice, corn, sorghum, millet, and teff, as well as pseudo-cereals like buckwheat, amaranth, quinoa, and canthus. Additionally, naturally gluten-free foods, including potatoes, legumes, oilseeds, tapioca, nuts, fruits, and vegetables, are essential dietary components (Pruska-Kedzior et al., 2008). Gluten-free bread formulated with rice flour and gums significantly affects dough properties. Among various gums, xanthan and xanthan–guar gum mixtures have been found to be the most effective in improving dough structure. The best firmness and specific volume values were observed in bread containing xanthan and xanthan-guar gum (Demirkesen et al., 2010). Several types of flour, such as barley, corn, cassava, and chickpea, have been extensively studied for their potential use in composite flour bread (Ali et al., 2000). For example, blending rice flour with corn and cassava starch can produce gluten-free bread with a well-structured crumb, enjoyable flavor, and appealing appearance (Lopez et al., 2004; Gujral & Rosell, 2004). However, traditional gluten -free bread often has a less flexible crumb that hardens quickly and is prone to crumbling. Additionally, the taste characteristics of these products vary depending on the ingredients used. According to the FAO, using composite flour in various food products could be economically beneficial by reducing or even eliminating the demand for wheat in bread and pastry production through the use of domestically grown ingredients (Jisha et al., 2008). Despite the availability of gluten-free products, their nutritional quality remains a significant concern due to typically lower levels of protein, fiber, and essential nutrients (Conte et al., 2019). However, ensuring the nutritional value of gluten-free products is crucial, given the growing population affected by celiac disease (Aprodu et al., 2016). Rice contains high levels of aspartic and glutamic acids, with lysine being the limiting amino acid (FAO, 2004). It is considered the primary staple food in many countries and serves as an important source of vitamins, including riboflavin, thiamine, and niacin. Additionally, rice is rich in aspartic acid, thiamine, niacin, vitamin B6, and folate (Watson, 1997). Maize flour also contains high levels of essential vitamins and minerals, including potassium, phosphorus, zinc, calcium, and iron. Among the gluten-free cereals available, lentils and chickpeas are winter legumes grown in the highlands of Upper Egypt (Kadria et al., 2021). Legume proteins can be incorporated into baked products to enhance protein content and improve amino acid composition (Bojňanská et al., 2012; Mohammed et al., 2012). Xu et al. (2019) investigated the thermal, pasting, and chemical properties of lentil, chickpea, and yellow pea flours, as well as their moisture adsorption characteristics. Their findings showed that total starch content de-

creased in germinated yellow pea and lentil, whereas germination had no significant effect on chickpeas. However, increasing the proportion of lentils and chickpeas in baked bread negatively impacted the qualitative parameters, particularly loaf volume and volume efficiency (Bojňanská et al., 2012). Chickpeas are an excellent source of protein and carbohydrates (Kaur & Singh, 2005). Akubor et al. (2003) formulated blends of plantain and legume flours, producing gluten-free products with high nutritional value and acceptable sensory properties. Proteins are essential structural and functional components of body cells (Michaelsen et al., 2003). They play a crucial role in the development of skin and muscles, as well as in hormone and enzyme production (Arnarson, 2017). Functional foods resemble conventional foods but provide additional physiological benefits, reducing the risk of chronic diseases beyond basic nutritional functions (Flamm et al., 2001). Traditionally, protein digestibility has been estimated through in vitro fermentation models, and in recent years, several trials have emerged assess the biological value of proteins (Marinangeli & House, 2017). The objective of this study was to develop a novel gluten-free flatbread suitable for individuals with gluten allergies, while enhancing its nutritional quality and functional properties using rice, corn, lentils, and chickpeas. The study aimed to evaluate the flatbread's physical, chemical, nutritional, and sensory properties, as well as its staling behavior, compared to a control sample. Additionally, it aimed to improve the biological value and digestibility of various cereallegume mixtures as a potential complementary gluten-free food.

2. Materials and Methods Materials

Broken rice from a low-amylose (15.27%) variety of Sakha 104 (Japonica qhti unit, Damietta, Egypt) was used. The percentage of broken rice kernels was 27%. Chickpeas, lentils, crushed fava beans, white beans, corn flour, salt, cumin, and sunflower oil were purchased from a local market in Cairo, Egypt.

Food Technology Research Journal, Vol. 7, issue 1, 44-61, 2025

Methods Preparation of rice and legume flours

Rice and legumes, including chickpeas, lentils, crushed fava beans, and white beans, were carefully cleaned to remove impurities and then washed with tap water. They were soaked in tap water for 8 hours at room temperature ($25 \pm 2^{\circ}$ C), following the method outlined by Khattab and Arntfield (2009). After soaking, the legumes were drained and chopped using a Moulinex 750-watt French blender. The chopped mixture was then dried to produce flakes, which were ground into flour using a Universal high-speed grinder (MDY-200, 3500W, 2800 rpm, China). To enhance flavor, 1.5% salt and 1.5% cumin (by weight) were added to the flour for each recipe. The flour formulations included 100% legume flour used individually or mixed in a 1:1 ratio with rice or corn flour. These flour combinations were then used to prepare flatbread.

Preparation of flatbread

Flatbread was prepared using a method similar to traditional Indian flatbreads such as dosai. Dosai is an Indian bread made from a mixture of rice and pulses, cooked on a hot surface (Nagaprabha & Prakash, 2009). The basic dough formula included flour, water, and a leavening agent (Brown, 1993).

To achieve the desired moisture content, warm water was added in a ratio of 220 to 250 of the flour weight for each legume or legume-cereal flour combination. The mixtures were stirred thoroughly to form a semi-liquid paste, which was then divided into equal 50±5g portions. Each portion was baked in a greased pan (10 cm diameter) with sunflower oil for 2–3 minutes on each side, following the method described by Prasad et al. (2012). Additionally, corn flour was mixed with water to prepare corn bread, which served as a control. All flatbread recipes, along with the control corn bread, were subjected to sensory evaluation to determine the most preferred formulations, as presented in Table 2.

Chemical analysis

The chemical composition of the various flatbread samples including moisture, protein, ash, and fat content was determined using the AOAC (2005) methods. Total carbohydrate content was calculated by difference using the following equation: Total available carbohydrates (%) = 100 - (protein% + fat% + moisture% + ash%+ available Fiber %) Energy value of the samples, was calculate according to FAO/WHO/UNU (1985). The equation used for this calculation is: Total valuable calories = 4 (protein %) + 4 (carbohydrates %) + 9 (fat %).

Textural profile analysis (TPA)

The texture profile analysis of flatbread was performed using a texture analyzer (Conetech, Model B, Taiwan) equipped with specialized software, as described by Bourne (2002). The texture parameters measured included hardness, resilience, cohesiveness, gumminess, chewiness, adhesiveness, and springiness.

Color measurement

Color measurements were conducted using a handheld portable colorimeter (CHROMA METER CR-400, Jaflat) according to Matos and Rosell (2013). The instrument was calibrated with a white tile, and color values were expressed in terms of lightness (L^*), redness (a^*), and yellowness (b^*).

Sensory evaluation of flat bread

Flatbread samples were evaluated for their sensory attributes. Half slices of each bread sample were presented to a panel of ten trained assessors on white disposable plates. The panelists assessed the samples based on color and appearance, taste, flavor, body and texture, and overall acceptability. The evaluation was conducted using a nine-point hedonic scale-based scorecard, as described by Srilakshmi (2007).

Amino acid profiles

The amino acid composition of different flatbread recipes was analyzed according to AOAC (2019) using an amino acid analyzer (Biochrom 30). Tryptophan was determined following the method of Hernandes and Bates (1969), using the formula:

Chemical amino acid score (%) = $100 \times [(mg \text{ of amino acid in 1 g test protein})/(mg \text{ of amino acid in reference pattern}) (FAO/WHO, 1991)$

Protein Efficiency Ratio (PER) was calculated using the equation suggested by Alsmeyer et al., (1974).

PER= -0.684+0.456 Leucine - 0.047 Proline

(g/100g protein).

Biological experiment

Two types of diets were tested: Diets containing 100% legume flour and diets containing a 50% legume-cereal flour mixture. The composition of these blends is presented in Table 1-a. The selection of ingredients was based on their protein content, ensuring that each combination met the WHO's recommended nutrient intake for rats.

Management of experimental animals

A biological evaluation of protein quality was conducted over 14 days using 35 adult male albino rats (initial body weight: $75\pm5g$). The experiment was conducted at the rat house of the Regional Center for Food and Feed (RCFF), Agricultural Research Center (ARC), Giza, Egypt. The rats were randomly divided into seven groups (Table 1-a), with five animals per group. Each rat was housed individually in stainless-steel metabolic cages (Model 1000, England; dimensions: $0.35 \times 0.22 \times$ 1.35m) equipped for separate urine and feces collection. The cages were maintained in a clean, wellventilated room at a temperature of $22-24^{\circ}C$ and 45 -55% relative humidity. Each group received a different diet: Control casein diet, cornbread (control recipe), various gluten-free flatbread recipes. A separate group was fed a protein-free basal diet to determine endogenous and metabolic nitrogen loss in feces and urine.

Composition of tested diets

Table 1-b showed that the experimental diets were formulated according to the method of Chapman et al. (1959). The basal diet included: Corn starch (90%), corn oil (5%), mineral mixture (4%) (Hegsted et al., 1941, vitamin mixture (1%) (Campbell, 1963; INC Pharmaceutical, Ohio, USA). The test diets were prepared by incorporating cornbread and different flatbread formulations as a protein source at a 10% level, replacing corn starch in the basal diet. The control diet contained casein as the protein source. After a 9-day acclimatization period, urine and feces were collected over five days. The collected samples were air-dried and weighed. Nitrogen content in urine and feces was determined using the micro-Kjeldahl method (AOAC, 2005). These values were used to calculate the biological value (BV%), true digestibility (TD%), and net protein utilization (NPU) for each tested recipe.

Group	Diets	Diets Composition - Casein basal diet
Control	Basel diet	(Casein)
G1	Lentil / rice (50%)	Rice/lentil flat bread 1:1
G2	Chickpea /rice (50%)	Rice/chickpea flat bread 1:1
G3	Lentil \Corn (50%)	Corn / lentil flat bread 1:1
G4	Chickpea \ Corn (50%)	Corn / chickpea flat bread 1:1
G5	lentil 100%	Lentil flat bread 100 %
G6	Chickpea 100%	Chickpea flat bread 100%

Table 1-a. Different experimental flat bread tested formulas

Parameters measured

The Biological Value (BV %), True Digestibility (TD %), and Net Protein Utilization (NPU %) of the tested flatbread recipes were determined following the method described by Eggum (1973). These parameters were calculated using the following equations:

 $BV = \frac{(mg \ N. \ consumed) - (mg \ N. \ feces - metabolic \ N) - (Urine \ N - 76) \times 100}{mg \ N. \ consumed - (mg \ N \ feces - metabolic \ N)}$ $TD = \frac{mg \ N. \ consumed - (mg \ faces - metabolic \ N) \times 100}{mg \ N. \ consumed - (mg \ faces - metabolic \ N) \times 100}$

mg N. consumed NPV

 $NPU = B V \times TD$

Food Technology Research Journal, Vol. 7, issue 1, 44-61, 2025

···· · · · · · · · · · · · · · · · · ·								
Diets	L/R (50%) 1	Ch/R (50%) 2	L/C (50%) 3	ChC9 (50%) 4	L (100%) 5	Ch (100%) 6	Casein basal diet diet 7	Protein-free basal diet 8
TF (gm)	65.00	70.00	67.70	68.00	40.80	41.70	-	-
Casein (85%)	-	-	-	-	-		11.76	-
Corn starch (%)	25.00	20.00	22.30	22.00	49.20	48.30	78.24	90.00
Minerals mixture (%)	4.00	4.00	4.00	4.0	4.00	4.00	4.00	4.00
*vitamins mixture (%)	1.00	1.00	1.00	1.0	1.00	1.00	1.00	1.00
Corn Oil (%)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00
*Total	100	100	100	100	100	100	100	100
Casein (85%) Corn starch (%) Minerals mixture (%) *vitamins mixture (%) Corn Oil (%) *Total	25.00 4.00 1.00 5.00 100	20.00 4.00 1.00 5.00 100	22.30 4.00 1.00 5.00 100	22.00 4.0 1.0 5.00 100	49.20 4.00 1.00 5.00 100	48.30 4.00 1.00 5.00 100	11.76 78.24 4.00 1.00 5.00 100	90.00 4.00 1.00 5.00 100

Table 1-b. Composition of tested diets

(TF), Tested Flat breads (gm) which provide 10 g protein, * vitamin A (800 IU); Vit. D3 (1,200 IU.); Vit. E (3 IU), Vit. K3-KASTAB (2mg); Vit. B2 - Riboflavin (3mg); Nicotinic acid (10mg); Pantothenic acid (150mg); Manganese (80mg); Zinc (50mg).Copper (2mg); Iodine (1.2mg); Cobalt (0.2mg), Selenium (0.1mg). R: rice, C: corn, L: lentil, Ch: check pea. Total = (TF+ Casein +Corn Starch + +Minerals mixture + vitamins mixture +Corn oil) =100%

Economic Study and cost reduction Analysis

The total cost (L.E.) and cost reduction percentage (%) of different gluten-free flatbread samples were calculated to evaluate the economic feasibility of the formulations. The total cost reduction (%) was determined by comparing the cost of glutenfree flatbread recipes with conventional alternatives.

Statistical analysis

The obtained results were statistically analyzed using a one-way analysis of variance (ANOVA) at a significance level of p < 0.05. Mean values were reported along with their corresponding standard deviations (SD). All statistical analyses were conducted using Assistat, Version 7.7, developed in Brazil, following the methodology proposed by Silva and de Azevedo (2009).

3. Results and Discussion

Sensory evaluation of gluten-free flatbread prepared from different blends

Sensory evaluation is a crucial tool for assessing food acceptability. It plays a significant role in product improvement, quality maintenance, and new product development (Kramer and Twigg, 1970). An exploratory experiment was conducted to evaluate gluten-free flatbread prepared using five different ingredient groups: Rice with legumes, corn with legumes, 100% legumes (individual types), mixed legumes and control (100% corn flour).

Sensory evaluation was used to select the bestperforming recipes, which were then further analyzed for quality properties (Figure 1, Table 2).

Gluten-free flatbread samples were evaluated for their sensory characteristics, and the results are summarized in Table 2.

Group 1 (Rice Blends): The highest-rated samples were rice flour with lentils (R/L) and rice flour with chickpeas (R/Ch), with no significant differences between them. Sensory scores for taste, color, aroma, texture, appearance, and overall acceptability were 8.05 & 8.05, 7.8 & 7.7, 8.4 & 7.65, 8.55 & 8.25, 8.6 & 8.15, and 8.3 & 8.15, respectively. These scores were comparable to those of the control (8.1, 8.45, 7.85, 8.25, 8.25, and 8.1).

Group 2 (Corn Blends): Corn flour with lentils (C/L) and corn flour with chickpeas (C/Ch) also performed well, with no significant differences compared to the control. Their overall acceptability scores were 8.0 and 7.9, respectively, compared to the control score of 8.1.

Group 3 (100% Legumes): Flatbreads made entirely from lentil or chickpea flour received high scores, indicating good sensory acceptance.

Groups 4 & 5 (Mixed Legumes & Control)

Mixed-legume recipes had lower sensory scores compared to the other groups. These findings align with Bojňanská et al. (2012), who reported varying acceptability levels in chickpea- and lentil-floursupplemented breads. Hefnawy et al. (2012) found that incorporating 30% corn flour maintained sensory qualities similar to those of the control. Similarly, Miñarro et al. (2012) observed good sensory acceptance in gluten-free bread containing chickpeas.

Selected samples for further analysis

Based on sensory evaluation, the following six samples were chosen for physical, chemical, and sensory analysis, alongside the control sample: R/L (50% rice & lentil), R/Ch (50% rice & chickpea),

C/L (50% corn & lentil), C/Ch (50% corn & chickpea), L (100% lentil), Ch (100% chickpea) and Control (100% corn). These selected formulations were further studied for their quality attributes, as illustrated in Figure 1.

Table 2. Sensory evaluation (of different flat bread blends
-------------------------------	--------------------------------

Recipes	Taste (9)	Color (9)	Aroma (9)	Body Texture (9)	Appearance (9)	Overall acceptability (9)
Corn bread	$8.15^{a} \pm 0.67$	$8.45^{a}\pm 0.68$	$7.85^{ab} \pm 9.44$	$8.25^{ab} \pm 0.79$	$8.25^{ab} \pm 0.79$	$8.1^{\ { m ab}}\pm 0.77$
Rice bread	$7.45^{abc} \pm 0.68$	$7.6^{\text{ abc}} \pm 0.84$	$7.65^{\text{ ab}}\pm\!0.72$	$7.55^{b} \pm 1.20$	$7.35^{\rm \ bc}\pm 0.88$	7.2 ^b ±0.42
		Gı	oup 1 - rice /Leg	gumes 50%		
L/rice bread	$8.05^{\mathrm{a}} \pm 0.82$	$7.80^{a} \pm 1.14$	$8.45^{a} \pm 0.45$	$8.55^{a} \pm 0.49$	$8.60^{a} \pm 0.51$	$8.30^{a} \pm 0.97$
Ch/rice bread	$8.05^{a} \pm 0.83$	$7.70^{a} \pm 1.14$	7. $75^{ab} \pm 0.75$	$8.25^{ab} \pm 0.79$	$8.15^{ab} \pm 0.88$	$8.15^{ab} \pm 0.88$
P/rice bread	$6.85^{bc} \pm 1.56$	$6.85^{bc} \pm 1.56$	$7.15^{b} \pm 1.20$	$7.4^{ab} \pm 1.24$	$6.05^{cd} \pm 1.30$	$7.25^{b} \pm 1.06$
b/rice bread	$7.34^{b} \pm 1.40$	$7.20^{b} \pm 1.00$	$6.60^{\circ} \pm 1.30$	7.22 ^b ±1.30	$6.69^{\circ} \pm 0.77$	$7.18^{b} \pm 1.80$
		G	roup2- corn /Leg	umes 50%		
L/Corn bread	$7.95^{a} \pm 1.01$	$7.85^{a} \pm 1.15$	$7.65^{ab} \pm 1.06$	$8.05^{ab} \pm 0.93$	$7.90^{abc} \pm 1.22$	$8.00^{ab}\pm 0.82$
Ch/Corn bread	$7.70^{abc} \pm 0.12$	$8.05^{ab}\pm 0.83$	$7.80^{\rm ab} \pm 1.05$	$7.85^{ab} \pm 1.03$	$7.95^{ m abc} \pm 0.76$	$7.95^{ab} \pm 0.76$
P/Corn bread	$7.15^{b} \pm 0.99$	$7.18^{b} \pm 0.75$	$7.17^{b} \pm 1.03$	7. $25^{ab} \pm 1.03$	$7.27^{b} \pm 1.06$	$7.28^{b}\pm0.88$
B/Corn bread	$7.26^{b}\pm0.87$	$7.25^{b} \pm 0.74$	$7.15^{b} \pm 1.15$	7. $25^{ab} \pm 1.06$	7. $25^{b} \pm 0.09$	$7.35^b\pm\!0.94$
			Group3 - Legun	nes 100%		
L bread	$7.40^{abc} \pm 1.20$	$7.15^{ab}\pm 1.37$	$7.25^{bc} \pm 1.27$	$7.55^{ab}\pm\!1.21$	$7.05^{bc} \pm 1.26$	$8.30^{a} \pm 1.30$
Ch bread	$7.35^{abc} \pm 0.90$	$7.00^{abc} \pm 0.84$	$7.35^{b} \pm 1.25$	$7.25^{bc} \pm 1.25$	$7.00^{\rm \ bc} \pm 0.97$	$8.15^{ab}\pm\!0.70$
Pea bread	$6.85^{bc} \pm 1.56$	$7.70^{bc} \pm 0.81$	7.1 ^b ±51.20	$7.40^{ab}\pm1.24$	$7.05^{\rm \ bc} \pm 1.30$	7.25 ^b ±1.06
Bean bread	$6.70^{\circ}\pm0.79$	$6.90^{b} \pm 0.77$	$7.05^{b} \pm .065$	$7.35^{b}\pm1.45$	$6.45^{\circ} \pm 1.06$	$7.2^{b} \pm 0.95$
			Group 4	- mix legumes 50	1%	
L\Ch bread	$7.15^{b} \pm 1.05$	7.65 ^{ab} ±1.05	7.85 ^{ab} ±1.05	7.90 ^{ab} ±1.12	$7.40^{\rm bc} \pm 1.37$	6.25°±1.05
L/P bread	7.25 ^b ±1.37	$7.45^{\text{ abc}} \pm 1.14$	$7.45^{b}\pm 0.89$	$7.80^{ab} \pm 0.78$	$7.50^{ m bc}\pm 1.05$	$6.28^{\circ} \pm 1.00$
L/b bread	$6.90^{bc} \pm 1.02$	$6.90^{bc} \pm 1.25$	$7.30^{b} \pm 1.31$	$7.55^{ab} \pm 1.12$	$6.95^{\circ} \pm 1.04$	$6.15^{\rm c}\pm0.97$
Ch/p bread	7.30 ^b ±1.20	7.25 ^b ±1.27	7.30 ^b ±1.43	$7.45^{ab} \pm 1.10$	$7.30^{bc} \pm 1.40$	6. $55^{bc} \pm 0.97$
Ch/b bread	$6.90^{bc} \pm 1.02$	$7.05^{bc} \pm 1.03$	$7.30^{b} \pm 1.22$	$7.30^{b} \pm 1.35$	$7.20^{b} \pm 1.21$	$6.25^{\circ} \pm 1.09$
P/b bread	6.75 °±1.16	$7.05^{bc} \pm 1.403$	7.45 ^b ±1.27	$7.65^{ab} \pm 1.02$	$7.20^{b} \pm 1.05$	$6.15^{\circ} \pm 0.94$

Control: corn 100%, R: rice, C: corn, p: Pea, B: bean L: lentil, Ch.: check pea

*Values are means of three replicates \pm SD. Each mean value, within the same raw, followed by the same letters is not significantly different at 0.05 level.



Corn bread 100%, Ch100%, L100%, Ch/rice50%, L/rice 50%, Ch/corn50%, L/Corn 50%

Figure 1. Different flat bread samples after sensory evaluation

Chemical composition of gluten-Free flatbread prepared from different blends

The chemical composition of gluten-free flatbread made from various blends was analyzed and is presented in Table 3. All recipes exhibited a significant increase in protein content, ranging from 13.7% to 24.5%, compared to the control flatbread made from corn flour (7.08%). The highest protein content was found in flatbreads made from 100% lentil and 100% chickpea flour, containing 24.5% and 23.96% protein, respectively. This increase can be attributed to the naturally higher protein levels in chickpeas and lentils. In terms of Dietary Reference Intake (DRI) for protein, each 100 g of dried flatbread provided 27.5%, 25.5%, 26.6%, 26.25%, 43.8%, and 42.8% of the daily protein requirement for a 19 to 50 year-old male weighing 65kg (based on a recommended intake of 56g of protein per day) for the following blends: Rice/lentil (50%), rice/ chickpea (50%), corn/Lentil (50%), corn/Chickpea (50%), 100% Lentil and 100% Chickpea. These findings align with Samah (2004), who reported that combining cereals and legumes enhances both protein content and protein quality in cereal-based complementary foods.

Fat content: Flatbreads made from 100% chickpea or mixed with corn or rice had higher fat content than other samples, likely due to the naturally higher fat content of chickpeas.

Ash content: A significant increase in ash content was observed in corn/chickpea (2.8%) and 100% chickpea (2.65%) samples, compared to other blends and the control.

Carbohydrate content: All gluten-free flatbread samples showed a notable decrease in carbohydrate content compared to the control (85.52%). The greatest carbohydrate reductions were observed in 100% lentil (61.18%) and 100% chickpea (61.36%) flatbreads. This decrease is attributed to the increase in protein content, which naturally lowers the proportion of carbohydrates.

Caloric Value

The control sample (100% corn flour flatbread) had the highest caloric value at 403.97Kcal. All oth-

er samples showed a significant caloric reduction, making them more suitable for dietary considerations. Overall caloric values ranged from 348.66 to 403.97Kcal. These findings are consistent with Gonzales et al. (2016), who stated that chickpea-based baked food products are good sources of nutrients, particularly energy, containing 105 to 526Kcal.

Color characteristics of gluten free flat bread prepared from different blends.

Color is a crucial characteristic influencing the acceptability of food products, as noted by Nithya et al. (2016). It is also one of the most important quality attributes of flatbread, as emphasized by Mahmoud et al. (2013). This study evaluates the color parameters lightness (L^*) , redness (a^*) , and yellowness (b^*) of gluten-free flatbreads, as presented in Table 4. The results demonstrated significant differences in color attributes across all flatbread samples compared to the control. Notably, flatbreads made from 100% chickpea or 100% lentil exhibited lower L^* values relative to the control, while their a^* and b^* values were higher, indicating increased redness and yellowness. Specifically, increasing the proportion of chickpea or lentil flour significantly reduced L^* values (54.43 for chickpea and 51.13 for lentil), producing darker bread. Concurrently, a^* values rose to 8.06 and 7.08, and b^* values increased to 36.32 and 35.83 for chickpea and lentil flatbreads, respectively, reflecting greater redness and yellowness. In contrast, the control sample displayed the highest L^* value (70.2) alongside significantly lower a^* and b^* values (1.81 and 32.69), indicating reduced redness and yellowness. These findings align with Olojede et al. (2020), who reported that adding chickpea flour to bread formulations darkens the product due to enhanced Maillard reactions, driven by lysine and sugars in chickpeas. Additionally, Sharima-Abdullah et al. (2018) observed that incorporating vegetable proteins and chickpea flour improves color and visual appeal, underscoring the importance of optimizing ingredient proportions to boost consumer acceptance.

Evaluation of Gluten-Free Flatbread Substituted with Some Cereals and Legumes

Recipes	Protein (%)	Fat (%)	Fiber (%)	Ash (%)	Total Carbohydrates (%)	Caloric Value (Kcal/100g)
Control	$07.08^{\rm c}\pm6.56$	3.73 ± 0.11	$02.15^{e}\pm 0.33$	$1.52^{cd} \pm 0.22$	$85.52^{a}\pm0.20$	$403.97^{\text{a}}{\pm}~2.00$
L100%	$24.50^{a} \pm 0.51$	$0.66^{d}\pm0.07$	$12.30^{\mathrm{a}} \pm 0.20$	$1.36^{\text{d}}\pm0.01$	$61.18^{e}\pm0.20$	$348.66^{\mathrm{f}}{\pm}~0.34$
Ch 100%	$23.96 {}^{a} \pm 0.40$	$4.83\ ^a\pm 0.72$	$07.20^{d} \pm 0.20$	$2.65^a\pm~0.13$	$61.36^{e}\pm0.30$	$384.75^{b} \pm 0.52$
L/rice (50%)	1 5.40 ^b ±0.36	$2.26^{\ c}\pm0.20$	$07.90^{c}\pm0.10$	$1.60^{\circ}\pm0.15$	$72.84^{c}\pm0.23$	$373.30^{d} \pm 1.82$
Ch/rice(50%)	$14.30^{0b} \pm 0.80$	$4.23^{ab}\pm 0.25$	$08.02^{\text{c}} \pm 0.61$	$2.26^{b}\pm 0.20$	71.19 ^c ±0.23	$380.03^{\text{c}}{\pm}\ 0.01$
L/Corn (50%)	14. 90 ^b ±0.60	$2.50^{\ c}\pm0.43$	$09.20^{\text{b}} \pm 0.20$	$2.46^{ab} \pm 0.05$	71.34 ^c ±0.22	$365.86^{e} \pm 0.55$
Ch/Corn (50%)	$1 4.70^{b} \pm 0.62$	$4.50^{a}\pm 0.70$	$10.00^{b}\pm 0.70$	$2.80^{a}\pm1.50$	$67.80^{d} \pm 0.28$	$371.30^{d} \pm 0.30$

Fable 3. Chemica	l composition a	of different fl	at bread recip	es (on dry	weight g/100g)
-------------------------	-----------------	-----------------	----------------	------------	----------------

Table 4. Color characteristics of gluten free flat bread prepared from different blends

Recipes	<i>L</i> *	a*	b*
Corn (100%)	$70.12^{a}\pm0.12$	$1.81^{e}\pm 0.81$	$32.69^{d} \pm 0.09$
Lentil (100%)	51.13 ^g ±0.13	$8.06^{a}\pm0.06$	$36.32^{a} \pm 0.32$
Chickpea (100%)	$54.43^{f} \pm 0.44$	$7.08^{b}\pm0.08$	$35.83^{b} \pm 0.03$
L/rice (50%)	$66.32^{\circ} \pm 0.32$	$5.36^{d} \pm 0.36$	$32.16^{e} \pm 0.16$
Ch/ rice (50%)	$67.47^{b}\pm0.47$	$5.56^{cd} \pm 0.11$	$31.08^{\rm f}{\pm}~0.08$
L/Corn (50%)	$63.9^{e} \pm 0.9$	$7.00^{b}\pm0.10$	$34.32^{c} \pm 0.32$
Ch/Corn (50%)	$65.32^{d}\pm0.32$	$6.07^{c}\pm0.07$	$32.83^{d} \pm 0.03$

Control: corn bread 100%, R: rice, C: corn, L: lentil, Cch : check pea

Values are means of three replicates \pm SD. Each mean value, within the same raw, followed by the same letters is not significantly different at 0.05 level. $L^{\underline{}}$ lightness color score are $a^{\underline{*}}$ = redness color score $b^{\underline{*}}$ = yellowness color score.

Texture profile analysis of gluten free flat bread prepared from different blends.

Hardness is a critical textural property in evaluating baked goods due to its direct association with the human perception of freshness (Karaoğlu & Kotancılar, 2009). The texture results of gluten-free flatbreads prepared from different recipes are shown in Table 5. Hardness varied significantly across all gluten-free flatbread samples, ranging from 4.3 to 11.15N. The highest hardness value (11.15±0.75N) was observed in the sample containing 100% corn flour, followed by the C/Ch 50% blend (corn/ chickpea at 7.62 ± 0.3 N), while the lowest hardness (4.3±0.26N) was recorded in the 100% chickpea sample. The results clearly demonstrate that incorporating chickpea or lentil flour significantly reduced hardness while increasing springiness and resilience. Conversely, adding corn or rice flour to blends significantly increased hardness and decreased springiness and resilience. Notably, the highest springiness (2.68±0.08mm) and resilience (0.16 ± 0.02) values were observed in the 100% chickpea sample, chickpea sample, which also improved quality parameters such as porosity, specific volume, and crust-to-crumb ratio. Texture and SEM analyses revealed that increasing chickpea flour concentration decreased crumb firmness and produced a continuous, smoother texture (Mohammad et al., 2014). This may be attributed to chickpea's high protein content, which promotes a more porous structure and softer crumb. Protein content in flour is a key factor influencing hardening and staling rates (Pateras et al., 2007). Table 5 also indicates that chewiness and gumminess values decreased as hardness diminished, following a similar trend to hardness. These findings align with Ibrahim (2011), who reported that gumminess and chewiness are hardness-dependent parameters. Chewiness and adhesiveness are texture attributes strongly correlated with sensory evaluations conducted by trained panels (Esteller et al., 2004). The use of chickpea as an ingredient in gluten-free foods has expanded due to its nutritional value and its ability to enhance product quality (Capriles & Arêas, 2014).

Recipes	Hardness (N)	Adhesive- ness (mj)	Resilience	Cohesiveness	Gumminess	Springiness (mm)	Chewiness
Control	$11.15^{a}\pm0.75$	5.6 ^a ±0.55	$0.03^{e}\pm 0.02$	$1.10^{e}\pm0.10$	$12.39^{a}\pm0.10$	$0.37 {}^{\rm a}\!\pm\!0.78$	$56.4^{a}\pm0.20$
L 100%	$5.74^{d}\pm 0.22$	$2.30^{cd} {\pm} 0.10$	$0.08^{\text{cde}} \pm 0.11$	$2.16^{b} \pm 0.05$	$12.28^{a}\pm0.07$	$2.58^{b}\pm0.50$	$33.90^{\circ} \pm 1.00$
Ch 100%	4.30 ^e ±0.26	$2.70^{b}\pm0.10$	$0.16^{bc} \pm 0.02$	$0.32^{\rm f}\pm0.21$	$14.40^{a}\pm0.22$	$2.68^d \pm 0.08$	$39.14^{b}\pm0.79$
L/rice (50%)	$6.02^{d}\pm 0.13$	$0.90^{e} \pm 0.01$	$0.05^{\text{de}} \pm 0.03$	$1.60^{c}\pm0.14$	9.90°±0.10	$2.60^{b} \pm 0.56$	$29.30^d\!\pm\!0.80$
Ch/rice (50%)	$7.07^{b}\pm 0.23$	$1.23^{e}\pm 0.15$	$0.02^{e}\pm 0.05$	$1.26^{d}\pm0.07$	$9.50^{d} \pm 0.10$	$2.55^{bc} \pm 0.16$	$21.50^{e}\pm 1.00$
L/Corn (50%)	$5.80^{d} \pm 0.90$	$2.30^{cd} \pm 0.10$	$0.05^{de} \pm 0.04$	1.75°±0.05	$10.48^{b}\pm 0.20$	$2.54^{b}\pm0.47$	$33.90^{\circ}\pm0.20$
Ch/Corn (50%)	7.62 ^b ±0.31	$2.60^{bc} \pm 0.10$	$0.11^{cd} \pm 0.05$	$1.24^{de} \pm 0.05$	$9.78^{cd} \pm 0.20$	$2.60^b{\pm}0.30$	$29.80^d{\pm}0.25$

Table 5. Texture profile analysis of gluten free flat bread prepared from different blends

Control: corn 100%, R: rice, C: corn L: lentil, Ch.: check pea.*Values are means of three replicates \pm SD. Each mean value, within the same raw, followed by the same letters is not significantly different at 0.05 level.

Desines		Hardness (Figure 2a))	
Recipes	Zero	24 h	48 h	72 h
Corn 100%	11.3 ^a ±1.13	$21.07^{a} \pm 0.07$	$41.04^{a}\pm0.07$	$76.90^{a} \pm 0.9$
L 100%	$5.07^{cd} \pm 0.07$	$6.99^{e} \pm 0.99$	$12.21^{f}\pm0.21$	$25.30^{g}\pm0.3$
Ch 100%	4.3 ^b ±0.20	$8.30^{d} \pm 0.30$	$15.4^{e}\pm0.40$	$29.00^{f} \pm 0.10$
L/rice 50%	5.6°±0.52	$13.08^{\circ} \pm 0.08$	$22.07^{d}\pm0.07$	$40.70^{e}\pm0.72$
Ch/rice50%	$7.07^{d}\pm0.07$	$17.07^{b} \pm 0.07$	$22.07^{d}\pm0.07$	$43.60^{d} \pm 0.5$
L/Corn 50%	$5.58^{\circ}\pm0.16$	$17.07^{b}\pm0.07$	$24.06^{b}\pm0.06$	$46.50^{\circ}\pm0.52$
Ch/Corn50%	$7.9^{d}\pm0.60$	$17.65^{b}\pm0.65$	$24.80^{b}\pm0.80$	$50.50^{b}\pm0.51$
		Stalling (Figure 2b)		
Corn 100%	$284.40^{a} \pm 21.1$	$392.80^{a}\pm0.80$	$484.7^{a}\pm0.72$	$598.50^{a} \pm 0.51$
L 100%	$193.50^{d} \pm 0.52$	$276.60^{d} \pm 0.60$	$334.6^{f} \pm 0.61$	$417.90^{e} \pm 0.90$
Ch 100%	273.57°±0.56	$276.60^{d} \pm 0.60$	334.6 ^f ±0.61	$420.60^{e} \pm 0.06$
L/rice 50%	$257.57^{bc} \pm 0.56$	$316.80^{\circ} \pm 0.80$	$341.4^{e}\pm0.40$	$452.90^{d} \pm 0.59$
Ch/rice50%	$284.50^{a}\pm0.50$	$355.50^{b}\pm0.50$	$408.2^{\circ}\pm0.30$	$510.70^{\circ} \pm 0.07$
L/Corn 50%	$273.60^{ab} \pm 0.55$	$318.40^{\circ}\pm0.42$	$348.3^{d}\pm0.32$	$456.20^{d}\pm0.52$
Ch/Corn50%	$288.57^{a} \pm 0.56$	$342.00^{bc} \pm 0.01$	$418.4^{b}\pm0.40$	$525.60^{b} \pm 0.60$

Control: corn bread 100%, R: rice, C :corn, L: lentil, C/ch: check pea., *Values are means of three replicates \pm SD. Each mean value, within the same raw, followed by the same letters is not significantly different at 0.05 level

Effect of Storage Time (hrs.) on Hardness & Staling of Different Flatbreads

Staling is a complex process influenced by multiple factors, including amylopectin retrogradation, reorganization of polymers in the amorphous region, moisture loss, redistribution of water between amorphous and crystalline zones, and changes in crumb macroscopic structure (Davidou et al., 1996; Rojas et al., 1999). A gradual decline in freshness (increased staling) was observed in all flatbreads over time. This may result from amylose crystallization post-baking or the inherent properties of lentil flour, which may delay staling. These findings align with Seleem (2000), Ammar et al. (2020), and Matsushita et al. (2020), who reported similar mechanisms in starch-based products. A negative correlation was observed between storage time and freshness, consistent with studies on tortillas by El-Tawil (1998) and Khorshid et al. (1997), who noted that (hardness) increased progressively during storage.



Figure 2a. Effect of storage time (hrs.) on hardness of different flat bread



Figure 2b. Effect of storage time (hrs.) on staling of different flat bread

Effect of storage time on hardness and staling

Figures 2a and 2b illustrate the hardness and staling trends of flatbread blends over storage periods of 0, 24, 48, and 72hours. At the initial timepoint (0 hours), the 100% chickpea and 100% lentil blends exhibited the lowest hardness and staling values, followed by rice-legume (RL, RCh) and corn-legume (CL, CCh) blends. In contrast, the control (corn flour bread) displayed the highest hardness and staling values. Both parameters increased progressively during storage, with the most pronounced rise observed in corn and corn-legume blends. Sidhu et al. (1997) reported that freshness, a holistic attribute encompassing taste, aroma, and texture is highly valued in flatbread. However, due to its lean formulation, flatbread is prone to rapid staling during storage, leading to a limited shelf life Staling involves physicochemical changes such as moisture redistribution and starch retrogradation, resulting in increased firmness and altered organoleptic properties. These changes render the bread tougher and less palatable, often deemed unacceptable by consumers (Gujral & Pathak, 2002). A direct correlation exists between prolonged storage time, increased firmness, and reduced freshness.

Amino acid profile and nutritional quality

The amino acid composition of various flatbread formulations (Table 7) revealed glutamic acid as the most abundant amino acid, with concentrations ranging from 16.1 to 18.44g/100g protein across 100% chickpea, 100% lentil, legume-cereal blends (50% chickpea/rice, 50% lentil/rice, 50% chickpea/corn, 50% lentil/corn), and the control (corn flatbread at 17.27g/100g protein). Aspartic acid ranked second, decreasing progressively from 11.61g (100% chickpea) to 8.42g (50% lentil/corn), while corn-containing blends exhibited leucine as the third predominant amino acid. Methionine and tryptophan were the least abundant, with methionine ranging from 0.9g (50% chickpea/rice) to 2.64 g (100% chickpea) and tryptophan varying between 0.65g (100% chickpea) and 1.2g (50% lentil/rice) per 100g protein.

ead recipe
ifferent br
n of d
protei
g/100g)
content (
acids
Amino
Table 7.

Amino acid	Corn 100%	Lentil 100%	Ch100%	L /rice %	Chick /rice	L/corn	Ch /corn
Isoleucine	$2.70^{i\pm0.10}$	$4.20^{h\pm} 0.11$	$03.80^{i}\pm0.13$	$4.65^8 \pm 0.05$	$4.45^{B\pm0.14}$	$3.75^{\mathrm{h}}\pm0.05$	$3.55^{ m h\pm}0.05$
Leucine	$10.10^{b}\pm0.10$	$7.10^{\rm d} \pm 0.13$	$07.20^{d}\pm0.10$	$7.70^{\circ}\pm0.023$	$8.60^\circ\pm0.01$	$8.60^{b}\pm0.10$	$8.65^{b}\pm0.05$
Valine	$4.72^{g}\pm 0.02$	$4.60^{1\pm} 0.15$	$4.20^8{\pm}0.12$	$4.85^{\mathrm{f}\pm0.05}$	$4.60^{\rm f}\pm0.19$	$4.66^{\rm f}\pm0.11$	$4.46^{\rm f}\pm0.02$
Threonine	$2.84^{i}\pm0.04$	$3.76^{k}\pm0.02$	$3.70^{\mathrm{i}}\pm0.16$	$3.70^{i}\pm0.10$	$3.60^{\mathrm{h}}\pm0.02$	$3.80^{\rm h}{\pm}0.10$	$3.77^{\rm g}\pm0.11$
Phenyl alanine	$5.06^{\rm f}\pm0.06$	$5.00^{\rm f} \pm 0.20$	$5.30^{\circ}\pm0.01$	$5.02^8 \pm 0.10$	$5.18^{\circ}\pm0.01$	$5.02^{\circ}\pm0.02$	$5.17^{\circ}\pm0.17$
Methionine	$2.64^{j}\pm0.04$	$0.90^{\circ}\pm0.10$	$1.20^{\rm m}\pm0.01$	$2.32^{\rm k}\pm 0.20$	$2.47^{i}\pm0.02$	$1.75^{j}\pm0.05$	$1.90^{~j}\pm0.10$
Tryptophan	$0.65^{m}\pm0.02$	$0.80^{p}\pm 0.11$	$0.90^{n}\pm0.10$	$1.15^{\rm m}\pm 0.01$	$1.20^{1}\pm0.05$	$0.71^{1}\pm0.01$	$0.81^k\pm0.05$
Lysine	$2.80^{i}\pm0.04$	$6.70^{\circ\pm}$ 0.15	$6.50^{\rm de}\pm0.18$	$4.80^{\rm f}\pm0.04$	$4.70^{\rm f}\pm0.01$	$4.70^{\rm f}{\pm}0.02$	$4.62^{\rm f}\pm0.02$
Histidine	$2.40^{\mathrm{k}}\pm0.10$	$2.60^{1}\pm0.12$	$2.80^{i\pm0.11}$	$2.60^{j}\pm0.07$	$2.50^{j}\pm0.03$	$2.50^i\pm0.10$	$2.62^i\pm0.05$
Serine	$2.40^{\mathrm{k}}\pm0.10$	$4.80^{i}\pm0.16$	$4.70^{\rm f}{\pm}0.01$	$5.09^{\circ}\pm0.07$	$5.04^{i}\pm0.05$	$4.04^{g}{\pm}0.04$	$3.75^{g}\pm0.05$
Cysteine	$2.08^{1}\pm0.01$	$1.10^{n}\pm0.02$	$1.50^{1}\pm0.11$	$1.30^{1}{\pm}0.04$	$2.28^{\mathrm{k}}\pm0.20$	$1.30^k{\pm}0.01$	$1.79^{j}\pm 0.01$
Tyrosine	$2.86^{i}\pm0.02$	$2.50^{\rm m}\pm0.11$	$2.20^{k}\pm0.21$	$2.71^{j}\pm0.03$	$2.60^i\pm0.17$	$2.68^{i}\pm0.10$	$2.53^{i}\pm0.03$
Aspartic acid	$6.08^{d}\pm0.04$	$10.70^{b} \pm 0.20$	$11.61^{b}\pm 0.01$	$8.94^{b}\pm0.04$	9.42 ^b ±0.40	$8.42^{b}\pm0.02$	$8.80^{\rm b}\pm0.16$
Glutamic acid	$18.44^{a}\pm0.50$	$16.10^{a}\pm0.11$	$16.60^{a}{\pm}0.20$	$17.50^{a}{\pm}0.02$	$17.17^{a}\pm 0.11$	$17.27^{a}\pm0.19$	$16.92^{a}\pm0.02$
Proline	8.34°±0.10	$03.90^{\text{j}\pm0.15}$	$4.40^8 \pm 0.21$	$4.30^{ m h}{\pm}0.01$	$4.70^{\rm f}\pm 0.10$	$6.17^{d}\pm0.12$	$6.37^{d}\pm0.03$
Glycine	$3.62^{\rm h}{\pm}0.20$	$04.00^{i}\pm0.17$	$3.60^{i}\pm0.10$	$3.64^{i}\pm0.11$	$3.18^{\rm h}\pm0.10$	$3.80^{\mathrm{h}}\pm0.10$	$3.70^{g}\pm0.02$
Alanine	$6.04^{d}\pm0.04$ ($04.20^{\mathrm{h}}\pm0.22$	$4.10^{\rm h}{\pm}0.12$	$5.17^{\circ}\pm0.09$	$5.02^{e}\pm0.33$	$5.12^{i}\pm0.13$	$5.04~^{\mathrm{ef}}\pm0.21$
Arginine	$5.26^{\circ}\pm0.20$	$07.60^{\circ}\pm0.22$	$9.00^0 \pm 0.16$	$7.10^d \pm 0.05$	$7.48^{\rm d}\pm0.15$	$6.43^{\rm c}\pm 0.03$	$7.50^{\circ} \pm 0.05$
Control: corn bread 100% same letters is not significar	, R: rice, C :Corn, ttly different at 0.05 1	L: lentil, Ch: chi level.	ckpea. *Values are mea	ns of three replicates $\pm S$	D. Each mean value,	within the same raw	7, followed by the

Evaluation of Gluten-Free Flatbread Substituted with Some Cereals and Legumes

Food Technology Research Journal, Vol. 7, issue 1, 44-61, 2025



Figure 3. Amino acid composition in different flat bread recipes

Figure 3 shows the Total Essential Amino Acids (TEAA), Total Non-Essential Amino Acids (TNEAA), and Total Amino Acids (TAA) of different flatbread recipes. TEAA values ranged from 33.91% to 37.3%, with the highest value observed in the 50% chickpea/rice blend and the lowest in the control (corn bread). TNEAA values varied between 55.12% and 57.7%, where the 100% chickpea recipe exhibited the highest TNEAA (57.7%), followed closely by the 50% chickpea/rice

(56.89%) and 50% chickpea/corn (56.4%) blends. The 100% lentil (L100) recipe recorded 54.9%, while lentil/corn and lentil/rice blends showed slightly lower values. For total amino acids (TAA), the 50% chickpea/rice blend had the highest value, followed by the 100% chickpea recipe, whereas the control corn bread had the lowest. Overall, formulations containing chickpea (either alone or blended with rice/corn) demonstrated superior TAA and TEAA values compared to other recipes.

Table 8.	Chemical score	and limiting	amino acids	of different	formulas o	f flat bread
----------	-----------------------	--------------	-------------	--------------	------------	--------------

Amino acid	Corn 100%	Lentil 100%	Ch 100%	L/rice 50%	Ch /rice (50%)	L/Corn (50%)	Ch/Corn (50%)	FAO/WHO, 1973
Threonine	71.00	94.00	92.50	92.50	90	95.00	94.25	4.00
Valine	94.40	92.00	84.00	97.00	92	93.20	89.20	5.00
Methionine	120.00*	40.90^{a}	54.50^{a}	105.50*	112.2*	79.50	86.30	2.20
Isoleucine	67.50	105.00*	95.00	116.25*	111.25*	93.75	88.75	4.00
Leucine	144.20*	101.40*	102.80*	110.00*	122.8*	123.00*	123.5*	7.00
Phenylalanine	180.70*	178.50*	189.30*	179.00*	185*	179.28*	184.6*	2.80
Lysine	51.50 ^a	123.00*	119.50*	88.00^{a}	86.4 ^a	86.40	85.00	5.44
Tryptophan	67.70	83.30	93.75	119.79*	125*	73.90 ^a	84.30 ^a	0.96
PER *	3.5#	2.37	2.39	2.60#	3.0#	2.9#	2.96#	Standard value

Composition supplies 100% or more of the requirement. a, first limiting amino acid.. PER= - 0.684+0.456 Leucine - 0.047 Proline(g/100g protein. # PER values (a standard value of casein protein). Chemical amino acid score % = $100 \times [(mg \text{ of amino acid in 1 g test protein}) / (mg \text{ of amino acid in reference pattern}).$

Table 8 presents the Chemical Score (CS), Protein Efficiency Ratio (PER), and limiting amino acids of various flatbread formulations. The data reveal that lysine is the limiting amino acid in corn bread, with a CS of 51.5%, followed by tryptophan (67.7%). In contrast, methionine serves as the limiting amino

acid in chickpea and lentil bread recipes, with CS values of 54.5% (chickpea) and 40.9% (lentil). Generally, lysine is the primary limiting amino acid in legume-cereal flatbread blends, while corn bread is rich in methionine, leucine, and phenylalanine—all exceeding 100% of dietary requirements.

Notably, all bread formulations met or surpassed 100% of phenylalanine requirements, and rice-based recipes exceeded recommended levels of lysine and tryptophan. Cereal proteins typically contain only 2% lysine, less than half the concentration recommended by the FAO/WHO (1973) for human nutrition. The Protein Efficiency Ratio (PER), a key indicator of protein quality, ranged from 2.37 to 3.0 across gluten-free flatbreads. All recipes except the 100% lentil (2.37) and 100% chickpea (2.39) exceeded the standard casein reference value (2.5), indicating high protein quality. The elevated PER values in blended recipes correlate with higher leucine concentrations (Table 8). According to Oluwaniyi et al. (2010) and Zengin et al. (2012), foods with PER values >2.0 are classified as high-quality protein sources, while those >2.5 (the casein benchmark) are considered excellent.

Experimental diets	BV %*	TD %**	NPU %***
Casine diet (control)	$84.50^{a} \pm 3.50$	$97.25^{a} \pm 0.50$	$82.17^{a} \pm 3.14$
L / rice 50%	$71.45^{cd} \pm 5.50$	$75.30^{bcd} \pm 0.65$	$53.80^{\text{cde}}\pm7.01$
Ch / rice50%	$70.10^{d} \pm 6.80$	$74.31^{bcd} \pm 0.67$	$52.09^{d} \pm 6.00$
L /Corn 50%	$74.57^{bcd} \pm 5.10$	$78.30^{\rm \ bcd}\pm0.80$	$58.38^{\circ} \pm 0.82$
Ch/Corn 50%	$70.90^{d} \pm 5.50$	$69.71^{bcd} \pm 0.50$	$49.42^{cd} \pm 0.16$
L100%	$79.50^{b} \pm 266$	$83.75^b \pm 0.70$	$66.60^{b} \pm 0.69$
Ch 100 %	$76.50^{bc} \pm 4.30$	$79.07^{\mathrm{bc}} \pm \ 0.40$	$60.50^{ m \ bc}\ \pm 0.57$

R: rice, C :corn, L: lentil, Cch: check pea .*Values are means of three replicates ±SD. Each mean value, within the same raw, followed by the same letters is not significantly different at 0.05 level. BV* biological value, (TD)** True digestibility (NPU) ***Net protein utilization

Protein Quality of Experimental Diets

The protein quality parameters true digestibility (TD), biological value (BV), and net protein utilization (NPU) of different flatbread formulations are presented in Table 9. NPU values differed significantly between all flatbread samples and the casein control. The casein diet yielded the highest NPU (82.2%), followed by the 100% lentil (L100%) recipe (66.6%), which surpassed the 100% chickpea (Ch100%) recipe (60.5%). All NPU values exceeded the 60% threshold recommended by PAG (1971) for legume-based blends. Biological value (BV) ranged from 70.1% to 79.5% across recipes. The 100% lentil (L100%) flatbread showed the highest BV (79.5%), followed by the 100% chickpea (Ch100%) recipe (76.5%), while the lowest values were observed in the 50% chickpea/rice (70.1%) and 50% chickpea/corn (70.9%) blends. These results align with Oser (1959), who classifies proteins with a BV of 70-100% as nutritionally adequate. For true digestibility (TD), the casein control (97.3%), L100% (83.75%), and Ch100% (79.07%) recipes exhibited the highest values. No significant differences were observed among the remaining samples: 50% lentil/rice (75.3%), 50% chickpea/rice (74.31%), 50% lentil/corn (78.3%), and 50% chickpea/corn (69.7%).

	()		()	8		1
Recipes	Cost of 100g sample (L.E)	20%	Total cost (L.E)	No. of .of Loaves	Cost of Loave (40-50g) (L.E)	Cost reduction (%)
Corn flour	8.5	1.70	10.20	5	2.04	
L100%	6.0	1.20	7.20	6	1.20	- 41.18
Ch 100%	8.0	1.60	9.60	6	1.60	- 21.56
R\L 50%	4.5	0.90	5.40	5	1.08	- 47.05
R\Ch 50%	5.5	1.10	6.60	5	1.32	- 35.30
C\L 50%	7.25	1.45	8.70	5.5	1.58	- 22.55
C\Ch50%	8.25	1.65	9.90	5.5	1.80	- 11.80

Table 10. Total cost (L.E) & cost reduction (%) of different gluten-free flat bread samples

R: rice, C: corn, L: lentil, Cch: check pea

. .

Production Cost Analysis

The production costs of gluten-free flatbread samples are detailed in Table 10. Results indicate that the total cost ranged from 5.40 to 9.90 pounds/100g, compared to the control bread (10.20 pounds/100g). Cost reductions for the formulated blends varied between 11.80% and 47.05%. The lentil-rice flatbread achieved the highest cost reduction (47.05%), followed by the 100% lentil flatbread (41.18%), while the chickpea-corn blend showed the lowest reduction (11.80%). Other samples exhibited cost reductions ranging from 21.56% to 35.30% relative to the control. Overall, all experimental blends were more cost-effective than the control formulation.

4. Conclusion

In response to the growing demand for functional gluten-free baked goods, this study demonstrates that lentil or chickpea flour-either used individually (100%) or blended with corn or rice flour-can successfully produce gluten-free flatbread with acceptable physicochemical properties. The findings reveal that lentil and chickpea flours are superior to yellow corn flour in terms of protein, fiber, and fat content, yielding final products enriched with crude protein, crude fiber, ash, and ether extract compared to the control. The formulated flatbreads are not only gluten-free but also costeffective, achieving production cost reductions of 11.80% to 47.05% relative to the control (corn bread). Notably, lentil-rice blends showed the highest cost efficiency. These results highlight the feasibility of preparing high-quality gluten-free flatbread using raw materials such as yellow corn, chickpea, rice, and lentil flours, offering a nutritious and affordable option suitable for individuals with celiac disease. Gluten-free bread made from lentils, chickpeas, or blended with corn/rice is recommended as a safe, nutritious alternative for individuals with wheat gluten allergies. Unlike traditional wheat bread, which can harm digestion and growth, this option provides high-quality protein, easy digestibility, and appealing taste, texture, and color. Compared to conventional corn bread, it serves as a complete meal, is cost-effective, and requires mini-

mal preparation time.

References

- AOAC (2005) Association of Official Analytical Chemist. Official Methods of Analysis. 18th Edition, Washington DC.
- AOAC (2019). Official Methods of Analysis (OMA). Association of Official Analytical Chemists. Official Methods of Analysis. AOAC, Washington, DC, USA. https://www.aoac.org/official-methods-of-analysis-21st-edition-2019/
- Akubor, P.I., Adamolekun F.O., Oba C.A., Obari H. and Abudu, I.O. (2003) Chemical composition and functional properties of cowpea and plantain flour blends for cookie production. Plant Foods Hum Nutr 58:1-9.
- Ali, H.K., Esam, H.M. and Fathy, M.D. (2000). Influence of malt on rheological and baking properties of wheat cassava composite flours. LWT- Food Science and Technology 33:159-164. .
- Alsmeyer, R.H., Alsmeyer, A.E. and Cunningham, M.L. Happich. (1974). Equations predict PER from amino acid analysis Food Technol., 28 pp. 34-40.
- Amin, R., Murphy, N., Edge, J., Ahmed, M.L., Acerini, C.L. and Dunger, D.B. (2002). A longitudinal study of the effects of a gluten-free diet on glycemic control and weight gain in subjects with type 1 diabetes and celiac disease. Diabetes care, 25(7), 1117-1122.
- Ammar, A.F., Siddeeg, A., Aqlan, F.M., Howladar, S.M., Refai, M.Y., Afifi, M. and Baeshen, M. (2020). Shelf life extension of wheat bread by alhydwan flour and Carboxymethylcellulose and improvement of their quality characteristics, dough rheological and microstructure. International journal of biological macromolecules, 156:851-857.
- Aprodu, I., Alexandra B.E. and Banu, I. (2016). Influence of protein and water addition on gluten-free dough properties and bread quality. International Journal of Food Engineering, 12(4):35.

. Arnarson, A. (2017). Signs and symptoms of

protein deficiency, Healthline.com. [Online]. Available, https://www.healthline.com/nutritio n/protein-deficiency-symptoms

- Barada, K., Bitar, A., Mokadem, M.A.R., Hashash, J.G. and Green, P. (2010). Celiac disease in Middle Eastern and North African countries: a new burden.World journal of gastroenterology: WJG, 16(12), 1449. –1457. / Google Scholar
- Bojňanská, T., Frančáková, H., Líšková, M. and Tokár, M. (2012). Legumes – the alternative raw materials for bread production. Journal of Microbiology, Biotechnology and Food Sciences 1: 876-886. Bourne, M. (2002). Food texture and viscosity: concept and measurement. Elsevier.
- Caio, G., Volta, U., Sapone, A., Leffler, D.A., De Giorgio, R., Catassi, C. and Fasano, A. (2019).Celiac disease: a comprehensive current review. BMC medicine, 17,1-20.
- Brown, J. (1993) Advances in Bread Making Technology. In: Kamel, B.S. and Stauffer, C.E., Eds., Advances in Baking Technology, Blackie Academic and Professional, Glasgow, 38-87.
- Campbell.J.A.(1963). Methodolgy of protein evaluation.PAG.Nutr.R.101 Add.37, June Meeting, New York.
- Capriles, V.D., and Arêas, J.A.G. (2014). Novel approaches in gluten-free breadmaking: interface between food science, nutrition, and health. Comprehensive Reviews in Food Science and Food Safety, 13(5), 871-890.
- Chapman, D.G., Castillo, R. and Campbell, J.A. (1959). Evaluation of protein in foods: 1.A method for the determination of protein efficiency ratios. Canadian Journal of Biochemistry and Physiology, 37(5), 679-686.
- Conte, P., Fadda, C., Drabinska, N. and Krupa-Kozak, U. (2019). Technological and nutritional challenges, and novelty in gluten-free breadmaking: A review. Polish Journal of Food and Nutrition Sciences, 69(1):5-21.
- Davidou, S., LeMeste, M., Debever, E. and Bekaert, D., (1996). A contribution to the study of staling of white bread: effect of water and hydrocolloid. Food Hydrocolloids 10 (4), 375–383.

- Demirkesen, I., Mert, B., Sumnu, G. and Sahin, S. (2010). Rheological properties of gluten-free bread formulations. Journal of food Engineering, 96(2), 295-303.
- Drulyte, D. and Orlien, V. (2019). The effect of processing on digestion of legume proteins. Foods, 8(6), 224.
- Eggum, B.O. (1973). A study of certain factors influencing protein utilization in rats and pigs.
- Miscellaneous: Beretning fra Forsoegs laboratoriet No.406, 173pp.
- El-Tawil, A.R (1998). Technological, Biological and Microbiological studies on produced from whole maize grain Ph.D. Thesis fac.of Agric.Cairo Univ.
- Esteller, M.S., Amaral, R.L. and Lannes, S.C.D.S. (2004). Effect of sugar and fat replacers on the texture of baked goods. Journal of texture studies, 35(4):383-393.
- FAO/WHO Committee (1973). Energy and protein requirements Expert Committee. FAO. Nut. Meeting Report Ceries No., 52.
- FAO/WHO/UNU (1985) Food and Agriculture Organization/World Health Organization/United Nations University Energy and Protein Requirements Report of a Joint Expert Consul tation. WHO Technical Report Series, No 724. Geneva.
- FAO/WHO (1991). Protein quality evaluation in human diets. Report of the Joint FAO/WHO Expert Consultation. FAO Food and Nutrition Paper No. 51. Rome.
- FAO (2004). Food and Agriculture Organization of United Nations.Economic and Social department.Available from FAOSTAT statistical database agriculture. Rome, Italy.
- Flamm, G., Glinsmann, W., Kritchevsky, D., Prosky, L. and Roberfroid, M. (2001). Inulin and oligofructose as dietary fiber: a review of the evidence. Critical reviews in food science and nutrition, 41(5):353-362.
- Gilani, G.S., Cockell, K.A. and Sepehr, E., (2005).Effects of antinutritional factors on protein digestibility and availability in foods, J. AOAC Int., 88(3):967–987.

- Gonzales, I.C., Gonzales, F.R., Quindara, H.L., Belino, P.L. and Botangen, E.T. (2016). Chickpea (Garbanzos) Its Nutritional and Economical Value. IOSR Journal of Economics and Finance, 7(4), 01-06.
- Gujral, H.S. and Pathak, A. (2002) Effect of Composite Flours and Additives on the Texture of Chapatti. Journal of Food Engineering, 55, 173-179.
- Gujral, H.S. and Rosell, C.M. (2004). Functionality of rice flour modified with a microbial transglutaminase. Journal of Cereal Science, 39(2), 225-230.
- Hernandes, H.H. and Bates, L.S. (1969). A simple method for tryptophan determination in maize. CYMMIT Research Bulletin, 13, 134-139.
- Hegsted, D.M., Mills, R.C., Elvehjem, C.A.F. and Hart, E.B. (1941). Choline in the nutrition of chicks. J. Biol. Chem. 138:459-465.
- Hefnawy, T.M.H., El-Shourbagy, G.A. and Ramadan, M.F. (2012). Impact of adding chickpea (Cicer arietinum L.) flour to wheat flour on the rheological properties of toast bread. International Food Research Journal, 19(2):521-525.
- Ibrahim, M.A.K. (2011). Chemical and biological studies on some bakery products (Doctoral dissertation, Ph.D. Thesis, Fac. Agric., Benha Univ., Egypt)., 184 p.
- Ijarotimi, O.S. (2008). Protein and hematological evaluations of infant formulated from cooking banana fruits (Musa spp, ABB genome) and fermented bambara groundnut (Vigna subterranean L. Verdc) seeds. Nutrition Research and Practice, 2(3), 165-170.
- Jisha, S., Padmaja, G., Moorthy, S.N. and Rajeshkumar, K. (2008). Pre-treatment effect on the nutritional and functional properties of selected cassava-based composite flours. Innovative Food Scienace & Emerging Technologies, 9(4), 587-592.
- Kadria M. El Sayed, M. Abd El-Sattar Ahmed, A.Salah Abu Donkol and Younes, M.H. (2021).Prospective of Local Cereal and Pulses for Free Gluten Toast in Egypt. Alex. J. Agric. Sci., 66:

(4), 105-119.

- Karaoğlu, M.M. and Kotancilar, H.G. (2009). Quality and textural behaviour of par-baked and rebaked cake during prolonged Storage
- storage. International Journal of Food Science & Technology, 44(1), 93-99.
- Kaur, M. and Singh, N. (2005). Studies on functional, thermal and pasting properties of flours from different chickpea (Cicer arietinum L.) cultivars. Food chemistry, 91(3):403-411.
- Khattab, R.Y., and Arntfield, S.D. (2009). Nutritional quality of legume seeds as affected by some physical treatments. Antinutritional factors. LWT- Food Science and Technology, 42 (6):1113-1118.
- Khorshid, A.M, Rabie, S.M., Eman, M.S. and EL-Azab, M.A. (1997), Effect of sorbate and packing material on quality and shelf life of corn tortilla bread .International control, 1-3 December, 1997, Sherton montaza, Alex, Egypt.
- Kramer, A. and Twigg, B.A. (1970). Quality control for the food industry. Westport, CT, USA.
- Luz Fernandez, M.A.R.I.A. and Berry, J.W. (1989). Rheological properties of flour and sensory characteristics of bread made from germinated chickpea. International Journal of Food Science & Technology, 24(1), 103-110
- López, A.C.B., Pereira, A.J.G. and Junqueira, R.G. (2004). Flour mixture of rice flour, corn and cassava starch in the production of gluten-free white bread. Brazilian archives
- Makanjuola, O.M. (2017). Amino Acids Profile of Powdered Fermented Maize Meal (Ogi) Fortified With Powdered Unfermented Locust Bean Seeds (Pakia Biglobosa). IOSR Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT), 11(11), 28-32.
- Marinangeli, C.P., and House, J.D. (2017). Potential impact of the digestible indispensable amino acid score as a measure of protein quality on dietary regulations and health. Nutrition reviews, 75(8), 658-66.
- Matos, M.E. and Rosell, C.M. (2013). Quality indicators of rice-based gluten-free bread-

like products: relationships between dough rheology and quality characteristics. Food and Bioprocess Technology, 6, 2331-2341.

- Matsushita, K., Tamura, A., Goshima, D., Santiago, D.M., Myoda, T., Takata, K., and Yamauchi, H. (2020). Effect of combining additional bakery enzymes and high pressure treatment on bread making qualities. Journal of food science and technology, 57, 134-142.
- Michaelsen, K.F., Weaver, L., Branca, F., and Robertson, A., (2003). Feeding and nutrition of infants and young children. Guidelines for the Who European Region with Emphasis on the Former Soviet Countries. Europe, UK:WHO Regional Office for Europe.
- Miñarro, B., Albanell, E., Aguilar, N., Guamis, B., & Capellas, M. (2012). Effect of legume flours on baking characteristics of gluten-free bread. Journal of cereal science, 56(2), 476-481.
- Mahmoud, R.M., Yousif, E.I., Gadallah, M.G.E., and Alawneh, A.R. (2013). Formulations and quality characterization of gluten-free Egyptian balady flat bread. Annals of Agricultural Sciences, 58(1), 19-25.
- Mohammed, I., Ahmed, A.R., & Senge, B. (2012). Dough rheology and bread quality of wheat chickpea flour blends. Industrial Crops and Products, 36(1), 196-202.
- Mohammad R, Jafar M. Milani and Gisoo Maleki (2014). Physical Properties of Gluten-Free Bread Made of Corn and Chickpea. International Journal of Food Engineering 10(3):467– 472.
- Nagaprabha, P. and Prakash, J. (2009) Development and Quality Assessment of Green Gram Based Instant Dosai Mix. Journal of Food Science and Technology, 46, 418-422.
- Nithya, D.J., Bosco, K.A.S., Saravanan, M., Mohan, R.J. and Alagusundaram, K. (2016). Optimization of process variables for extrusion of rice– Bengal gram blends. Journal of Scientific and Industrial Research, 75, 108-114.
- Olojede, A.O., Sanni, A.I. and Banwo, K. (2020). Effect of legume addition on the physiochemi-

cal and sensorial attributes of sorghum-based sourdough bread. Lwt, 118, 108769.

- Oluwaniyi, O.O., Dosumu O.O. and Awolola G.V. (2010). Effect of local processing methods (boiling, frying and roasting) on the amino acid composition of four marine fishes commonly consumed in Nigeria Food Chem., 123 (2010), pp. 1000-1006.
- Oser, B.L. (1959). An integrated essential amino acid index for predicting the biological value of proteins. Protein and amino acid nutrition lbanese, A.A.A. by Academic Press, New York, 281.
- Pateras IMC. Bread spoilage and staling. In: Cauvain S.P. and Young L.S. (2007). Technology of bread making. New York: Springer, 275–99.
- Prasad, K., Singh, Y. and Anil, A. (2012). Effects of grinding methods on the characteristics of Pusa 1121 rice flour. Journal of Tropical Agriculture and Food Science, 40(2), 193-201.
- PAG. (1971). Guidelines on protein rich mixtures for use in weaning foods. Protein Advisory Group.
- Pruska-Kędzior, A., Kędzior, Z., Gorący, M., Pietrowska, K., Przybylska, A. and Spychalska, K. (2008). Comparison of rheological, fermentative and baking properties of gluten-free dough formulations. European Food Research and Technology, 227, 1523-15.
- Rojas, J.A., Rosell, C.M. and De Barber, C.B. (1999). Pasting properties of different wheat flour-hydrocolloid systems. Food hydrocolloids, 13(1), 27-33.
- Samah, A.S. (2004). Review Article: Nutrient composition in fortified complementary foods. Egyptian Journal of Agricultural Research, 82(3):75-100.
- Sarwar Gilani, G., Wu Xiao, C. and Cockell, K.A., (2012). "Impact of anti -nutritional factors in food proteins on the digestibility of protein and the bioavailability of amino acids.
- Seleem, H.A.S. (2000). Studies on addition of some corn and sorghum varieties to wheat flour and balady bread characteristic (Doctoral dissertation, Ph.D. Thesis, Fac. Agric., Cairo

Univ., Egypt).

- Sharima-Abdullah, N., Hassan, C.Z., Arifin, N. and Huda-Faujan, N. (2018). Physicochemical properties and consumer preference of imitation chicken nuggets produced from chickpea flour and textured vegetable protein. International. Food Research Journal, 25(3), 1016-1025.
- Sidhu, J.S., Al-Saqer, J. and Al-Zenki, S. (1997) Comparison of Methods for Assessment of the Extent of Staling in Bread. Food Chemistry, 58, 161-167.
- Silva, F.D.A.S., and de Azevedo, C.A.V. (2009). Principal Components Analysis in the Software Assis tat-Statistical Assistance. In 7th World Congress on Computers in Agriculture Conference Proceedings, 22-24 June 2009, Reno, Nevada (p1). American Society of Agricultural and Biological Engineers.
- Srilakshmi, B. (2007). Food science, 4th Ed. New age internation (P) Ltd. New Delhi. 170-174.
- Watson, S.A. (1997). Structure and composition. In S.A. Watson and P.E. Ramstad (Eds.). Corn: Chemistry and technology, p. 311–349. Saint Paul, MN, USA: AACC.
- Xu, M., Jin, Z., Simsek, S., Hall, C., Rao, J. and Chen, B. (2019). Effect of germination on the chemical composition, thermal, pasting, and moisture sorption properties of flours from chickpea, lentil, and yellow pea. Food chemistry, 295, 579-587.
- Zengin, G., Aktumsek, A., Guler, G.O., Cakmak, Y.S., Giron-Calle, ' J., Alaiz, M. and Vioque, J. (2012). Nutritional quality of protein in the leaves of eleven Asphodeline species (Liliaceae) from Turkey. Food Chem. 135, 1360–1364.