Journal of Applied Veterinary Sciences, 10 (2): 18-27 (April, 2025). ISSN: Online: 2090-3308, Print: 1687-4072 Journal homepage : https://javs.journals.ekb.eg

Effect of Effervescent Formulation of Silymarin against Experimental Hepatotoxicity in Rats: Involvement of NRF2/HO-1, PI3K/AKT and TLR4/NFκB Pathways

 Hany G. Attia^{1*}, Bassim M.S.A. Mohamed^{2*}, Saeed M. Alasmari³, Sahar S AbdelRahman⁴, Mohamed Aleraky⁵, Abdulwahab Alqahtani⁶, Marawan A. Elbaset² and Hany M. Fayed²
 ¹Department of Pharmacognosy, College of Pharmacy, Najran University, Najran 66462, Saudi Arabia
 ²Department of Pharmacology, Medical Research and Clinical Studies Institute, National Research Centre, Cairo, P.O. 12622 Bohouth St. Dokki, Egypt

 ³Department of Biology, College of Science and Arts, Najran University, Najran 1988, Saudi Arabia
 ⁴Department of Pathology, Faculty of Veterinary Medicine, Cairo University, Giza, Egypt
 ⁵Department of Clinical Pathology, College of Medicine, Najran University, Najran 1988, Saudi Arabia
 ⁶Department of Pediatrics, College of Medicine, Najran University, Najran 1988, Saudi Arabia
 *Corresponding Authors: Bassim M.S.A. Mohamed, E-Mail: bassim.mohamed@umontreal.ca Hany G. Attia, E-Mail: hgattia@nu.edu.sa

ABSTRACT

Despite the exploration of pharmaceutical and natural liver-protecting drugs, the number of hepatotoxicity deaths continues to rise. Milk thistle flavonoids, _ silymarin, have been widely investigated and demonstrated liver disease protection. Poor solubility and absorption restrict silvmarin bioavailability. Effervescent formulations may boost silvmarin absorption by improving gastrointestinal solubility. Our study compared the effectiveness of silymarin effervescent formulation in protecting the liver from thioacetamide (TAA)-induced oxidative damage at both low and high doses. Study findings may have an implication on chemotherapy-induced off-target harmful oxidative insult. 24 adult male rats were separated into normal control, hepatotoxic (TAA) (100 mg/kg body weight), and TAA plus silymarin (50 and 100 mg/kg b.wt) groups. Serum liver enzyme, hepatic antioxidant, lipid peroxidation, and inflammatory indicators were measured. The modest dose's great bioavailability was shown by its potency being identical to the doubled dose. TAA caused hepatic injury, as evidenced by elevated liver enzymes (ALT and AST) and tissue levels of MDA, TLR4, TGFβ1, TNF-α, IL-6, NF-κB, and p-NFkB. TAA also decreased tissue SOD, GSH, and HO-1 levels as well as Nrf2 level and expression. Increased gene expressions of Akt, PI3K, and TLR4 were observed. Silymarin's antioxidant and anti-inflammatory actions reduced hepatotoxicity via upregulating Nrf2/HO-1 and downregulating PI3K/Akt and TLR4/NFkB pathways. This study shows that effervescent silymarin formulation is a powerful bioavailable treatment for liver oxidative toxicity in chemotherapyrelated toxicities.

Keywords: Chemotherapy, Nrf2/ HO-1, Oxidative Damage, PI3K/Akt, Silymarin effervescent formula, Thioacetamide, TLR4/ NFκB.

INTRODUCTION

The liver plays a significant role in the body's process of detoxifying dangerous medications, xenobiotics, and organic pollutants (Shirani *et al.*, 2020; Taleb *et al.*, 2023). The prevalence and mortality of liver disease remain major global issues. The main factors known to induce liver damage are viral hepatitis, antibiotics, chemotherapeutic drugs, hepatotoxins including carbon tetrachloride (CCI4), and TAA (Teksoy *et al.*, 2020; Bashandy *et al.*, 2022; Elbaset *et al.*, 2023b).

Original Article: DOI:https://dx.doi.org/10.21608/j avs.2025.348491.1503 Received : 08 January, 2025. Accepted: 18 February, 2025. Published in April, 2025.

This is an open access article under the term of the Creative Commons Attribution 4.0 (CC-BY) International License . To view a copy of this license, visit:

http://creativecommons.org/licenses/by/4

J. Appl. Vet. Sci., 10(2): 18-27.

Thioacetamide (TAA) is a thio-sulfurcontaining chemical (Abd El-Rahman and Fayed 2019). One dosage of TAA administration induces acute liver damage (Chen *et al.*, 2008). Reactive oxygen species (ROS) and TAA intermediates are formed when cytochrome P450 metabolically activates TAA. Following their binding to physiologically significant molecules, these ROS can amplify cellular oxidative stress, lipid peroxidation, and glutathione depletion (Pallottini *et al.*, 2006). TAA has been a widely used



model for research on liver damage and fibrosis (**Sarma and Hanzlik 2011**). TAA offers a liver-damaging model with a high ability for regeneration that is similar to that of a human with protracted injury.

Further, the TAA liver toxicity is similar to the pathological oxidative burden induced bv chemotherapeutic agents (Delire et al., 2015). A significant event in the pathogenesis of many hepatic diseases is oxidative stress. Hepatocytes' mitochondria and endoplasmic reticulum are where the cvtochrome P-450 enzymes mostly create ROS (Arauz et al., 2016). Liver cells can regulate the level of oxidative stress and the balance of oxidant and antioxidant molecules under physiologically normal circumstances. While under stressful circumstances, oxidative stress is increased, and the liver's capacity to maintain this balance is compromised (Cichoz-Lach and Michalak 2014).

Natural remedies are increasingly popular for preventing and treating liver disease (Abdelbaset et al., 2014; Ayoub et al., 2022; Ogaly et al., 2022; Taleb et al., 2023). There is evidence that oxidative stressrelated liver damage can be mitigated by antioxidants and free radical-scavenging substances (Bruck et al., 2004; Elbaset et al., 2022; Moussa et al., 2022). It is generally known that silymarin, derived from the milk thistle (Silybum marianum), has hepatoprotective and antioxidant activities (Vargas-Mendoza et al., 2014). Silvmarin is composed of the flavonoids silvbin, silydianin, and silychristin. (Wu et al., 2009). Saller et al., (2001) showed that silymarin protects the liver from oxidative stress caused by alcohol. Additionally, silymarin has been demonstrated to protect the liver against carbon tetrachloride (CCl₄) damage and the kidney from acetaminophen damage (Lamia et al., 2021). These effects are primarily attributable to decreased lipid peroxidation (Chen et al., 2012).

Manna et al., (1999) showed that silymarin has limited bioavailability due to poor solubility and absorption. This means a large portion of the ingested silymarin isn't absorbed by the body. Since silymarin solubility is essential for achieving an ideal and effective bioavailability, formulation techniques aimed at enhancing its solubility are vital. According to the reviewed research, silvmarin-based formulations have become more advanced and efficient over time, which makes sense given the development of nanotechnologies and nano systems intended to enhance the delivery of active key bioactive molecules that are poorly soluble in water (Di Costanzo and Angelico 2019). Drug delivery is an active area of research (Zahin et al., 2020; Ahmad et al., 2021).

Further, effervescent formulations offer a potential solution by enhancing silymarin's dissolution in the gastrointestinal tract, leading to increased

absorption (Ahmad *et al.*, 2023). These findings motivate us to investigate silymarin's effervescent formulation as a potent bioavailable hepatoprotective role in TAA toxicity as well as molecular approaches upon toxicity.

MATERIALS AND METHODS

1. Animals

At the National Research Centre (NRC, Egypt), 24 Wistar rats, males (180-220 g; 7 weeks) were acquired from the "Animal House." The rats were housed in an environment with a constant temperature of 25 °C and 12-hour light/dark cycles. Adherence to both national and international ethical standards was ensured in the treatment of the animals. Approval for all experimental procedures was obtained from the Institutional Animal Care and Use Committee at Cairo University. (Approval certificate: Vetcu23052022441).

2. Chemicals

The supplier of thioacetamide was "Sigma-Aldrich, USA". Silymarin, Na HCO3, sodium citrate, citric acid compounds were provided with the best analytical-grade ingredients. Silymarin effervescent formula: Silymarin 78%, Na HCO3 19%, sodium citrate 1%, citric acid 2%.

3. Experimental design

Four sets of six rats each were randomly distributed; The rats in the first set were designated as the control group, received intraperitoneal injections of normal saline three times weekly for a duration of two weeks (0.5 ml/ 100 g rat). The rats in the second set (referred to as the TAA-treated group) were administered intraperitoneal injections of TAA (100 mg/kg body weight) three times weekly over a period of two consecutive weeks. The third sets of rats were given silymarin at a dosage of 50 mg/kg/day orally for two weeks following the TAA injections in a similar manner to the second group. The fourth set of rats were administered silvmarin at a dosage of 100 mg/kg/day orally for a period of two weeks after receiving TAA injections following the same protocol as the second group.

4. Preparation of blood samples and tissue specimens

Twenty-four hours after the end of the experiment, during the administration of ketamine for sedation, blood samples were collected from the tail vein of each rat. Serum samples intended for biochemical analysis were maintained at a temperature of -20°C. Following the extraction of blood samples, rats were euthanized through cervical dislocation, and expeditiously, their livers were extracted, cooled using ice-cold saline, and desiccated. The left lobe of the liver from each rat was excised and preserved in buffered neutral formalin at a concentration of 10% for

subsequent histological investigations. A portion of this tissue was also weighed and stored at -80°C for advanced molecular and biological scrutiny.

5. Evaluation of biomarkers for liver function

Colorimetric analysis was performed on the aspartate aminotransferase (AST) (Catalog #AS1061, Biodiagnostic®, Egypt) and alanine aminotransferase (ALT) (Catalog #AL1031, Biodiagnostic®, Egypt) activities in serum, as well as the serum albumin levels (Catalogue #AB1010, Biodiagnostic®, Egypt).

6. Evaluation of the markers of oxidative stress

The levels of "glutathione (GSH) content, malondialdehyde (MDA), and superoxide dismutase (SOD)" were assessed in liver tissue samples using particular colorimetric assays (BioVision, Milpitas, CA, USA; Cat: No: K464-100, K739-100, and K335-100, respectively), in order to assess oxidation stress and antioxidant defense.

7. Biomarkers of liver inflammation and profibrosis by enzyme-linked immunosorbent assay (ELISA)

Using particular Rat ELISA kits (SunLong Biotech Co., LTD, China; Catalogs Number: SL0722Ra, SL0411Ra, SL0537Ra, and SL0705Ra, respectively), the important signals in liver injury and inflammation, tumor necrosis factor (TNF- α), interleukin-6 (IL-6), nuclear factor-kappa B (NF- κ B), and transforming growth factor-beta (TGF- β 1) were assayed in liver homogenate as per the manufacturer's guidelines. Toll like receptor 4 "TLR4" in hepatic homogenate was tested using the production protocols

of Sunlong Biotec Com. LTD in Zhejiang, China (Catalog No. SL0699Ra).

8. Nrf2 and HO-1 in the liver using enzymelinked immunosorbent assays (ELISA)

The ELISA kits (Sunlong Biotech Co., Zhejiang, China) were used to quantify the liver level of nuclear factor erythroid 2-related factor-2 (Nrf2) (SL0985Ra) and heme oxygenase 1 (HO-1) (SL0341Ra).

9. QRT-PCR for the expression Nrf2, AKT, PI3K and TLR4

RNA extraction

For the reverse transcription and PCR of the isolated RNA, the SuperScript IV One-Step RT-PCR kit (Cat# 12594100) was provided by Thermo Fisher Scientific, located in Waltham, Massachusetts, USA. A thermal profile was executed utilizing a 96-well plate StepOne equipment from Applied Biosystems, USA, the initial denaturation phase is comprised of 40 cycles, with each cycle lasting 10 s at 98 °C, 10 s at 55 °C, and 30 s at 72 °C for the amplification step. Following this, the reverse transcription process extends over a duration of 10 minutes at 45 °C, succeeded by RT inactivation lasting 2 minutes at 98 °C. Upon completion of the RT-PCR procedure, the data concerning the target genes and housekeeping genes were presented using the cycle threshold (Ct) method. Oligonucleotide sequences for the forward and reverse primers are detailed in Table (1). The mRNA levels of tested genes Nrf2, AKT, PI3K, and TLR4 were adjusted for variability using the mean critical threshold (CT) expression.

Gene		Sequence (5'-3')		
Nrf2	FD	AAAATCATTAACCTCCCTGTTGAT	NM_010902.5	
	RE	CGGCGACTTTATTCTTACCTCTC		
AKT	FD	CGGATACCATGAACGACGTAG	NM 033230.2	
	RE	GCAGGCAGCGGATGATAAAG	1001_033230.2	
PI3K	FD	TTAAACGCGAAGGCAACGA	VM 022808071 1	
	RE	CAGTCTCCTCCTGCTGTCGAT	AWI_052696971.1	
TLR4	FD	CCGTCACCACATACTGCCTTTA	NM_001109668.1	
	RE	GCAGTTTGGACTATTGAAATACGAAA		

Table 1: List of "oligonucleotide primers" used in qPCR.

10. Examination of the histopathology

Samples collected from the livers of rats from different groups underwent standardized processing to create paraffin sections following a twenty-four-hour fixation in a 10% concentration of neutral buffered formalin. Subsequently, the specimens were rinsed with distilled water, dehydrated using ethanol solutions, and clarified with xylene. Subsequent to this, sections measuring four to five micrometers in thickness were generated from the paraffin blocks. Following the placement of the tissue sections on glass slides, deparaffinization was carried out, followed by staining with hematoxylin and eosin (H&E). An experienced evaluator conducted all histopathological examinations, throughout blinding maintaining the sample identification process to prevent bias.

The assessment of histological impairments was conducted utilizing the METAVIR test, which categorizes the extent of fibrosis and includes various indicators of liver damage such as bile duct proliferation, steatosis, portal vein dilatation, and fibrosis. These lesions were classified as follows: F0 =absence of lesions (-), F1 = presence of mild lesions (+), F2 = presence of moderate lesions (++), and F3 =presence of severe lesions (+++).

11. Statistical analysis

Prior to proceeding with the statistical analysis, the normality assumption was validated using the Shapiro-Wilk test. The results are expressed in the form of means \pm standard error. Subsequently, the data underwent analysis through one-way ANOVA, followed by the Tukey-Kramer post hoc test. The statistical analysis and graphical representations were carried out using GraphPad Prism software (version 9, USA). A significance level of P<0.05 was chosen for all statistical tests.

RESULTS

Impact of silymarin on the serum levels of AST, ALT, and albumin in rats receiving TAA

As markers of liver damage, serum ALT, AST, and albumin were routinely measured. Rats given TAA injections showed a substantial increase in serum AST and ALT activities, whereas serum albumin was reduced in comparison to the non-treated control group. Silymarin administration significantly reduced the levels of AST and ALT and significantly elevated the albumin level as compared to the hepatotoxic group (**Fig.1 a, b, c**).



Fig.1: The impact of silymarin administration on the liver function indicators (a) ALT, (b) AST, and (c) albumin in rats subjected to thioacetamide-induced intoxication. Data are presented as mean \pm SE. * vs. normal control group, [@] vs. TAA group, [#] vs. Silymarin groups at *p*<0.05.

Impact of silymarin on oxidation stress in rats receiving TAA

Figure 2 (a, b, c) clearly showed how TAA had a negative impact on the liver's antioxidant defenses and significantly increased the lipid peroxidation product MDA in the liver tissue. According to the table, silymarin was able to restore the liver's antioxidant status by markedly elevating both the liver's SOD activity and GSH levels. Malondialdehyde levels were markedly reduced by silymarin to normal levels compared to the hepatotoxic group.



Fig. 2: The impact of silymarin administration on the liver oxidative stress indicators (a) GSH, (b) MDA, and (c) SOD in rats subjected to thioacetamide-induced intoxication. Data are presented as mean \pm SE. * vs. normal control group, [@] vs. TAA group, [#] vs. Silymarin groups at *p*<0.05.

Impact of silymarin on liver's profibrotic and inflammatory marker content in rats receiving TAA

In comparison to the negative control group, TAA dramatically elevated the levels of TLR4, TNF- α , IL-6, TGF β 1, NF- κ B, and p-NF- κ B in the rats' livers (**Fig. 3 a-f**). Treatment with silymarin significantly and dose-dependently decreased the levels of aforementioned parameters in the rat livers compared to the TAA-treated group. Silymarin at a dose of 100 mg/kg revealed the best effects in inhibition of inflammatory response.



Fig. 3: The impact of silymarin administration on the liver profibrotic and inflammatory indicators (a) TLR 4, (b) TNF- α , (c) II-6, (d) TGF1 β -1, (e) NF- κ B, and (f) p-NF- κ B in rats subjected to thioacetamide-induced intoxication. Data are presented as mean \pm SE. * vs. normal control group, [@] vs. TAA group, [#] vs. Silymarin groups at *p*<0.05.

Impact of silymarin on the liver levels of Nrf2 and HO-1 in rats receiving TAA

Hepatic Nrf2 and HO-1 levels significantly decreased in the TAA group compared to the negative control group (**Figs. 4 a, b**). Rats given both dosages of silymarin, as opposed to the TAA group, exhibited a marked rise in Nrf2 and HO-1 levels.

Impact of silymarin on genes expression of Nrf2, TLR4, AKT, and PI3K in rats receiving TAA

In hepatic tissue, the relative expression of the mRNAs for Nrf2 (**Fig.5d**) showed that TAA significantly downregulated their expression; meanwhile, the mRNAs for TLR4, Akt, and PI3K expressions were significantly upregulated when compared to the control groups (**Fig. 5 a, b, c**). Silymarin administrations could up regulate Nrf2 expression and down regulate PI3K, Akt, and TLR4 expressions, compared with the TAA group.



Fig.4: The impact of silymarin administration on the liver antioxidant proteins (a) Nrf2 and (b) HO-1 in rats subjected to thioacetamide-induced intoxication. Data are presented as mean \pm SE. * vs. normal control group, [@] vs. TAA group, [#] vs. Silymarin groups at *p*<0.05.



Fig. 5: The impact of silymarin administration on the liver mRNA expression (a) TLR4, and (b) AKT, (c) PI3K, and (d) Nrf2 in rats subjected to thioacetamide-induced intoxication. Data are presented as mean \pm SE. * vs. normal control group, [@] vs. TAA group, [#] vs. Silymarin groups at *p*<0.05.

Impact of silymarin on liver histopathology in rats intoxicated with TAA

Liver specimens from the control group rats exhibited a normal histological arrangement of central veins, portal regions, and hepatic cell cords (**Fig. 6**). Conversely, liver specimens from the TAA group rats demonstrated a profound fibrotic response. The hepatic capsule displayed distinct corrugation, accompanied by the absence of typical hepatic lobulation. Fibrous proliferation extensively enlarged the portal triads, characterized by mononuclear inflammatory cell infiltration, proliferation of cholangiolar epithelium leading to multiple newly formed bile ductules, and vascular congestion



Fig. 6: Impact of silymarin on liver histopathology in rats intoxicated with TAA

Table 2: Showing scoring of the histologicallesions observed.

Itom	Control	TAA	TAA+Sil	TAA+Sil	
nem	group	group	(50mg/kg)	(100mg/kg)	
Fibrosis	-	+++*	+@	+@	
Bile duct proliferation	-	+++*	++*	+@	
Steatosis	-	+++*	++*	+@	
Dilatation of portal vein	-	+++*	+@	+@	

Data are presented as mean \pm SE. * vs. normal control group, [@] vs. TAA group, [#] vs. Silymarin groups at p < 0.05.

METAVIR score

Severity of liver fibrosis were graded as (- equal no lesion), (+++ equal severe fibrosis including nodules, sepat, and portal tract fibrosis), (++ equal moderate fibrosis including sepat and portal tract fibrosis), (+ mild portal tract fibrosis) (**Table 3**).

Table 3: showing grading of histological fibrosisusing the METAVIR score.

score	Histological character	Normal control group	TAA group	TAA+Sil (50mg/kg)	TAA+Sil (100/kg)	
		The number of animals showed the lesion per group				
F0	No fibrosis	6	-	-	-	
F1	Portal fibrosis	-	-	2	5	
F2	Portal fibrosis with few septa	-	-	4	1	
F3	Portal fibrosis with numerous septa	-	6	-	-	
F4	Cirrhosis	-	-	-	-	

DISCUSSION

Silymarin, a group of flavonoids found in milk thistle, has been extensively studied and shown protective activity in liver diseases (Gillessen and Schmidt 2020). However, silymarin has limited bioavailability due to poor solubility and absorption. This means a large portion of the ingested silvmarin isn't absorbed by the body. Effervescent formulations offer a potential solution by enhancing silvmarin's dissolution in the gastrointestinal tract, leading to increased absorption. Our study showed that the small dose and doubled dose of silymarin have a very similar potency of reversing the liver toxicity induced by TAA, reflecting the potent bioavailable efficacy of the small dose of the effervescent formulation. A model that can give inference to the potent bioavailable effect in treating pathological oxidative burden of chemotherapeutic drugs. The liver's biotransformation of TAA into TAA sulfoxide and thioacetamide-S. Sdioxide makes TAA a powerful centrilobular hepatotoxicant. Due to their ability to alter lipid and protein structure, these reactive metabolites are extremely harmful to hepatocytes (Hajovsky et al., 2012; Ayoub et al., 2022; Elbaset et al., 2022; Abdel-Rahman et al., 2023).

Accordingly, our study found that TAA rats had elevated levels of hepatic injury markers and disrupted hepatic lobules, both of which suggested injured hepatic cells. The current study's findings, along with those of earlier studies, showed that TAA can cause liver oxidative damage that is shown as lipid peroxidation and impairment of the liver's antioxidant defense mechanisms (Ramadan et al., 2018; Abd El-Rahman and Fayed, 2019). According to reports, silymarin exhibits protective effects across a variety of tissues and exhibits antioxidant activity (Vargas-Mendoza et al., 2014). Treatment with silymarin significantly reduced the levels of MDA, an indication of oxidative stress. In contrast, after silymarin administration, the activities of SOD and GSH in the liver tissues of hepatotoxic rats were significantly elevated.

TAA has the capacity to induce the release of several hepatotoxic cytokines from immune cells (**Jiao** *et al.*, **2022**) i.e., TNF- α , IFN- γ , and IL-6. Through the TLRs signal pathway, these cytokines further contribute to the development of hepatocyte injury by stimulating and activating immune cells (**Xu** *et al.*, **2013**). A class of receptors known as toll-like receptors (TLRs) is implicated in inflammation and innate immunity. TLRs play a role in liver fibrosis, which is brought on by a variety of common liver infections, including parasite, viral, and toxin-induced hepatitis (**Cengiz** *et al.*, **2015**). Kupffer cells, hepatic stellate cells, hepatocytes, sinusoidal endothelial cells, biliary epithelial cells, and

hepatic dendritic cells are only a few of the liver cells that express TLRs broadly (**Xu** *et al.*, **2011**), which are crucial for the health of the liver (**Kim** *et al.*, **2010**).

Many therapeutic medications that prevent liver injury may interact with the TLR4 signaling system, according to previous studies that suggest that TLRmediated signals are considered to be involved in practically all liver illnesses (Zhai et al., 2016). When the TLR4 signal pathway is activated, a series of things happen, such as the nuclear factor kappa B (NF- κ B) p65 moving into the nucleus, which increases the production and release of inflammatory cytokines (TNF- α , TGF β 1, and IL-6) (Lawrence 2009). Further, we inspected the TLR4 expression. We show that we found that effervescent silvmarin formulation in lower dose administration after TAA toxicity was more potent than the doubled dose in the anti-inflammatory properties due to its significant suppression of TLR4 expression and blockage of the NF-kB signaling pathway. Our findings demonstrated that low-dose effervescent silymarin suppressed the generation of inflammatory cytokines, which are involved in TLR4-NF- kB signaling.

We thoroughly examined alternative potential mechanisms and found that effervescent silymarin formulation in lower dose administration after TAA toxicity with potency as the doubled dose altered the expression of many different genes in both up- and down-regulation. The cellular kinase PI3K produces the intracellular second messenger phospholipid PIP3 by upstream activating other signaling molecules like protein kinase B "Akt" (Troutman et al., 2012). Recently, a number of biological processes, including inflammation and cell differentiation, have been connected to the PI3K/Akt pathway (Mohamed and Hafez 2019). Although the significance of PI3K/Akt in inflammation is debatable (Troutman et al., 2012), our results, along with others, suggested that this pathway has a positive effect on inflammation in the liver (Hazeki et al., 2007; Kidd et al., 2008). Silymarin downregulated the expression of the PI3K/Akt gene in hepatic tissue.

The hepatic tissue's normal pattern and antioxidant state were successfully restored by the lowdose effervescent silymarin. It was shown that an essential mechanism in normal liver cells to counteract oxidative stress is mediated by nuclear factor erythroid 2-related factor (Nrf2) and its antagonist kelch-like ECH-associated protein 1 (Keap1) (**Wang et al., 2016**). The antioxidant response element (ARE), which modifies detoxification and controls the production of antioxidant genes, is thought to be primarily directed by Nrf2 (**Raslan et al., 2021**). In a healthy state, Keap1 and Nrf2 interact, and Nrf2 remains in the cytoplasm. Oxidative stress causes Nrf2 to separate from Keap1 and migrate to the nucleus, where it binds to ARE and activates downstream particular genes, such as quinone oxidoreductase 1 (NQO1), heme oxygenase (HO-1), and glutathione, which work to combat oxidative stress. Therefore, treatments that might activate Nrf2 may be helpful in reducing liver damage caused by oxidative stress. The current study concentrated on lower-dose effervescent silymarin's capacity to upregulate the antioxidant genes HO-1 and Nrf2 in comparison to controls and to the doubled-dose treatments. Rasha et al., who attributed silymarin's protective effect against TAA-induced acute liver damage to its induction for the Nrf2/HO-1 pathway, provide strong support for our findings (Hussein et al., 2021). Another investigation examined silymarin's potential to shield rats from acute kidney damage brought on by paraquat by inhibiting Nrf2, which reduced oxidative stress and cellular damage (Jia et al., 2022).

Silymarin effervescence at a relatively low dose (50 mg/kg body weight) was potent and had a strong bioavailable effect when comparing the treated TAA group to the naive control, regarding the normalization of Nrf2/HO-1, PI3K/Akt, and TLR4/NF- κ B values of the TAA silymarin-treated group to almost normal control levels. Compared to non-formulated silymarin (200 mg/kg body weight) in previous reports (**Tsai** *et al.*, **2008**).

It's worth mentioning that it has been reported that albino rabbits were used in in vivo experiments to quantify the pharmacokinetics of both the enhanced effervescent formulation with 200 mg of silymarin and the regular standard dosage form. After oral administration, the plasma concentration of silymarin was measured. The formulation's Cmax increased by 180%. For the formulation, the area under the curve was 310% greater (**Ahmad** *et al.*, **2023**).

Regarding the clinical relevance of using silymarin as a protective adjunct in chemotherapeutic treatments, silymarin may have the ability to modulate the xenobiotic system by affecting the activity of metabolizing phase I and phase II enzymes, leading to the protection of healthy cells from the insults of chemotherapeutic drugs. Additionally, silymarin and its primary bioactive ingredients block organic anion transporters (OAT) and ATP-binding cassettes (ABC), which helps to fight possible chemoresistance. Silymarin and its derivatives have two main roles: they prevent cancer cells from progressing through various stages, which forces them to evolve toward cell death, and they accumulate cancer cells in a particular stage of the cell cycle, which enables a particular anticancer agent to target a larger quantity of tumor cells (Delmas et al., 2020).

CONCLUSION

Silymarin effervescent low dose was potent and had a strong bioavailable effect compared to the doubled dose in reversing oxidative hepatotoxicity via its antioxidant and anti-inflammatory effects attributed to the upregulation of Nrf2/HO-1, downregulation of PI3K/Akt, and TLR4/NFkB pathways. Further, the present study highlights effervescent silvmarin as a potential formulation giving protection against chemotherapy-related toxicities in a clinical setting.

Conflicts of interest

The authors declare that there is no conflict of interest regarding the research data and tools used in this study.

REFERENCES

- ABDEL-RAHMAN, R. F., FAYED, H. M., MOHAMED, M. A., HESSIN, A. F., ASAAD, G. F., ABDELRAHMAN, S. S., SALAMA, A. A., ARBID, M. S., and OGALY, H. A., 2023. Apigenin role against thioacetamide-triggered liver fibrosis: Deciphering the $PPAR\gamma/TGF-\beta1/NF-\kappa B$ and the HIF/FAK/AKT pathways. Journal of Herbmed Pharmacology, 12: 202-213. https://doi.org/10.34172/jhp.2023.21
- ABD EL-RAHMAN, S. S. and FAYED, H. M., 2019. Targeting AngII/AT1R signaling pathway by perindopril inhibits ongoing liver fibrosis in rat. Journal of Tissue Engineering and Regenerative Medicine, 13(12): 2131-2141..https://doi.org/10.1002/term.2940
- ABDELBASET, M., SAFAR, M. M., MAHMOUD, S. S., NEGM, S. A., and AGHA, A. M., 2014. Red yeast rice and coenzyme Q10 as safe alternatives to surmount atorvastatin-induced myopathy in hyperlipidemic rats. Canadian Journal of Physiology and Pharmacology, 92: 481-489. https://doi.org/10.1139/cjpp-2013-0430
- AHMAD, A., MUBARAK, N. M., JANNAT, F. T., ASHFAQ, T., SANTULLI, C., RIZWAN, M., NAJDA, A., BIN-JUMAH, M., ABDEL-DAIM, M. M., HUSSAIN, S., and ALI, S., 2021. A Critical Review on the Synthesis of Natural Sodium Alginate Based Composite Materials: An Innovative Biological Polymer for Biomedical Delivery Applications. Processes, 9(1). https://doi.org/10.3390/pr9010137
- AHMAD, S., KHAN, J. A., KAUSAR, T. N., MAHNASHI, M. H., ALASIRI, A., ALQAHTANI, A. A., ALQAHTANI, T. S., WALBI, I. A., ALSHEHRI, O. M., ELNOUBI, O. A., MAHMOOD, F., and SADIQ, A., 2023. Preparation, Characterization and Evaluation of Flavonolignan Silymarin Effervescent Floating Matrix Tablets for Enhanced Oral Bioavailability. Molecules. 28(6).

https://doi.org/10.3390/molecules28062606

- ARAUZ, J., RAMOS-TOVAR, E., and MURIEL, P., 2016. Redox state and methods to evaluate oxidative stress in liver damage: From bench to bedside. Ann Hepatol, 15(2): 160-173.
- AYOUB, I. M., EL-BASET, M. A., ELGHONEMY, M. M., BASHANDY, S. A. E., IBRAHIM, F. A. A., AHMED-FARID, O. A. H., EL GENDY, A. E.-N. G.,

AFIFI, S. M., ESATBEYOGLU, T., FARRAG, A. R. H., FARAG, M. A., and ELSHAMY, A. I., 2022. Chemical Profile of Cyperus laevigatus and Its Protective Effects against Thioacetamide-Induced Hepatorenal Toxicity in Rats. Molecules, 27: 6470.https://doi.org/10.3390/molecules27196470

- BASHANDY, S. A. E., ABDELHAMEED, M. F., AHMED-FARID, O. A. H., MORSY, F. A., IBRAHIM, F. A. A., and ELBASET, M., 2022. The Pivotal role of Cerium oxide nanoparticles in thioacetamide induced hepatorenal injury in rat. Egyptian Journal of Chemistry, 65: 267-278. https://doi.org/10.21608/eichem.2022.115482.5242
- BRUCK, R., AEED, H., AVNI, Y., SHIRIN, H., MATAS, SHAHMUROV, М., AVINOACH, Z., Ι.. ZOZULYA, G., WEIZMAN, N., and HOCHMAN, A., 2004. Melatonin inhibits nuclear factor kappa B activation and oxidative stress and protects against thioacetamide induced liver damage in rats. J Hepatol, 86-93. https://doi.org/10.1016/S0168-40(1): 8278(03)00504-X
- CENGIZ, M., OZENIRLER, S., and ELBEG, S., 2015. Role of serum toll-like receptors 2 and 4 in nonalcoholic steatohepatitis and liver fibrosis. J 1190-1196. Gastroenterol Hepatol, 30(7): https://doi.org/10.1111/jgh.12924
- CHEN, I. S., CHEN, Y.C., CHOU, C.H., CHUANG, R.F., CHIU, C.-H., L.Y., and 2012. SHEEN, Hepatoprotection of silymarin against thioacetamideinduced chronic liver fibrosis. Journal of the Science of Food and Agriculture, 92(7): 1441-1447. https://doi.org/10.1002/jsfa.4723
- CHEN, T. M., SUBEO, Y. M., LEE, R. P., CHIOU, T. W., and HSU, B. G., 2008. Single dose intravenous thioacetamide administration as a model of acute liver damage in rats. Int J Exp Pathol, 89(4): 223-231. https://doi.org/10.1111/j.1365-2613.2008.00576.x
- CICHOZ-LACH, H., and MICHALAK, A., 2014. Oxidative stress as a crucial factor in liver diseases. 8082-World Gastroenterol, 20(25): J 8091.https://doi.org/10.3748/wjg.v20.i25.8082
- DELIRE, B., STARKEL, P., and LECLERCO, I., 2015. Animal Models for Fibrotic Liver Diseases: What We Have, What We Need, and What Is under Development. J Clin Transl Hepatol. 3(1): 53-66. https://doi.org/10.14218/JCTH.2014.00035
- DELMAS, D., XIAO, J., VEJUX, A., and AIRES, V., 2020. Silymarin and Cancer: A Dual Strategy in Both in Chemoprevention and Chemosensitivity. Molecules, 25(9).https://doi.org/10.3390/molecules25092009
- DI COSTANZO, A., and ANGELICO, R., 2019. Formulation Strategies for Enhancing the Bioavailability of Silymarin: The State of the Art. Molecules.

24(11).https://doi.org/10.3390/molecules24112155

ELBASET, M. A., MOHAMED, B. M. S. A., GAD, S. A., S. ESATBEYOGLU, AFIFI. М., Т., ABDELRAHMAN, S. S., and FAYED, H. M., 2023a. Erythropoietin mitigated thioacetamide-induced renal injury via JAK2/STAT5 and AMPK pathway. Scientific Reports, 13: 14929. https://doi.org/10.1038/s41598-023-42210-1

- ELBASET, M. A., MOHAMED, B. M. S. A., MOUSTAFA, P. E., MANSOUR, D. F., AFIFI, S. M., ESATBEYOGLU, T., ABDELRAHMAN, S. S. M., and FAYED, H. M., 2023b. Erythropoietin Suppresses the Hepatic Fibrosis Caused by Thioacetamide: Role of the PI3K/Akt and TLR4 Signaling Pathways. Oxidative Medicine and Cellular Longevity, 2023: 1-14.https://doi.org/10.1155/2023/5514248
- ELBASET, M. A., NASR, M., IBRAHIM, B. M. M., AHMED-FARID, O. A. H., BAKEER, R. M., HASSAN, N. S., and AHMED, R. F., 2022. Curcumin nanoemulsion counteracts hepatic and cardiac complications associated with high-fat/high-fructose diet in rats. Journal of Food Biochemistry. https://doi.org/10.1111/jfbc.14442
- ELBASET, M. A., SALEM, R. S., AYMAN, F., AYMAN, N., SHABAN, N., AFIFI, S. M., ESATBEYOGLU, T., ABDELAZIZ, M., and ELALFY, Z. S., 2022. Effect of Empagliflozin on Thioacetamide-Induced Liver Injury in Rats: Role of AMPK/SIRT-1/HIF-1α Pathway in Halting Liver Fibrosis. Antioxidants, 11: 2152. <u>https://doi.org/10.3390/antiox11112152</u>
- GILLESSEN, A., and SCHMIDT, H. H., 2020. Silymarin as Supportive Treatment in Liver Diseases: A Narrative Review. Adv Ther, 37(4): 1279-1301. https://doi.org/10.1007/s12325-020-01251-y
- HAJOVSKY, H., HU, G., KOEN, Y., SARMA, D., CUI, W., MOORE, D. S., STAUDINGER, J. L., and HANZLIK, R. P., 2012. Metabolism and toxicity of thioacetamide and thioacetamide S-oxide in rat hepatocytes. Chem Res Toxicol, 25(9): 1955-1963. https://doi.org/10.1021/tx3002719
- HAZEKI, K., NIGORIKAWA, K., and HAZEKI, O., 2007. Role of phosphoinositide 3-kinase in innate immunity. Biol Pharm Bull, 30(9): 1617-1623.<u>https://doi.org/10.1248/bpb.30.1617</u>
- HUSSEIN, R. M., SAWY, D. M., KANDEIL, M. A., and FARGHALY, H. S., 2021. Chlorogenic acid, quercetin, coenzyme Q10 and silymarin modulate Keap1-Nrf2/heme oxygenase-1 signaling in thioacetamideinduced acute liver toxicity. Life Sci, 277: 119460.https://doi.org/10.1016/j.lfs.2021.119460
- JIA, C., ZHANG, Z., WANG, J., and NIE, Z., 2022. Silymarin protects the rats against paraquat-induced acute kidney injury via Nrf2. Human & Experimental Toxicology, 41: 09603271221074334. https://doi.org/10.1177/09603271221074334
- JIAO, F., ZHANG, Z., HU, H., ZHANG, Y., and XIONG, Y., 2022. SIRT6 Activator UBCS039 Inhibits Thioacetamide-Induced Hepatic Injury In Vitro and In Vivo. Front Pharmacol., 13: 837544. https://doi.org/10.3389/fphar.2022.837544
- KIDD, L. B., SCHABBAUER, G. A., LUYENDYK, J. P., HOLSCHER, T. D., TILLEY, R. E., TENCATI, M., and MACKMAN, N., 2008. Insulin activation of the phosphatidylinositol 3-kinase/protein kinase B (Akt) pathway reduces lipopolysaccharide-induced inflammation in mice. J Pharmacol Exp Ther, 326(1): 348-353. https://doi.org/10.1124/jpet.108.138891
- KIM, H. Y., KIM, J. K., CHOI, J. H., JUNG, J. Y., OH,
 W. Y., KIM, D. C., LEE, H. S., KIM, Y. S., KANG,
 S. S., LEE, S. H., and LEE, S. M., 2010. Hepatoprotective effect of pinoresinol on carbon

tetrachloride-induced hepatic damage in mice. J Pharmacol Sci, 112(1): 105-112.https://doi.org/10.1254/jphs.09234FP

- LAMIA, S. S., EMRAN, T., RIKTA, J. K., CHOWDHURY, N. I., SARKER, M., JAIN, P., ISLAM, T., GIAS, Z. T., SHILL, M. C., and REZA, H. M., 2021. Coenzyme Q10 and Silymarin Reduce CCl4-Induced Oxidative Stress and Liver and Kidney Injury in Ovariectomized Rats-Implications for Protective Therapy in Chronic Liver and Kidney Diseases. Pathophysiology, 28(1): 50-63. https://doi.org/10.3390/pathophysiology28010005
- LAWRENCE, T. 2009. The nuclear factor NF-kappaB pathway in inflammation. Cold Spring Harb Perspect Biol, 1(6): <u>https://doi.org/10.1101/cshperspect.a001651</u>
- MANNA, S. K., MUKHOPADHYAY, A., VAN, N. T., and AGGARWAL, B. B., 1999. Silymarin suppresses TNFinduced activation of NF-kappa B, c-Jun N-terminal kinase, and apoptosis. J Immunol, 163(12): 6800-6809. https://doi.org/10.4049/jimmunol.163.12.6800
- MOHAMED, M. Z., and HAFEZ, H. M., 2019. PI3K/Akt and Nrf2/HO-1 pathways involved in the hepatoprotective effect of verapamil against thioacetamide toxicity in rats. 38(4): 381-388. https://doi.org/10.1177/0960327118817099
- MOUSSA, S. A. A., IBRAHIM, F. A. A., ELBASET, M. A., AZIZ, S. W., ABDELLATIF, N. A., ATTIA, A. M., EL TOUMY, S. A., SALIB, J. Y., and BASHANDY, S. A. E., 2022. Efficacy of goldenberry extract in chelated iron overload induced by obesity: Novel safety concept for the treatment of iron overloads diseases. Journal of Applied Biology and Biotechnology, 10: 92-100. https://doi.org/10.7324/JABB.2022.100413
- OGALY, H. A., ABDEL-RAHMAN, R. F., MOHAMED, M. A. E., O, A. A., KHATTAB, M. S., and ABD-ELSALAM, R. M., 2022. Thymol ameliorated neurotoxicity and cognitive deterioration in a thioacetamide-induced hepatic encephalopathy rat model; involvement of the BDNF/CREB signaling pathway. Food Funct, 13(11): 6180-6194.https://doi.org/10.1039/D1FO04292K
- PALLOTTINI, V., MARTINI, C., BASSI, A. M., ROMANO, P., NANNI, G., and TRENTALANCE, A., 2006. Rat HMGCoA reductase activation in thioacetamide-induced liver injury is related to an increased reactive oxygen species content. J Hepatol, 44(2): 368-374. https://doi.org/10.1016/j.jhep.2005.06.011

RAMADAN, A., AFIFI, N., YASSIN, N. Z., ABDEL-RAHMAN, R. F., ABD EL-RAHMAN, S. S., and FAYED, H. M., 2018. Mesalazine, an osteopontin inhibitor: The potential prophylactic and remedial roles in induced liver fibrosis in rats. Chemico-Biological Interactions, 289: 109-

118.<u>https://doi.org/10.1016/j.cbi.2018.05.002</u>

RASLAN, M., ABDEL RAHMAN, R., FAYED, H., OGALY, H., and FIKRY, R., 2021. Metabolomic Profiling of Sansevieria trifasciata hort ex. Prain Leaves and Roots by HPLC-PAD-ESI/MS and its Hepatoprotective Effect via Activation of the NRF2/ARE Signaling Pathway in an Experimentally Induced Liver Fibrosis Rat Model. Egyptian Journal of Chemistry, 64(11): 6647-6671.<u>https://doi.org/10.21608/ejchem.2021.78970.3877</u>

- SALLER, R., MEIER, R., and BRIGNOLI, R., 2001. The use of silymarin in the treatment of liver diseases. Drugs, 61(14): 2035-2063.https://doi.org/10.2165/00003495-200161140-00003
- SARMA, D., and HANZLIK, R. P., 2011. Synthesis of carbon-14, carbon-13 and deuterium labeled forms of thioacetamide and thioacetamide S-oxide. J Labelled Comp Radiopharm, 54(13): 795-798. https://doi.org/10.1002/jlcr.1933
- SHIRANI, K., YOUSEFSANI, B. S., SHIRANI, M., and KARIMI, G., 2020. Protective effects of naringin against drugs and chemical toxins induced hepatotoxicity: A review. 34(8): 1734-1744. <u>https://doi.org/10.1002/ptr.6641</u>
- TALEB, S. A. A., ISMAIL, S. A., MOHAMED, M., MOURAD, R. M., and EL-HASHEMY, H. A., 2023. Promising synthesized bis (arylmethylidene) acetone polymeric PCL emulsified nanoparticles with enhanced antimicrobial/antioxidant efficacy: in-vitro and in-vivo evaluation. OpenNano, 11: 100139. https://doi.org/10.1016/j.onano.2023.100139
- TEKSOY, O., CENGIZ, M., KOROGLU, E., SAHINTÜRK, V., INAL, B., KACAR, S., and AYHANCI, A., 2020. Hepatoprotective effect of silymarin against thioacetamide induced hepatic injury in rats. Applied Biochemistry and Biotechnology, 191:528–539.
- TROUTMAN, T. D., BAZAN, J. F., and PASARE, C., 2012. Toll-like receptors, signaling adapters and regulation of the pro-inflammatory response by PI3K. Cell Cycle, 11(19): 3559-3567.https://doi.org/10.4161/cc.21572
- TSAI, J. H., LIU, J. Y., WU, T. T., HO, P. C., HUANG, C.
 Y., SHYU, J. C., HSIEH, Y. S., TSAI, C. C., and LIU,
 Y. C. 2008. Effects of silymarin on the resolution of liver fibrosis induced by carbon tetrachloride in rats. Journal of Viral Hepatitis, 15(7): 508-514. https://doi.org/10.1111/j.1365-2893.2008.00971.x
- VARGAS-MENDOZA, N., MADRIGAL-SANTILLÁN, E., MORALES-GONZÁLEZ, A., ESQUIVEL-SOTO, J., ESQUIVEL-CHIRINO, C., GARCÍA-LUNA, Y. G.-R. M., GAYOSSO-DE-LUCIO, J. A., and MORALES-GONZÁLEZ, J. A., 2014. Hepatoprotective effect of silymarin. World J Hepatol, 6(3): 144-149. <u>https://doi.org/10.4254/wjh.v6.i3.144</u>
- WANG, W., GUAN, C., SUN, X., ZHAO, Z., LI, J., FU,
 X., QIU, Y., HUANG, M., JIN, J., and HUANG, Z.,
 2016. Tanshinone IIA protects against acetaminopheninduced hepatotoxicity via activating the Nrf2 pathway.

Phytomedicine, 589.https://doi.org/10.1016/j.phymed.2016.02.022

WU, J. W., LIN, L. C., and TSAI, T. H., 2009. Drug-drug interactions of silymarin on the perspective of pharmacokinetics. J Ethnopharmacol, 121(2): 185-193.<u>https://doi.org/10.1016/j.jep.2008.10.036</u>

23(6):

- XU, J., ZHANG, X., MONESTIER, M., ESMON, N. L. and ESMON, C. T., 2011. Extracellular histones are mediators of death through TLR2 and TLR4 in mouse fatal liver injury. J Immunol, 187(5): 2626-2631. https://doi.org/10.4049/jimmunol.1003930
- XU, X., HU, Y., ZHAI, X., LIN, M., CHEN, Z., TIAN, X., ZHANG, F., GAO, D., MA, X., LV, L. and YAO, J., 2013. Salvianolic acid A preconditioning confers protection against concanavalin A-induced liver injury through SIRT1-mediated repression of p66shc in mice. Toxicol Appl Pharmacol, 273(1): 68-76. https://doi.org/10.1016/j.taap.2013.08.021
- ZAHIN, N., ANWAR, R., TEWARI, D., KABIR, M. T., SAJID, A., MATHEW, B., UDDIN, M. S., ALEYA, L. and ABDEL-DAIM, M. M., 2020. Nanoparticles and its biomedical applications in health and diseases: special focus on drug delivery. Environ Sci Pollut Res Int, 27(16): 19151-19168. https://doi.org/10.1007/s11356-019-05211-0
- ZHAI, K.-F., DUAN, H., CAO, W.-G., GAO, G.-Z., SHAN, L.-L., FANG, X.-M. and ZHAO, L., 2016. Protective effect of Rabdosia amethystoides (Benth) Hara extract on acute liver injury induced by Concanavalin A in mice through inhibition of TLR4-NF-κB signaling pathway. Journal of Pharmacological Sciences, 130(2): 94-100. https://doi.org/10.1016/j.jphs.2015.12.006

How to cite this article:

Hany G. Attia, Bassim M.S.A. Mohamed, Saeed M. Alasmari, Sahar S AbdelRahman, Mohamed Aleraky, Abdulwahab Alqahtani, Marawan A. Elbaset and Hany M. Fayed, 2025. Effect of Effervescent Formulation of Silymarin against Experimental Hepatotoxicity in Rats: Involvement of NRF2/HO-1, PI3K/AKT and TLR4/NFκB Pathways. Journal of Applied Veterinary Sciences, 10 (2): 18-27.DOI:https://dx.doi.org/10.21608/javs.2025.348491.1503