

#### ARCHIVES OF AGRICULTURE SCIENCES JOURNAL

Volume 7, Issue 3, 2024, Pages 141-166

Available online at https://aasj.journals.ekb.eg

DOI: https://dx.doi.org/10.21608/aasj.2024.337748.1178

# Improving yield, quality, and storability of Thompson seedless grape by foliar application of potassium

Mansour A. H. A.<sup>a\*</sup>, Omar Nagwa M.<sup>a</sup>, Mohamed Asmaa A.<sup>b</sup>, Aldaby Eman S. E.<sup>c</sup>

<sup>a</sup>Fruit Handling Research Department, Horticultural Research Institute, Agricultural Research Center, Giza, Egypt <sup>b</sup>Viticulture Research Department, Horticulture Research Institute, Agricultural Research Center, Giza, Egypt <sup>c</sup>Botany and Microbiology Department, Faculty of Science, Assiut University, 71516 Assiut, Egypt

#### Abstract

This study was carried out during the two successive seasons 2022 and 2023 on 15-years-old "Thompson seedless grapes" grown at Assiut governorate, Egypt, to study the effect of four foliar spray treatments: control (spraying with water, T1), potassium nitrate (36% K<sub>2</sub>O, T2), potassium citrate (40% K<sub>2</sub>O, T3), and potassium thiosulfate (36% K<sub>2</sub>O, T4) as preharvest on yield, berry quality, storability quality parameters, and enzyme activity of Thompson Seedless Grapes. All vines were sprayed three times during the growing season (end of May, mid-June, and first of July). Treated clusters were stored at  $2\pm2^{\circ}$ C and 85-90% RH for 5 weeks. The results showed that all treatments significantly improved the quantity and quality at harvest of "Thompson Seedless Grapes" potassium citrate or potassium thiosulfate treatment at 0.2% K<sub>2</sub>O increased cluster weight, yield (kg/vine) and yield (ton/ha), berry weight, and berry diameter, while potassium nitrate increased leaf area (cm<sup>2</sup>), juice weight %. Throughout cold storage at  $2\pm2^{\circ}$ C and 85-90% RH for 5 weeks, foliar application of different sources of potassium decreased weight loss %, shattering %, decay %, total loss % and maintained quality and increased antioxidant enzyme activities in the end storage period. The best potassium spray treatments were potassium citrate or potassium thiosulfate to improve the yield and quality at harvest, as well as reduce weight loss, shattering %, decay %, total loss % and the best quality during storage.

Keywords: antioxidant enzymes, grapes, malondialdehyde (MDA), potassium, storage.

\*Corresponding author: Mansour A. H. A., *E-mail address:* dr.mansour2015@yahoo.com



### 1. Introduction

Grapes are of great importance, both in local of consumption terms and exportation. The total area of vineyards in the world reached 11.0 million hectares with a total production of 90.0-million tons (FAO, 2019). In Egypt, it ranks third behind citrus and mango in terms of the cultivated area and the magnitude of the yield. The total area reached was 190,486 feddan, and the fruiting area was 174,715 feddan that produced about 1,594,781 tons of fruits, or about 9.128 tons per feddan. In addition, grapes are considered the most important export horticultural crop, and its export value is about 10% while the quantity is about 3% of total horticultural export (M.A.L.R., 2019). Thompson Seedless grapevines are a popular grape grown in Egypt. Increasing demand for grape production requires a new strategy. However, it is smaller in size. Also, numerous physiological and biochemical processes in the fresh grape, including nutrient intake, increased respiration, and water loss, cause the grape quality to decline during storage (Sabir and Sabir, 2013). Potassium intake is linked to human health. (Cowan et al., 2020). Adequate supply of potassium to crops is essential for the production of high-quality foods (Holland et al., 2019; Pettigrew, 2008; Valentinuzzi et al., 2018). Therefore, adequate supplies of K fertilizer are needed not only to produce foods with sufficient K<sup>+</sup> concentration for a balanced human diet (Holland et al., 2019; Valentinuzzi et al., 2018; Wu et al., 2014), but also to improve fruit quality, as

potassium availability is associated with a rise in concentrations of sugars, soluble solids (Valentinuzzi et al., 2018), and antioxidants such as phenolics (Preciado-Rangel *et al.*, 2018). In crops,  $K^+$ maintains adequate leaf inclination by turgor control (Carroll et al., 1994), enhances photosynthetic assimilation, and improves nutrient uptake (Sustr et al., 2019). Thus, potassium has a key role in the mechanism that controls the transport of nutrients, water, and metabolites across tissues plant and organs. also. maintenance of osmotic homeostasis and plant defense against oxidative stresses (Cuin et al., 2018; Lu et al., 2016; Srivastava et al., 2019). Adequate K nutrition has been associated with increased shelf life, storage, and shipping quality of many horticultural crops (Geraldson, 1985; Lester et al., 2007). Previous studies have shown different effects of different forms of potassium on yield, fruit quality, and post-harvest traits, with different results depending on methods and frequency of application, as well as crop type and stages of fruit growth (Awad et al., 2014; Yousuf et al., 2018). However, no previous studies compared, among K sources, and K source could be applied to grapes. Based on the above, the aim of this paper is to study the physiological and levels of bioactive compounds effects of preharvest foliar application of different sources of potassium: potassium nitrate, potassium sulphate, and potassium citrate's effects on fruit yield, quality, and the storability of Thompson seedless grapes.

## 2. Materials and methods

### 2.1 Experimental site and treatments description

The field trial was conducted during two successive growing seasons, 2022 and 2023, at a private vineyard in Assiut, Egypt. The soil has a clay texture and is well drained, and a surface irrigation system was used to water the tested vineyard. The trial was conducted on a vigorous, fruitful 'Thompson Seedless' (*Vitis vinefera* L.) cultivar with a 15-year-old grapevine and was planted at a distance of 2 m along the row and 2.5 m between the row (2000 vines ha<sup>-1</sup>).

### 2.2 Experimental design

The experiment was performed as a randomized complete block design with three replications, with each replicate consisting of two uniform grapevines. The experiment was set up with four foliar spray treatments: control (spraying with water, T1), potassium nitrate (36% K<sub>2</sub>O, T2), potassium citrate (40% K<sub>2</sub>O, T3), and potassium thiosulfate (36% K<sub>2</sub>O, T4). The treatment solution was sprayed at the rate of 0.2% K<sub>2</sub>O three times during the growing season (end of May, mid-June, and first of July).

### 2.3 Samples collections

At the harvest stage, clusters were collected from each replicate and picked up in the earlier morning and then transported to the Plant Physiology Laboratory, Faculty of Science, Assiut University, Egypt. The samples collected were divided into two groups. The first group consists of two clusters for each replicate to measure the quantity and quality parameters. While the second group was used for storage experiments.

### 2.4 Vegetative growth and yield parameters

The average leaf area (cm<sup>2</sup>) was measured according to the equation reported by Ahmed and Morsy (1999). The sample consisted of 20 leaves for each grapevine from those opposite to basal clusters. At harvest time, the grapevine productivity was recorded (kg/vine). Two randomly selected clusters of each grapevine were selected to measure cluster weight (g), berry weight (g), and berry length and width (mm).

### 2.5 Storage experiment

In this experiment, five storage periods were done, and the duration of each storage period was one week. The harvested clusters were packed in perforated plastic bags and divided into two groups; the first group included two clusters for each replicate/storage period to measure physiological parameters. However, the second group consists of one cluster for each replicate/storage period to perform the biochemical analysis at each storage period. The prepared plastic bags were stored at  $2\pm 2$ °C and 85-90% RH.

### 2.5.1 Physiological parameters

Weight loss (%) = (Initial cluster weight -Final cluster weight / Initial cluster weight)  $\times$  100. The shattered berries were calculated by dividing the shattered berries weight by cluster weight before the shaking and expressed as percentage. Decayed berries were separated, weighted, and expressed as percentage per cluster weight. Total loss in cluster weight percentage was calculated by percentage sum of cluster weight loss, berry shatter, and decayed berries.

### 2.5.2 Biochemical analysis

The following biochemical analyses were performed at harvest time and at each storage period. Soluble solid contents (TSS) of berry juice were measured using the hand refractometer apparatus and expressed as a percentage. Titratable acidity was determined as mg of tartaric acid equivalent using NaOH (0.1 N) in 100 ml of berry juice according to A.O.A.C. (2000). TSS: acid ratio was calculated.

### 2.5.3 Determination of malondialdehyde

Malondialdehyde (MDA), a reactive metabolite of 2-thiobarbituric acid (TBA), was used to measure the amount of lipid peroxidation in the grapes fruit tissues. In short, 2.5 ml of 0.1% w/v trichloroacetic acid (TCA) was used to homogenize 0.45 g of samples. This was followed by 5 minutes of centrifugation at 2400 g. Four milliliters of 20% w/v TCA containing 0.5% w/v TBA were combined with an aliquot of supernatant (1 mL). An ice bath was used to quickly chill the mixture after it had been heated to 95°C for 30 minutes. The mixture was centrifuged for 15 minutes at 2400 g, and the supernatant's absorbance was measured at 450, 532, and 600 nm. The following equation (Hodges *et al.*, 1999) was used to express the MDA concentration as  $\mu$ mol g<sup>-1</sup> DW.

### 2.5.4 Determination of antioxidant enzymes

After homogenizing fresh weight in potassium phosphate buffer (50 mM, pH polyvinylpyrrolidone with and 7) ethylenediamine tetra acetic acid (0.2 mM), centrifugation at 4800 g for 20 minutes was performed. Ascorbate peroxidase, peroxidase, and catalase were all quantified using the supernatant. The Lowry technique was used to determine the extract's protein content (Lowry et al., 1951). The activity of catalase (CAT, EC 1.11.1.6) was determined using Aebi's technique (Aebi, 1984). 100 µL of the enzymatic extract and 2.85 mL of phosphate buffer (PB) (50 mM, pH 7) made up the reaction media. 50  $\mu$ L of 10 mM H<sub>2</sub>O<sub>2</sub> was added to start the reaction, and the absorbance at 240 nm decreased for 60 seconds. The rate at which ascorbic acid is oxidized in a hydrogen peroxidedependent manner is known as ascorbate peroxidase (APX, EC 1.11.1.11) activity (Nakano and Asada, 1987). 2.85 mL of PB (50 mM, pH 7) was combined with 100  $\mu$ l of enzyme extract, H<sub>2</sub>O<sub>2</sub> (100  $\mu$ L, 5 mM), Na<sub>2</sub>-EDTA (0.1 mM), and 144

ascorbic acid (50  $\mu$ L, 0.5 mM) in the reaction mixture. The decrease in absorbance at 290 nm for 60 seconds was used to determine the amount of ascorbic acid that was oxidized. The rate at which guaiacol dehydrogenates at 436 nm was used to calculate the activity of peroxidase (POD) (Pütter, 1974). 3 mL of PB (50 mM, pH 7) was combined with 50  $\mu$ L of guaiacol (20 mM), 30  $\mu$ L of H<sub>2</sub>O<sub>2</sub> (12 mM), and 100  $\mu$ l of enzyme extract in the reaction mixture.

### 2.6 Statistical analysis

To check the significance of differences among treatments, the obtained data of both seasons were subjected to analysis of variance using Statistix 8.1 software (Analytical Software, 2005). Means were compared for significant differences using the Tukey test at P = 0.05. Data were presented as the means  $\pm$  standard deviations.

### 3. Results

### 3.1 Vegetative growth and yield parameters

### 3.1.1 Vegetative growth (Leaf area, cm<sup>2</sup>)

It is clear from the data in Figure (1) that foliar application of different sources of potassium significantly resulted in an obvious promotion on the leaf area ( $cm^2$ ). Similar results were announced during both seasons. The maximum values recorded on vines that sprayed with Knitrate were 203.33 and 206.00 cm<sup>2</sup> during both seasons, respectively. Also, there was no significant difference or effect between spraying grape vines with potassium thiosulfate and potassium citrate in both seasons of the study. On the other hand, the minimum values of these traits were recorded on spraying vine with water (control) at 184.00 and 186.33 cm in the two studied seasons, respectively.

# 3.1.2 Cluster and yield traits: cluster weight (g), yield per vine (kg), and yield per ha (ton)

Figure (2) shows that spraying K-nitrate, K-citrate, or K-thiosulfate significantly with improved the yield expressed in cluster weight (g), yield (kg/vine), and yield (ton/ha) of Thompson Seedless Grapes in the 2022 and 2023 seasons. The experimental results showed the highest values for yield parameters, viz., cluster weight (442 and 451.67 g), yield (kg/vine) (11.49 and 12.20 kg/vine), and yield (ton/ha) (22.98 and 24.39 ton/ha) in vines that were sprayed with K-citrate during the 1<sup>st</sup> and 2<sup>nd</sup> seasons, respectively, compared to other treatments. The lowest cluster weight (348.00 and 350.67 g), yield (kg/vine) (9.05 and 9.47 kg/vine), and yield (ton/ha) (18.10 and 18.94 ton/ha) were recorded in the control in the first and second seasons, respectively. The highest increase percentage of yield (ton/ha) compared to control was observed when grapes vines were treated with Kcitrate, which was 27.01 and 28.80%, then treated with K-thiosulfate, which amounted to (25.48 and 25.76%), followed by treated with K-nitrate (18.20 and 18.63%) in the first and second seasons, respectively.



Figure (1): Influence of foliar application of different sources of potassium on leaf area (cm<sup>2</sup>) of Thompson seedless grapes in the 2022 and 2023 seasons. Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the *P*<0.05 level according to the Tukey test.



Figure (2): Influence of foliar application of different sources of potassium on (A) cluster weight (g), (B) yield (kg/vine), and (C) yield (ton/ha) Thompson seedless grapes in the 2022 and 2023 seasons. Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the *P*<0.05 level according to the Tukey test.

3.1.3 Physical properties of berries (berry weight (g), length (mm), diameter (mm), and juice weight (%)

As shown in the data presented in Figure (3), all treatments increased significantly berry weight, length, diameter, and juice weight % compared to control (water sprayed), The highest significant values were 1.70 and 1.74 g for berry weight and 14.93 and 15.17 mm for berry diameter recorded due to spraying with K-citrate in the two examined seasons, respectively. Likewise, the significantly lowest values of 1.37 and 1.37 g for berry weight and

12.00 and 11.87 cm for berry diameter were recorded for the control of the two seasons, respectively. The lowest values were recorded in berry length (13.57 and 13.47 mm) with unsprayed grape vines. On the other side, spraying K-nitrate (58.77 and 58.40) was significantly preferable to using the other two sources of potassium, followed by K-citrate (58.44 and 56.86). While the lowest treatments for spraying potassium in juice weight are K-thiosulfate (56.20 and 55.57). The lowest values were recorded in juice weight (53.85 and 53.56) in both seasons, respectively.



Figure (3): Influence of foliar application of different sources of potassium on (A) Berry weight (g), (B) length (mm), (C) diameter (mm), and (D) juice weight % of Thompson seedless grapes in the 2022 and 2023 seasons. Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the *P*<0.05 level according to the Tukey test.

# 3.1.4 Chemical properties of berries (TSS, TA, and TSS/TA)

Data listed in Table (1) show that foliar application of sources of potassium was significantly very effective in improving fruit quality in terms of increasing T.S.S. % and TSS/TA and reducing total acidity % in relative to the control treatment. Spraying K-thiosulfate was significantly preferable to using the other two sources of potassium in improving fruit quality. As it gave the highest values TSS/TA (47.03 and 48.68) in both seasons, respectively. On the other hand, the control treatment gave the lowest values of TSS/TA (37.48 and 40.94) in the first and second seasons, respectively.

Table (1): Influence of foliar application of different sources of potassium on (TSS%), acidity (%) and TSS/acid ratio of Thompson seedless grapes in the 2022 and 2023 seasons.

Treatments	(TSS	5 %)	Acidit	y (%)	TSS /Acid ratio		
	Season 2022	Season 2023	Season 2022	Season 2023	Season 2022	Season 2023	
Control	20.47±0.50°	21.27±0.46°	0.55±0.02ª	0.52±0.02ª	37.48±1.98°	40.94±1.77°	
K-nitrate	23.67±0.42ª	24.33±0.58ª	$0.50 \pm 0.0^{b}$	$0.50 \pm 0.01^{ab}$	41.11±6.28 <sup>b</sup>	43.26±5.18 <sup>b</sup>	
K-citrate	22.30±0.10 <sup>b</sup>	22.60±0.20 <sup>b</sup>	$0.42{\pm}0.0^{d}$	0.41±0.01°	44.37±4.30ª	45.96±6.14ª	
K-thiosulfate	23.27±0.64 <sup>ab</sup>	23.80±0.20ª	0.47±0.01°	0.47±0.01 <sup>b</sup>	47.03±1.35 <sup>ab</sup>	48.68±1.49 <sup>b</sup>	

Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the P<0.05 level according to the Tukey test.

### 3.2 Storage period evaluation

### 3.2.1 Weight loss percentage

Data presented in Figure (4) clear a continued increase in weight loss percentage as the cold storage period

increased. As a result, the highest percentage of weight loss % was obtained at the end of the storage period (after five weeks). In Figure (4), clusters treated with different sources of potassium were less in weight loss % than those untreated during different cold storage periods. Moreover, the percentage of weight loss was lower when sprayed with K-citrate. Compared to other potassium spray treatments (10.33 and 9.41% in  $1^{st}$  and  $2^{nd}$  seasons, respectively) at five weeks.



Figure (4): Influence of foliar application of different sources of potassium on weight loss% of Thompson seedless grapes during storage at (2±2 °C) and 85 - 90% RH in 2022 and 2023 seasons. Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the *P*<0.05 level according to the Tukey test.

### 3.2.2 Shatter %

Data presented in Table (2) show a continued increase in shatter percentage as the cold storage period increased. As a result, the highest percentage of shatter % was obtained at the end of the storage period (after five weeks). The highest percentage of shatter % was obtained from the control treatment compared with other treatments, which were 0.93 and 0.64% at 1<sup>st</sup> week, 1.32 and 0.74% at 2<sup>nd</sup> week, 2.46 and 1.85% at 3rd week, 5.65 and 4.36% at 4<sup>th</sup> week; however, at 5<sup>th</sup> weeks, it was 11.19 and 10.52% in both seasons, respectively. From Table (2), the results also show spraying grape vines with potassium nitrate, potassium citrate, and potassium thiosulfate to reduce the shatter percentage during storage at  $2\pm 2$ °C and 85-90 % RH in the 2022 and 2023 seasons. In addition, the difference between spraying treatments was very little significant.

### 3.2.3 Decay %

The effect of foliar application of different sources of potassium on the percentage decay of Thompson seedless grapes at different time samplings during storage was presented in Table (3). In general, decay % was gradually increased with the advancing cold storage periods in both seasons. But no decayed berry was observed in foliar application with Kcitrate or K-thiosulfate in the first week of 2022 season and also in the first and second weeks of the 2023 season. We also note as in Table (3) that all potassium spraying treatments reduce the percentage of decay berries compared to the control during different storage periods. Control treatment recorded the maximum percentage of decay, which were 5.82 and 4.92% during the 2022 and 2023 seasons, respectively, at 5<sup>th</sup> week. Moreover, treatment of K-citrate or

K-thiosulfate showed the highest reduction percentage in decayed percentage during the 2022 and 2023 seasons.

Table (2): Influence of foliar application of different sources of potassium on shatter % of Thompson seedless grapes during storage at  $(2\pm 2 \text{ °C})$  and 85-90% RH in the 2022 and 2023 seasons.

Treatment			2022			2023					
	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	
Control	$0.93{\pm}0.05^{a}$	1.32±0.12 <sup>a</sup>	2.46±0.36 <sup>a</sup>	5.65±0.60 <sup>a</sup>	$11.19{\pm}0.14^{a}$	0.64±0.01 <sup>a</sup>	$0.74{\pm}0.09^{a}$	$1.85{\pm}0.13^{a}$	4.36±0.29 <sup>a</sup>	$10.52{\pm}0.19^{a}$	
K-nitrate	$0.40{\pm}0.07^{b}$	0.93±0.05 <sup>b</sup>	1.59±0.10 <sup>b</sup>	3.93±0.02 <sup>b</sup>	9.97±0.25 <sup>b</sup>	$0.11 \pm 0.20^{b}$	$0.55{\pm}0.09^{ab}$	1.33±0.12 <sup>b</sup>	3.14±0.08 <sup>b</sup>	8.65±0.29 <sup>b</sup>	
K-citrate	$0.47 \pm 0.09^{b}$	0.64±0.10°	$1.05 \pm 0.15^{b}$	2.94±0.50°	9.35±0.25 <sup>b</sup>	0.13±0.11b	0.21±0.03°	1.08±0.12b	2.03±0.44°	8.64±0.27 <sup>b</sup>	
K-thiosulfate	$0.44 \pm 0.07^{b}$	0.73±0.13 <sup>bc</sup>	1.31±0.08 <sup>b</sup>	3.40±0.08 bc	10.22±0.5 <sup>b</sup>	0.27±0.24 <sup>ab</sup>	0.41±0.07 <sup>b</sup>	1.12±0.06 <sup>b</sup>	2.88±0.23 <sup>b</sup>	9.53±0.40 <sup>b</sup>	

Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the P<0.05 level according to the Tukey test.

Table (3): Influence of foliar application of different sources of potassium on decay % of Thompson seedless grapes during storage at  $(2\pm 2 \text{ °C})$  and 85-90% RH in the 2022 and 2023 seasons.

Treatment			2022		2023					
	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks
Control	0.52±0.11ª	1.23±0.09 <sup>a</sup>	$1.87 \pm 0.10^{a}$	3.22±0.12 <sup>a</sup>	5.82±0.85ª	$0.00 \pm 0.00$	$0.80{\pm}0.06^{a}$	1.22±0.15ª	2.77±0.12ª	4.92±0.35 <sup>a</sup>
K-nitrate	0.30±0.01b	$0.62{\pm}0.09^{ab}$	1.25±0.25 <sup>b</sup>	2.25±0.29 <sup>b</sup>	$4.88{\pm}0.07^{ab}$	$0.00 \pm 0.00$	$0.49{\pm}0.06^{b}$	$0.84 \pm 0.10^{b}$	1.91±0.06 <sup>b</sup>	4.36±0.10 <sup>ab</sup>
K-citrate	$0.00{\pm}0.00^{\circ}$	0.33±0.29b	0.55±0.02°	$1.04{\pm}0.10^{d}$	2.97±0.21°	$0.00 \pm 0.00$	$0.00{\pm}0.00^{\circ}$	0.33±0.13°	$0.73{\pm}0.06^{d}$	2.90±0.36°
K-thiosulfate	$0.00 \pm 0.00^{\circ}$	0.39±0.34 <sup>b</sup>	$0.95 \pm 0.14^{bc}$	1.68±0.06°	3.77±0.76 <sup>bc</sup>	$0.00 \pm 0.00$	$0.00{\pm}0.00^{\circ}$	$0.61 \pm 0.10^{bc}$	1.19±0.08°	3.79±0.09 <sup>b</sup>

Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the P<0.05 level according to the Tukey test.

### 3.2.4 Total loss %

Effects of K-nitrate, K-citrate, and Kthiosulfate on total loss were shown in Figure (5). Clusters treated with different sources of potassium were less in total loss % than those untreated during different cold storage periods. Moreover, data also showed that the foliar application of different sources of potassium decreased total loss compared to the water sprayed (control). The best treatment concerned total loss was using K-citrate treatment, which gave the lowest values of 22.65 and 20.95%, followed by K-thiosulfate, which gave 25.08 and 23.12, followed by K-nitrate, which gave 26.13 and 23.26 at the end of the storage period during the 2022 and 2023 seasons, respectively. Data also showed that the total loss increased with the progression of the cold storage period during the two studied seasons. The highest values of total loss were recorded at the end of the storage period.

### 3.2.5 Juice weight %

Data presented in Table (4) show all spraying treatments with potassium reduced the percentage decrease in juice weight compared to the control. The highest rate of decrease in the percentage of juice weight at clusters untreated (control), where the percentage of juice weight decreased from 53.85 and 53.56 to 40.84 and 41.53 at the end of the storage period, *i.e.*, a decrease rate of 24.16 and 22.45%. Wile, the lowest rate of decrease in the percentage of juice weight was at clusters treated with K-citrate, where the percentage of juice weight decreased from 58.44 and 56.86 to 48.25 and 47.70 at the end of the storage period, *i.e.*, a decrease rate of 17.43 and 16.10% compared to the treatments of the other potassium spraying, which decreased rate of 17.88 and 17.44% at clusters treated with K-thiosulfate (18.95 and 17.47%) at clusters treated with K-nitrate after five weeks during the two studied seasons, respectively.



Figure (5): Influence of foliar application of different sources of potassium on Total loss % of Thompson seedless grapes during storage at (2±2 °C) and 85-90% RH in the 2022 and 2023 seasons. Data were presented as means ± SDs, and the different upper letters indicate significant differences at the P<0.05 level according to the Tukey test.

Table (4): Influence of foliar application of different sources of potassium on juice weight % of Thompson seedless grapes during storage at  $2\pm 2$  °C and 85-90% RH in the 2022 and 2023 seasons.

Treatment			2022			2023					
	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	
Control	50.90±0.10°	48.53±0.16 <sup>d</sup>	46.73±0.24°	44.81±0.22 <sup>c</sup>	40.84±0.14 <sup>c</sup>	51.22±0.10 <sup>d</sup>	48.79±0.22 <sup>d</sup>	47.02±0.07 <sup>d</sup>	43.23±0.24 <sup>d</sup>	41.53±0.25 <sup>d</sup>	
K-nitrate	57.05±0.65 <sup>a</sup>	53.72±0.07 <sup>b</sup>	52.12±0.07 <sup>a</sup>	50.22±0.10 <sup>a</sup>	47.64±0.15 <sup>a</sup>	56.43±0.06 <sup>a</sup>	54.21±0.25 <sup>a</sup>	52.21±0.09 <sup>a</sup>	49.91±0.08 <sup>a</sup>	48.20±0.11 <sup>a</sup>	
K-citrate	56.70±0.10 <sup>a</sup>	54.72±0.04ª	52.22±0.07 <sup>a</sup>	50.03±0.07 <sup>a</sup>	48.25±0.05 <sup>a</sup>	54.97±0.04 <sup>b</sup>	52.83±0.10 <sup>b</sup>	51.14±0.14 <sup>b</sup>	49.36±0.05b	47.70±0.28 <sup>b</sup>	
K-thiosulfate	54.14±0.17 <sup>b</sup>	51.47±0.05°	49.22±0.07 <sup>b</sup>	47.75±0.05 <sup>b</sup>	46.15±1.02 <sup>b</sup>	53.73±0.10°	51.43±0.06°	49.24±0.66°	47.75±0.06°	45.88±0.13°	

Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the P<0.05 level according to the Tukey test.

### 3.6 TSS %

Data in Table (5) show the influence of foliar application of different sources of potassium on the content of the total soluble solids during storage at  $2\pm2$  °C and 85-90% RH in the 2022 and 2023 seasons. The percentage of TSS at one, two, three, four, and five weeks during cold storage was measured (Table 5).

Obtained data revealed that TSS was gradually increased with the advancement of cold storage duration in the 2022 and 2023 seasons. The results also show that all potassium spraying treatments had higher values of TSS than the control during different storage periods. The minimum value of TSS was recorded in the juice of Thompson Seedless Grapes untreated (control), which were 20.57, 20.80, 20.97, 21.50, and 22.60 in the 1<sup>st</sup> seasons and 22.07, 22.83, 22.97, 24.83, and 25.20 at one, two, three, four and five weeks in the 2<sup>nd</sup> seasons, respectively.

However, K-nitrate treatment showed the highest TSS content, which were 23.83, 24.03, 24.13, 24.20 and 24.83 in the 1<sup>st</sup> seasons and 24.57, 24.83, 25.43, 25.67, and 26.60 in the 2<sup>nd</sup> seasons at one, two, three, four, and five weeks, respectively. However, the rate of increase in TSS values was higher in the control (9.44 and 15.61%) compared to the rest of the treatments, and the lowest rate of increase was in treatment with K-citrate (3.32 and 6.87%) from harvest until the end of the storage period in the two studied seasons, respectively.

Table (5): Influence of foliar application of different sources of potassium on TSS % of Thompson seedless grapes during storage at  $2\pm2$  °C and 85-90% RH in 2022 and 2023 seasons.

Treatment			2022			2023					
Treautient	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	
Control	20.57±0.06 <sup>d</sup>	20.80±0.10 <sup>d</sup>	20.97±0.06 <sup>d</sup>	21.50±0.50°	22.60±0.20 <sup>d</sup>	22.07±0.50 <sup>b</sup>	22.83±0.15°	22.97±0.06°	24.83±0.15 <sup>ab</sup>	25.20±0.20°	
K-nitrate	23.83±0.06 <sup>a</sup>	24.03±0.15 <sup>a</sup>	24.13±0.12 <sup>a</sup>	24.20±0.10ª	24.83±0.06 <sup>a</sup>	24.57±0.12 <sup>a</sup>	24.83±0.29ª	25.43±0.81 <sup>a</sup>	25.67±0.58 <sup>a</sup>	26.60±0.10 <sup>a</sup>	
K-citrate	22.43±0.06°	22.53±0.15°	22.77±0.15°	22.93±0.21b	23.07±0.15°	23.70±0.26 <sup>a</sup>	23.80±0.20 <sup>b</sup>	24.07±0.12 <sup>bc</sup>	24.17±0.06 <sup>b</sup>	24.27±0.06 <sup>d</sup>	
K-thiosulfate	23.50±0.10 <sup>b</sup>	23.57±0.06b	23.77±0.06b	24.00±.00ª	24.13±0.06b	24.07±0.12 <sup>a</sup>	24.37±0.15 <sup>ab</sup>	24.67±0.15 <sup>ab</sup>	24.80±0.20 <sup>ab</sup>	25.67±0.12b	

Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the P<0.05 level according to the Tukey test.

### 3.2.7 TA %

Data in Figure (6) show the influence of foliar application of different sources of potassium on acidity %. The obtained results indicated that total acidity % (TA %) gradually decreased with increasing storage period during the two studied seasons. The unsprayed clusters (control) recorded the highest total acidity values at the end of storage periods (0.43 and 0.41) compared to the clusters sprayed with potassium, except for the clusters sprayed with K-nitrate (0.44 and 0.43%) in 1<sup>st</sup> and

### 2<sup>nd</sup> seasons, respectively.

### 3.2.8 TSS/acid ratio

Data illustrated in Table (6) show that the TSS/acid ratio increased with an increased storage period. Regarding the effect of foliar spray treatments on the TSS/acid ratio (Table 6), it was significantly affected in both tested seasons. All treatments of foliar application of different sources of potassium result in an increase in values of TSS/TA during different storage periods compared to the control.



Figure (6): Influence of foliar application of different sources of potassium on TA% of Thompson Seedless Grapes during storage at  $2\pm2$  °C and 85-90% RH in the 2022 and 2023 seasons. Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the *P*<0.05 level according to the Tukey test.

Table (6): Influence of foliar application of different sources of potassium on TSS/acid ratio of Thompson seedless grapes during storage at  $2\pm 2$  °C and 85-90% RH in the 2022 and 2023 seasons.

Treatment			2022			2023					
	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	1 weeks	2 weeks	3 weeks	4 weeks	5 weeks	
Control	39.30±0.54°	41.35±1.44 <sup>d</sup>	41.95±0.94°	45.42±0.64°	52.16±0.37°	42.71±1.03 <sup>d</sup>	47.26±1.45°	48.55±1.46 <sup>d</sup>	56.02±1.01°	61.50±1.99 <sup>b</sup>	
K-nitrate	47.99±0.51 <sup>b</sup>	50.09±1.28°	50.99±0.54b	53.39±0.50 <sup>b</sup>	56.90±1.55 <sup>b</sup>	49.80±0.72°	51.38±0.61 <sup>bc</sup>	54.51±2.22°	57.47±1.79 <sup>bc</sup>	61.39±0.62 <sup>b</sup>	
K-citrate	51.78±0.76 <sup>a</sup>	54.53±1.06b	55.99±0.95ª	58.32±1.36 <sup>a</sup>	61.25±1.10 <sup>a</sup>	58.29±1.48 <sup>a</sup>	59.58±2.48 <sup>a</sup>	62.25±0.83 <sup>a</sup>	65.92±0.89 <sup>a</sup>	69.37±1.84 <sup>a</sup>	
K-thiosulfate	53.82±0.92 <sup>a</sup>	56.57±0.65ª	57.51±0.74 <sup>a</sup>	59.51±0.84 <sup>a</sup>	62.42±0.87 <sup>a</sup>	53.50±1.21 <sup>b</sup>	55.81±0.43 <sup>ab</sup>	57.82±1.09 <sup>b</sup>	60.01±1.27 <sup>b</sup>	64.78±2.81 <sup>ab</sup>	

Data were presented as means  $\pm$  SDs, and the different upper letters indicate significant differences at the P<0.05 level according to the Tukey test.

3.2.9 Malondialdehyde (MDA), and specific activities of catalase (CAT), ascorbate peroxidase (APX), peroxidase (POD), and total soluble protein

The MDA level in the grapes was found to be somewhat impacted by the foliar application of various K-salts, although there were no appreciable variations between the various K-salts when compared to the control, which was applied foliar with water. For the seasons 2022 and 2023, the same response was Conversely, (Figure 7). the seen antioxidant enzyme's reaction varies depending on the potassium salt found in the foliar application. When compared to

the control, POD, APX, and CAT (Figure 8, 9 and 10) activities showed noticeably greater values at the various treatments. Over the course of two seasons, the usage of potassium nitrate, citrate. and thiosulfate significantly increased the activities of APX, POD, and CAT. At thiosulfate, nitrate, and citrate, the antioxidant enzymes (ASA and CAT) showed the greatest increase of their activity compared with the control. When compared to the control. foliar administration of K-thiosulfate and Kcitrate salts induced almost the same improved rate of POD activities, which are more than enhanced as detected by Knitrate at two seasons (2022 and 2023) as

follows: (0.041, 0.0497, 0.0559, 0.0571 U/mg protein/g FW/min) at season 2022 and (0.04, 0.048, 0.058, and 0.058 U/mg protein/g FW/min) at season 2023, respectively (Figure 11). During the 2022 season, CAT activities were approximately 48% higher in the foliar containing Kthiosulfate than in the control group (Figure 10). In the first season, potassium salts (K-thiosulfate, nitrate, and citrate) applied topically resulted in increased ASA activities of 35%, 13%, 11%, 28%, 11% and 16% as compared to the control (foliar application with water), respectively (Figure 9). The foliar spray of several K-



Figure (7): Variations in malondialdehyde (MDA) under different potassium salts treatments. Different letters over columns indicate significant differences at the p<0.05.



Figure (9): Specific activities of ascorbate peroxidase (APX) under different potassium salt treatments. Different letters over columns indicate significant differences at p<0.05.

salts improved the protein concentration in the grapes; nevertheless, there were notable variations between the treatments. In the first season of 2022, the rate of protein enhancement was measured at the various K-salts, k-nitrate, k-thiosulfate, and k-citrate in the following order as compared to the control (foliar with water): 12.4, 11.07, 9.4, and 7.6 mg/g Fw, respectively. The responses from the two seasons of 2022 and 2023 were almost identical. Comparing the protein content to the control, potassium nitrate, thiosulfate, and citrate increased it to around 64, 46 and 23.5%, respectively (Figure 11).



Figure (8): Specific activities of guicacol peroxidase (POD) under different potassium salt treatments. Different letters over columns indicate significant differences at p<0.05.



Figure (10): Specific activities of catalase (CAT) under different potassium salt treatments. Different letters over columns indicate significant differences at p<0.05.



Figure (11): Total soluble protein under different potassium salt treatments. Different letters over columns indicate significant differences at p<0.05.

### 4. Discussion

Grapes are considered one of the most significant fruits in the world. Thompson Seedless Grapes is one of the most important varieties of seedless grapes, popular in Egypt. Thompson seedless grape is attracting huge interest for its better. Eating quality and higher returns to the grower, but it is smaller in berry size. Also, the grape quality is reduced during storage due to many physiological and biochemical reactions in the grape, including consumption of nutrients, increased decay. water and loss. Potassium (K) is an essential macronutrient that is known to be required for higher plants. In cell expansion, stomatal movements, and turgor-driven movements, it is essential for maintaining osmoregulation. Additionally, plants that are subjected to a wide range of biotic and abiotic challenges benefit from potassium (Hakerlerler et al., 1997; Johnson et al., 2022; Marschner, 1995; Sanyal et al., 2020; Sardans and Penuelas, 2021; Wang et al., 2013). High K levels were also found to strengthen plant defenses. Potassium has many positive effects on cultivation. In fruit addition. the application of K promotes the increase of fruit weight, fruit diameter, fruit length, and fruit yield. Also, it is improving fruit quality and disease resistance, as well as improving storability. Its role is well documented increasing in enzyme activity, improving the synthesis of carbohydrates, protein, and fats, translocating of photosynthates, enhancing their ability to resist pests and diseases, and also potassium is considered a major osmotically active cation of the plant cell (Mehdi et al., 2007). Adequate K nutrition has been associated with increased fruit yield, fruit weight, fruit size, increased solids soluble and ascorbic acid concentrations. increased shelf life. storage, and shipping quality of many horticultural crops (Geraldson, 1985; Lester et al., 2007). Our results showed spraying that all treatments with potassium led to an increase in leaf area,

and the best treatment in this regard was treatment with potassium nitrate. The increase in leaf area of Thompson Seedless grapes can be attributed to the physiological role of K in carbohydrate formation, which is translocation and accumulation within the organs of the plant and the turgor pressure of cells (Meyer and Anderson, 1970). K also contributes to the growth of meristematic cells, the stimulation of the young tissue, and cell enlargement (Mengel and Kirkby, 1987). In addition to the synergistic effect between potassium and indole acetic acid (IAA) and the enhancement of K on the effect of gibberellin and cytokinin on plant growth (Cocucci and Rosa, 1980). The effect of K in increasing the growth of leaf area was confirmed by Gowda et al. (2022). The obtained results indicated that foliar application of different sources of potassium appeared to have the greatest effect on berry weight, length, diameter, increased cluster weight, and increased yield in comparison to the control trees. Moreover, the maximum cluster weight and yield were produced from grape vines treated with citrate potassium or thiosulfate, followed potassium bv potassium nitrate treatment. An increase in weight of berry with potassium application might be due to the fact that potassium plays an important role in translocation of photo-assimilates to sink organs (fruits), thus resulting in changes to osmosis inside these organs, which allow the continuous import of carbohydrates and water to fruits, leading to higher fruit weight. Another probable explanation for these increases in weight of berries is higher fruit water content (juice weight %) in the potassium treatments. According to our results, it was observed that alteration in fruit water content is closely associated with the fruit's growth development (Haggag et al., 2013). The positive effect of potassium citrate is either due to the role of K in photosynthesis osmoregulation, and allowing an import of the assimilates from source to fruits, which in turn leads to increased fruit weight, or due to the role of citric acid in respiration pathways and the production of important energy (ATP synthesis) for all vital reactions inside the cell (Taiz et al., 2002). The results presented in our research were similar to (Alebidi et al., 2021; Baiea et al., 2015; Ben Mimoun et al., 2013; Bibi et al., 2019; Chidananda et al., 2020; Fattahi et al., 2021; Gowda et al., 2022; Haggag et al., 2016; Jadhav et al., 2019; Jawandha et al., 2017; Shen et al., 2016; Shinde et al., 2018; Solhjoo et al., 2017; Vijay et al., 2017; Yousuf et al., 2018; Zayan et al., 2016), which show that treatment with potassium leads to an increase in fruit weight and yield. Our results also show that all potassium treatments led to an increase in TSS and TSS/TA and a decrease in TA at harvest. It might be because k is known to enhance photophosphorylation and some reactions of photosynthesis, resulting in increased accumulation of carbohydrates. Also, the higher sugar content of fruits might be due to the role of potassium in translocation of sugars from leaves to fruits and maybe be

because of the higher assimilating power of leaves over a long period, resulting in increased availability of sugars in the fruits. The high percentage of sugar in fruits is often the main reason for the increasing percentage of TSS. Generally, all forms of potassium applications resulted in a reduced fruit acidity at harvest in comparison to control. This may be attributed to the role of potassium in neutralizing acids and reducing fruit acidity (Habib, 1998), while the increase in fruit acidity caused by the potassium nitrate's application being a source of nitrogen, which caused an increase in fruit acidity %, as a result of increased synthesis of amino acids, proteins, and other metabolites as well as their subsequent translocation to the fruits (Mohit et al., 2017). These findings are in conformity with those obtained by (Abou El-Wafa et al., 2020; Alebidi et al., 2021; Aly et al., 2020; Baiea et al., 2015; Bibi et al., 2019; Chidananda et al., 2020; Fattahi et al., 2021; Gowda et al., 2022; Haggag et al., 2016; Hamouda et al., 2015; Jadhav et al., 2019; Khayyat et al., 2012; Manivannan et al., 2015; Kumar et al., 2011; Pandey and Singh., 2016; Sindhe et al., 2018; Tehranifar and Tabar, 2009; Yousuf et al., 2018). Regarding the TSS during storage at 2±2 °C and 85-90% RH in the 2022 and 2023 seasons., in most cases, the TSS was increased as the storage period prolonged; this increase may be due to the solubilization of compounds other than carbohydrate to sugars. On the other hand, results showed a decrease in acidity, which was caused by the use of acids in fruits as a source of energy and the conversion of organic acids to sugars (Wills et al., 1998). It may also be due to the loss of water from the berry, and this is consistent with what we found in this study, where the percentage of juice decreases with progress in the storage period. Similar results were seen by several researchers (Jawandha et al., 2017; Nireshkumar et al., 2020; Okba et al., 2021; Prasad et al., 2015; Shen et al., 2016). Also, this study shows that applying several foliar treatments of potassium salts (nitrate, citrate, and thiosulfate) to grape plants has a beneficial effect. This is achieved at all foliar potassium treatments by an increase in the total soluble protein content. The protein content was increased bv potassium nitrate, thiosulfate, and citrate to approximately 64, 46, and 23.5%, respectively, of the control (water foliar). K is responsible for a number of critical functions. biological including the stabilization of protein synthesis, neutralization of negative charges on proteins, and enzyme activation (Ragel et al., 2019; Saadati et al., 2021). Because the synthesis of polypeptides in ribosomes high concentration demands а of potassium, the synthesis of protein increased with the usage of K under stress (Jones and Pollard, 1983; Saadati et al., expansion, stomatal 2021). In cell movements, and turgor-driven movements, it is essential for maintaining osmoregulation (Hakerlerler et al., 1997; Johnson et al., 2022; Marschner, 1995; Sanyal et al., 2020; Sardans and Penuelas,

2021; Wang et al., 2013). According to the current investigation, all K-salts had a neglected impact on grape MDA during the two seasons. This study is compatible with Hernandez et al. (2012), which recorded that potassium treatments enhance plant development while lowering lipid peroxidation. Grapes with K-salts have higher levels of antioxidant enzymes on their foliar. During the first season of 2022, K-thiosulfate had the greatest enhancement of the antioxidant enzymes ASA, POD, and CAT, followed by K-citate and K-nitrate. The availability of K-salts can act as a form of defense against biotic and abiotic stress as well as oxidative damage. It has been discovered that potassium applied foliar improves grapes' antioxidant defense system by lowering reactive oxygen species (ROS) and activating antioxidant enzymes. This result is compatible with the research conducted by García-Martí et al. (2019), which demonstrated that elevated potassium buildup in plant cells can mitigate oxidative stress by increasing the activities of key antioxidant enzymes like oxidase reduction, NADPH dehydroascorbate reductase (DHAR), glutathione reductase (GPX), superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX), peroxidase (POX), and ascorbate reductase (APX). Potassium (K) is an essential macronutrient that plays important roles in plant development, stress adaptation, cotransport of sugars, osmoregulation, and membrane potential regulation (Johnson et al., 2022; Sanyal et al, 2020; Sardans and Penuelas, 2021). According to Johnson et al. (2022), potassium is essential for the upregulation of K+, which lowers the generation of ROS in plants. In these results, K+ lowers the production of ROS through increasing the antioxidant enzymes activities, which lowers lipid peroxidation, which was also recorded by Cakmak (2005) and García-Martí et al. (2019). Foliar application of different sources of potassium reduced the weight loss % and values tended to be statistically equal in most cases. The weight loss is due to the loss of water, and this can be confirmed by looking at the percentage of juice, as it also decreases during the storage period. Earlier studies showed that the pre-harvest potassium salt sprays decreased the weight loss per cent (Nireshkumar et al., 2020; Okba et al., 2021; Venkatachalam, 2015). Regarding the fruit decay % during storage at 2±2 °C and 85-90% RH in 2022 and 2023 seasons., in most cases, all potassium sources recorded a lower percentage of decay than the control treatment, which registered the highest value. Decrease in the percentage of fruit decay in most potassium treatments, which showed significant differences with the control, may be due to their effective role in limiting lipid peroxidation by reducing fruit MDA content, either after harvest or at the end of the storage period (Kanpure et al., 2019; Okba et al., 2021). The positive effect of potassium citrate in decreasing fruit decay % may be attributed to the role of citric acid in minimizing the reduction of totalphenolics

(Abbasi et al., 2013; Venkatachalam et al., 2015) in loquat and longkong fruits, considered anti-microbial which is (Pandey and Rizvi. 2009). and additionally its role in suppressing the activities of oxidoreductase enzymes (PPO and POD) which cause the fruit's surface to become more fragile (Venkatachalam *et al.*, 2015) and susceptible to cold injury, while the positive effect of potassium nitrate on reducing the percentage of decay in Thompson seedless grapes may be through several factors; potassium accumulation in tissues in combination with increase the nitrogen uptake and induction of defense genes. Increasing potassium in tissues, which has a good role in disease resistance by enhanced silicification of cell walls (Prabhu et al., 2007), or due to its direct role in inhibiting the growth of mycelium of the fungus (Sugimoto et al., 2009). Our results are in line with the findings of Kanpure et al. (2019).

# 5. Conclusion

As a whole, it is recommended to spray potassium citrate and potassium thiosulfate in Thompson seedless grapes farms to obtain a high yield and high quality, maintaining the quality of the clusters during storage.

# References

Abbasi, N. A., Akhtar, A., Hussain, A. and Ali, I. (2013), "Effect of antibrowning agents on quality changes of loquat [*Eriobotrya japonica* (Thunb.) Lindley] fruit after harvest", *Pakistan Journal of Botany*, Vol. 45, pp. 1391–1396.

- Abou El-Wafa, M., El-Husseiny, A. M. and El-Soda, A. S. (2020), "Influence of ethrel and potassium thiosulfate spraying on improving fruit coloration and quality of wonderful pomegranate", *Journal of Horticulture Science & Ornamental Plants*, Vol. 12 No. 3, pp. 193–199.
- Aebi, H. (1984), "Catalase *in vitro*", In: *Methods in Enzymology*, Academic Press, USA, pp. 121–126
- Ahmed, F. F. and Morsy, M. H. (1999), "A new method for measuring leaf area in different fruit crops", *Minia Journal of Agriculture Researches & Development*, Vol. 19 No. 6, pp. 97– 105.
- Alebidi, A., Almutairi, K., Merwad, M., Mostafa, E., Saleh, M., Ashour, N., Al Obeed, R. and Elsabagh, A. (2021), "Effect of spraying algae extract and potassium nitrate on the yield and fruit quality of Barhee date palms", *Agronomy*, Vol. 11, Article No. 922.
- Aly, M. A., Harhash, M. M., Bassiony, S.
  S. and Felifal, M. M. S. (2020), "Effect of foliar spray of sitofex, moringa leaves extract and some nutrients on productivity and fruit quality of Thompson seedless grapevine", *Journal of the Advances in Agricultural Researches*, Vol. 25

No. 1, pp. 112–129.

- Analytical software (2005), *Statistix 8.1* for Windows analytical software, Analytical software, Tallahassee, Florida, USA.
- AOAC (2000), Official Methods of Analysis, 17<sup>th</sup> ed., The Association of Official Analytical Chemist (AOAC) International, Washington, DC, USA, pp. 16–20.
- Awad, M. A., Ismail, S. M. and Al-Qurashi, A. D. (2014), "Effect of potassium soil and foliar spray fertilization on yield, fruit quality and nutrient uptake of Seweda date palm grown in sandy loam soil", *Journal of Food, Agriculture and Environment*, Vol. 12, pp. 305–311.
- Baiea, M. H. M., El-Sharony, T. F. and Abd El- Moneim, E. A. A. (2015), "Effect of different forms of potassium on growth, yield and fruit quality of mango cv. Hindi", *International Journal of ChemTech Research*, Vol. 8 No. 4, pp 1581–1587.
- Ben Mimoun, M. and Marchand, M. (2013), "Effects of potassium foliar fertilization on different fruit tree crops over five years of experiments", *Acta Horticulture*, Vol. 984, pp. 211–218.
- Bibi, F., Ahmad, I., Bakhsh, A., Kiran, S.,Danish, S., Ullah, H., and Rehman,A. (2019), "Effect of foliar application of boron with calcium and potassium on quality and yield of

mango cv. Summer Bahisht (SB) Chaunsa", *Open Agriculture*, Vol. 4, pp. 98–106.

- Cakmak, I. (2005), "The role of potassium in alleviating detrimental effects of abiotic stresses in plants", *Journal of Plant Nutrition and Soil Science*, Vol. 168, pp. 521–530.
- Carroll, M. J., Slaughter, L. H. and Krouse, J. M. (1994), "Turgor potential and osmotic constituents of Kentucky Bluegrass leaves supplied with four levels of potassium", *Agronomy Journal*, Vol. 86, Article No. 1079.
- Chidananda, G., Shivakumar, B. S., Salimath, S. B., Girish, R. and Chaitanya, H. S. (2020), "Impact of foliar application of potassium on quality of pomegranate cv. Bhagwa", *International Journal of Current Microbiology and Applied Sciences*, Vol. 9 No. 10, pp. 2885–2893.
- Cocucci, M. C. and Rosa, S. D. (1980), "Effects of canavanine on IAA- and fusicoccin-stimulated cell enlargement, proton extrusion and potassium uptake in maize coleoptiles", *Plant Physiology*, Vol. 48, pp. 239–242.
- Cowan, A. E., Jun, S., Tooze, J. A., Eicher-Miller, H.A., Dodd, K. W., Gahche, J. J., Guenther, P. M., Dwyer, J. T., Potischman, N., Bhadra A. (2020), "Total usual micronutrient intakes compared to the dietary reference intakes among U.S", *Adults by Food Security Status. Nutrients*, Vol. 12, Article No. 38.

- Cuin, T. A., Dreyer, I. and Michard, E. (2018), "The role of potassium channels in *Arabidopsis thaliana* long distance electrical signaling: AKT2 modulates tissue excitability while GORK shapes action potentials", *International Journal of Molecular Sciences*, Vol. 19, Article No. 926.
- Dalal, R. P. S., Vijay and Beniwal, B. S. (2017), "Influence of Foliar Sprays of Different Potassium Fertilizers on Leaf Mineral Quality and Composition Sweet Orange of Jaffa", (Citrus sinensis) cv. International Journal of Pure & Applied Biosciences, Vol. 5 No. 5, pp. 587–594.
- FAO (2019), *Production Yearbook*, Food and Agriculture Organization of the United Nations, Rome, Italy, p. 66.
- Fattahi, E., Jafari, A. and Fallahi, E. (2021), "Foliar Application of Potassium Nitrate and Silicate Can Improve the Fruit Quality of Pomegranate cv. Malase Yazd", *Journal of Plant Productions*, Vol. 43 No. 4, pp. 527–534.
- García-Martí, M., Piñero, M. C., García-Sanchez, F., Mestre, T. C., López-Delacalle, M., Martínez, V. and Rivero, R. M. (2019), "Amelioration of the oxidatice stress generated by simple or combined abiotic stress through the K and ca<sup>+2</sup> supplementation in tomato plants", *Antioxidants*, Vol. 8, Article No. 81.

Geraldson, C. M. (1985), "Potassium

nutrition of vegetable crops", In: Munson RS (ed) *Potassium in Agriculture*, ASA-CSSA-SSSA, Madison, WI, USA, pp. 915–927.

- Gill, P. P. S., Ganaie, M. Y., Dhillon, W. S. and Singh, N. P. (2012), "Effect of foliar sprays of potassium on fruit size and quality of 'Patharnakh' pear", *Indian Journal of Horticulture*, Vol. 69 No. 4, pp. 512–516.
- Gowda, A. M., Ali, H. A. M. and El-Bolok, T. Kh. (2022), "Effect of foliar spraying with different sources of potassium on growth, leaf mineral composition, yield and fruit quality of Picual olive trees", *Egyptian Journal of Horticulture*, Vol. 49, No. 1, pp. 103–114.
- Habib, R. (1998), "Modeling peach fruit acidity nitrogen and potassium nutrition", In *Proceedings of the XXV International Horticultural Congress, Part 2: Mineral Nutrition and Grape and Wine Quality*, Acta 512, Brussels, Belgium.
- Haggag, L. F., Fawzi, M. I. F., Shahin, M.
  F. M. and El-Hady, E. S. (2016), "Effect of yeast, humic acid, fulvic acid, citric acid, potassium citrate and some chelated micro-elements on yield, fruit quality and leaf minerals content of "Canino" apricot trees", *International Journal of ChemTech Research*, Vol. 9, pp. 7–15.
- Haggag, L., Shahin, M. F. M., Genaidy, E.A. E. and Fouad, A. A. (2013),"Changes in fruit weight, dry matter, moisture content and oil percentage

during fruit development stages of two olive cultivars", *Middle East Journal*, Vol. 2, pp. 21–27.

- Hakerlerler, H., Oktay, M., Eryuce, N. and Yagmur, B. (1997), "Effect of potassium sources on the chilling tolerance of some vegetable seedlings grown in hotbeds", In: Johnston, A. E. (Ed.), *Food security in the Wana region, the essential need for balanced fertilization*, International Potash Institute, Basel, Switzerland, pp. 353–359.
- Hamouda H. A., Elham Zinhoum, A. and Zahran Nagwa, G. (2015)."Nutritional status and improving quality fruit by potassium, magnesium and manganese foliar application in pomegranate", International Journal of ChemTech Research, Vol. 8 No. 6, pp. 858-867.
- Hernandez, M., Fernandez-Garcia, N., Garcia-Garma, J., Rubio-Asensio, J.
  S., Rubio, F. and Olmos, E. (2012), "Potassium starvation induces oxidative stress in Solanum lycopersicum L. roots", Journal of Plant Physiology, Vol. 169 No. 14, 1366–1374.
- Hodges, D. M., DeLong, J. M., Forney, C.
  F. and Prange, R. K. (1999), "Improving the thiobarbituric acidreactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds", *Planta*, Vol. 207, pp. 604–611.

Holland, J. E., Hayes, R. C., Refshauge,

G., Poile, G. J., Newell, M. T. and Conyers, M. K. (2019), "Biomass, feed quality, mineral concentration and grain yield responses to potassium fertilizer of dual-purpose crops", *New Zealand Journal of Agricultural Research*, Vol. 62 No. 4, pp. 476–494.

- Jadhav, S., Salvi, V. G., Deshmukh, U. and Jadhav, S. C. (2019), "Yield and quality of Alphonso mango as influence by foliar application of nutrients in lateritic soil", *The Pharma Innovation Journal*, Vol. 8 No. 10, pp. 14–18.
- Jawandha, S. K., Gill, P. P. S., Singh, H. and Thakur, A. (2017), "Effect of potassium nitrate on fruit yield, quality and leaf nutrients content of plum", *Vegetos*, Vol. 30, pp. 325– 328.
- Johnson R., Vishwakarma K., Hossen Md. Sh., Kumar V., Shackira A. M., Puthur J. T., Abdi G., Sarraf M., Hasanuzzaman M. (2022), "Potassium in plants: Growth regulation, signaling, and environmental stress tolerance", *Plant Physiology and Biochemistry*, Vol. 172, pp. 56–69.
- Jones, R. G. and Pollard, A. (1983), "Proteins enzymes and inorganics ions", In: Lauchli, A., Bileski, R. L. (Eds.), *Inorganic Plant Nutrition*, Springer, New York, NY, USA, pp. 528–562.
- Kanpure, R. N., Tiwari, R., Barholia, A. K., Kushwah, S. S., Mishra, S. N. and Lekhi, R. (2019), "Effect of pre

harvest spray of gibberellic acid, calcium nitrate and potassium sulphate on post-harvest behaviour of winter guava (*Psidium guajava* L.) fruits", *Journal of Pharmacognosy and Phytochemistry*, Vol. 8 No. 3, pp. 3955–3959.

- Khayyat, M., Tehranifar, A., Zaree, M., Karimian, Z., Aminifard, M. H., Vazifeshenas, M. R., Amini, S., Noori, Y. and Shakeri, M. (2012), "Effects of potassium nitrate spraying on fruit characteristics of "malas yazdi" pomegranate", *Journal of Plant Nutrition*, Vol. 35, pp. 1387–1393.
- Kumar, C. N., Sathyanarayana, B. N., Naresh, P. and Lakshmipathy, M. (2011), "Effect of certain pre harvest treatments in improving the yield and quality of banana cv. Nanjangudu Rasabale", *Plant Archives*, Vol. 11 No. 2, pp. 677–681.
- Lester, G. E., Jifon, J. L. and Stewart, W. M. (2007), "Foliar potassium improves cantaloupe marketable and nutritional quality", *Better Crops*, Vol. 91, pp. 24–25.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. and Randall, R. J. (1951), "Protein measurement with the Folin phenol reagent", *Journal of Biological Chemistry*, Vol. 193, pp. 265–275.
- Lu, Z. F., Lu, J. W., Pan, Y. H., Li, X. K., Cong, R. H. and Ren, T. (2016), "Genotypic variation in photosynthetic limitation responses to K deficiency of Brassica napus is

associated with potassium utilization efficiency", *Functional of Plant Biology*, Vol. 42, pp. 880–891.

- M.A.L.R. (2019), *Ministry of Agriculture* and Land Reclamation Publishes, Economic Affairs Sector, Ministry of Agriculture and Land Reclamation, Egypt.
- Mandal, G., Dhaliwal, H. S. and Mahajan, B. V. C. (2012), "Effect of preharvest application of NAA and potassium nitrate on storage quality of winter guava (*Psidium guajava*)", *Indian Journal of Agricultural Sciences*, Vol. 82 No. 1, Article No. 985.
- Manivannan, M. I., Irulandi, S. and Thingalmaniyan, K. S. (2015), "Studies on the effect of pre-harvest application of plant growth regulators and chemicals on yield and quality of guava (*Psidium guajava* L.) cv. L-49", *International Journal* of Agriculture Science, Vol. 11 No. 1, pp. 138–140.
- Marschner, H. (1995), Mineral Nutrition of Higher Plants, 2<sup>nd</sup> ed., Academic Press, San Diego, California, USA.
- Mehdi, S. M., Sarfraz, M. and Hafeez, M. (2007), "Response of rice advance line PB-95 to potassium application in saline sodic soil", *Pakistan Journal of Biological Science*, Vol. 10, pp. 2935–2939.
- Mengel, K. and Kirkby, E. A. (1987), *Principle of Plant Nutrition*, 4<sup>th</sup> ed., International Potash Institute, Berne,

Switzerland.

- Meyer, B. S. and Anderson, D. B. I. (1970), *Plant Physiology*, 2<sup>nd</sup> ed., East West Press Pvt. Ltd., New Delhi, India, pp. 473.
- Mohit, M., Verma, M. L. and Thakur, J. (2017), "Effect of different nitrogenous fertilizers on fruit quality yield of apricot (Prunus and L.)", Journal armeniaca of Pharmacognosy and Phytochemistry, Vol. 6, pp. 217–220.
- Nakano, Y. and Asada, K. (1987), "Purification of ascorbate peroxidase in spinach chloroplasts; its inactivation in ascorbate-depleted medium and reactivation by monodehydroascorbate radical", *Plant and Cell Physiology*, Vol. 28, pp. 131–140.
- Nireshkumar, N., Saraswathy, S., Subbiah, A. and Venkatesan, K. "Yield (2020),and quality enhancement as influenced by preharvest nutrient spray in grapes (Vitis *vinifera L.*) var. Muscat Hamburg by adopting double pruning and double cropping system under Cumbum condition", Valley Journal of Pharmacognosy and Phytochemistry, Vol. 9 No. 5, pp. 2240–2246.
- Okba, S. K., Mazrou, Y., Elmenofy, H. M., Ezzat, A. and Salama, A. (2021), "New insights of potassium sources impacts as foliar application on 'Canino' apricot fruit yield, fruit anatomy, quality and storability", *Plants*, Vol. 10, Article No. 1163.

- Pandey, D. and Singh, N. P. (2016), "Effect of foliar feeding of potassium nitrate on yield and fruit quality parameters of litchi cv. Dehradun", *International Journal and Chemical Studies*, Vol. 6 No. 6, pp. 662–666.
- Pandey, K. B. and Rizvi, S. I. (2009), "Plant polyphenols as dietary antioxidants in human health and disease", *Oxidative Medicine and Cellular Longevity*, Vol. 2 No. 5, p. 270–278.
- Pettigrew, W. T. (2008), "Potassium influences on yield and quality production for maize, wheat, soybean and cotton", *Plant Physiology*, Vol. 133, pp. 670–681.
- Prabhu, A. M. S., Fageria, N. K., Huber,
  D. M. and Rodrigues, F. A. (2007),
  "Potassium and Plant Disease", In Mineral Nutrition and Plant Disease,
  American Phytopathological Society Press, USA, pp. 57–78.
- Prasad, B., Dimri, D. C. and Bora, L. (2015), "Effect of pre-harvest foliar spray of calcium and potassium on fruit quality of Pear cv. Pathernakh", *Scientific Research and Essays*, Vol. 10 No. 11, p. 376–380.
- Preciado-Rangel, P., Salas-Pérez, L., Gallegos-Robles, M. A., Ruiz-Espinoza, F. H., Ayala-Garay, A. V., Fortis-Hernández, M. and Murillo-Amador, B. (2018), "Increasing doses of potassium increases yield and quality of muskmelon fruits under greenhouse", *Horticultera Brasilieria*, Vol. 36, pp. 184–188.

- Pütter, J. (1974), "Peroxidases", In: Methods of enzymatic analysis, Elsevier, USA, pp. 685–690
- Ragel P., Raddatz N., Leidi E. O., Quintero F. J., Pardo J. M. (2019), "Regulation of K nutrition in plants", *Frontiers in Plant Science*, Vol. 10, Article No. 281.
- Ruiz, R. (2006), "Effects of different potassium fertilizers on yield, fruit quality and nutritional status of "Fairlane" nectarine trees and on soil fertility", *Acta Horticulturae*, Vol. 721, pp. 185–190.
- Saadati, S., Baninasab, B., Mobli, M. and Gholami, M. (2019), "Measurements of freezing tolerance and their relationship with some biochemical and physiological parameters in seven olive cultivarsc", *Acta Physiologiae Plantarum*, Vol. 41, Article No. 51.
- Saadati, S., Baninasab, B., Mobli, M. and Gholami, M. (2021), "Foliar application of potassium to improve the freezing tolerance of olive leaves by increasing some osmolite compounds and antioxidant activity", *Scientia Horticulturae*, Vol. 276, Article No. 109765.
- Sabir, F. K. and Sabir, A. (2013), "Quality response of table grapes (*Vitis vinifera* L.) during cold storage to postharvest cap stem excision and hot water treatments", *International Journal of Food Science & Technology*, Vol. 48, pp. 999–1006.

- Sardans, J. and Peñuelas, J. (2021), "Potassium control of plant functions: ecological and agricultural implications", *Plants*, Vol. 10, Article No. 419.
- Shen, C., Ding, Y., Lei, X., Zhao, P., Wang, S., Xu and Y., Dong, C. (2016), "Effects of foliar potassium fertilization on fruit growth rate, potassium accumulation, yield, and quality of 'Kousui' Japanese pear", *Hort technology*, Vol. 26, pp. 270– 277.
- Shinde, A. K., Dabke, D. J., Keluskar, M. H. and Bhave, S. G. (2018), "Effect of different doses and sources of potassium on yield, spongy tissue and nutrient content of Alphonso mango", Advanced Agricultural Research & Technology Journal, Vol. 2 No. 2, pp. 2581–3749.
- Solhjoo, S., Gharaghani, A. and Fallahi, E. (2017), "Calcium and potassium foliar sprays affect fruit skin color, quality attributes, and mineral nutrient concentrations of 'Red Delicious' apples", *International Journal of Fruit Science*, Vol. 17, pp. 358–373.
- Srivastava, A. K., Shankar, A., Chandran, A. K. N., Sharma, M., Jung, K. H., Suprasanna, P. and Pandey, G. K. (2019), "Emerging concepts of potassium homeostasis in plants", *Journal of Expermintal Botany*, Vol. 71, pp. 608–619.
- Sugimoto, T., Watanabe, K., Furiki, M., Walker, D. R., Yoshida, S., Aino, M.,

Kanto, T. and Irie, K. (2009), "The effect of potassium nitrate on the reduction of phytophthora stem rot disease of soybeans, the growth rate and zoospore release of phytophthora sojae", *Journal of Phytopathology*, Vol. 157, pp. 379–389.

- Sustr, M., Soukup, A. and Tylova, E. (2019), "Potassium in root growth and development", *Plants*, Vol. 8, pp. 435.
- Taiz, L. and Zeiger, E. (2002), "Respiration and Lipid Metabolism", *In Plant Physiology*, 3<sup>rd</sup> ed., Sinauer Associates, Sunderland, MA, USA, pp. 223–258.
- Tehranifar, A. and Tabar, S. M. (2009), "Foliar application of potassium and boron during pomegranate fruit development can improve fruit quality", *Horticulture, Environment, and Biotechnology*, Vol. 50 No. 3, pp. 1-6.
- Valentinuzzi, F., Maver, M., Fontanari, S., Mott, D., Savini, G., Tiziani, R., Pii, Y., Mimmo, T. and Cesco, S. (2018), "Foliar application of potassium-based fertilizer improves strawberry fruit quality", In VIII International Symposium on Mineral Nutrition of Fruit Crops, Vol. 1217, pp. 379–384.
- Venkatachalam, K. (2015), "The different concentrations of citric acid on inhibition of Longkong pericarp browning during low temperature storage", *International Journal of Fruit Science*, Vol. 15, pp. 353–368.

- Vijay, V., Dalal, R. P. S., Beniwal, B. S. and Saini, H. (2017), "Effect of foliar application of potassium and its spray schedule on yield and yield parameters of sweet orange (*Citrus* sinensis Osbeck) cv. Jaffa", Journal of Applied and Natural Science, Vol. 9 No. 2, pp. 786–790.
- Wang, M., Zheng, Q., Shen, Q. and Guo, S. (2013), "The critical role of potassium in plant stress response", *International Journal of Molecular Sciences*, Vol. 14 No. 4, pp. 7370–7390.
- Wills, R. B. H., Me Glasson, W. B., Graham, D. and Joyce, D. (1998), Post-harvest Introduction to the physiology and handling of fruits. vegetable and ornamentals, 4<sup>th</sup> New South Wales University Press Limited, Kensington, England.
- Wu, L. Q., Cui, Z. L., Chen, X. P., Zhao, R. F., Si, D. X., Sun, Y. X. and Yue, S. C. (2014), "High-Yield maize production in relation to potassium uptake requirements in China", *Agronomy Journal*, Vol. 106 No. 4, pp. 1153–1158.
- Yang, C., Chen, T., Shen, B., Sun, S., Song, H., Chen, D. and Xi, W. (2019), "Citric acid treatment reduces decay and maintains the postharvest quality of peach (*Prunus persica* L.) fruit", *Food Science & Nutrition*, Vol. 7 No. 11, pp. 3635–3643.
- Yousuf, S., Sheikh, M. A., Chand, S. and Anjum, J. (2018), "Effect of different

sources of potassium on yield and quality of apple (cv. Red Delicious) in temperate conditions", *Journal of Applied and Natural Science*, Vol. 10 No. 4, pp. 1332–1340.

Zayan, M. A., Mikhael, G. B. and Okba,

S. K. (2016), "Treatments for improving tree growth, yield and fruit quality and for reducing double fruit and deep suture incidence in "Desert red" peach trees", *International Journal of Horticultural Science*, Vol. 22 No. 3-4, pp. 7–19.