

EFFECT OF POTASSIUM FERTILIZATION ON ANNA APPLE TREES GROWN IN SANDY SOILS OF EGYPT

EMAN. S. ATTALA

Horticultural Research Institute, Agricultural Research Centre, Giza . Egypt.

(Manuscript received 18 February 1997)

Abstract

The effect of applying different doses from 0 to 2 kgs of potassium sulphate/tree, divided in four applications, was investigated on Anna apple trees on M.M. 106 stock grown at newly reclaimed sandy soils under a drip irrigation system. No significant effect in leaf N and P content occurred in response to fertilizer addition yet, K, Fe, Zn and Mg content were increased by application of 1.5 or 2.0 Kgs of K_2SO_4 . Likewise, obvious increases in leaf total soluble and non soluble carbohydrates were evident. Significant increase in fruit set and decrease in fruit drop were also noticed. Fruit weight, volume, flesh firmness and T.S.S. values were increased by 1.5 and 2.0 Kgs rate of the fertilizer rendering higher values of T.S.S. /acidity that caused earlier fruit ripening. However, acid content was not affected. Yield was highly increased by the high rates of potassium sulphate additions.

INTRODUCTION

Production of low chilling apple cultivars has spread extensively in the newly reclaimed sand soils in Egypt during the last decades. Many difficulties including nutritional problems, have been met by the growers. Optimum rates and time of application of major elements are not exactly determined. However, it has been mentioned that the optimum concentration of major mineral elements in mature apple leaves were around: N 1.8-2.4%, P 0.08-0.30, K 1.3-1.8%, Ca 1.0-1.3%, Mg 0.35 -0.50%, S 2%, Zn 35-50 p.p.m, Mn 35-50 p.p.m, Cu 7.5 -20 p.p.m, Fe 50 p.p.m, B 35-50 p.p.m (Swietlikand Miklos, 1984).

It has been reported that increasing potassium application resulted in only slight variation in vegetative and reproductive development of Gravensteiner apples (Schumacher Stadler, 1991).

The highest average yield of Goldenspur apples was obtained from plots receiving N, P and K (as potassium chloride) each at 60 kgs/ha. (Zhigarevich and Treglzoza, 1991). However, it has been determined that 540 kg K_2O /ha was the optimum fertilization rate. Best results in terms of growth, yield and fruit stability of Goldenspur apple were produced when 60 kgs N + 60 kgs P + 60 kgs K/ha were applied (Treglzoza and Fataliev, 1994).

In an experiment carried out on young Anna apple trees growing in newly reclaimed sand soil at El-Nubaria Egypt, soil application of (450 or 900g K_2SO_4 /tree) in 3 equal split doses: March 10th, May 1st and June 1st resulted in significant increase of yield, fruit weight, fruit size and fruit firmness (Kilany and Kilany, 1991).

Effect of potassium application on mineral leaf content of apples was reviewed by Hansen (1975) and citrus by Resse and Koo (1975) respectively. The aim of this work is to determine the optimum K requirements for young Anna apple trees grown in newly reclaimed sandy soil.

MATERIALS AND METHODS

This investigation was carried out during two successive seasons, i.e 1994 and 1995 on young Anna apple trees on M.M. 106 stock of seven and eight years old planted 5m x 5m apart (168 trees/Feddan). Trees are grown in sand soil of El-Kasassen experimental orchard. Drip irrigation, using 3000 M^3 of water/feddan/year, is practiced. Analyses of soil and water before and after treatments are presented in Table 1.

The trees were equally subjected to the same cultural and fertilization practices (except for K). fifteen M^3 /fed. of cow manure + 200 kg super phosphate (15.5% P_2O_5) were applied/feddan in shallow trenches dug 50 cm away from the trunks. Nitrogen fertilization was carried out by adding 210 units in December of Nitrogen as Ammonium nitrate (33.5%) per feddan.

The experiment included four treatments of potassium fertilization by adding 0.0, 0.75, 1.5 and 2.0 Kgs of potassium sulphate/tree (K_2O) divided into 4 split

doses (March, April, May and June) and added to the soil surface under the irrigation drippers. Each treatment comprised 4 replicates each of 3 individual trees. Guard rows of untreated trees surrounded each replicate. The following determinations were carried on the vegetative growth in the two seasons of investigation. Samples of leaves were taken from medium portion of current season growth in July according to the method described by Westwood, 1978. Leaves were dried at 70°C for 72 hrs. The following analyses were carried on dry weight bases. Total nitrogen, phosphorus and potassium were determined according to the method described by Koch and Meekin (1924), Troug and Meyer, (1939), and Brown and Lilleland (1946), respectively Also, Fe, Zn and Mn were determined using the atomic absorption apparatus model (Thermo Jarrell Ash smith-Hieftje 1000) (Barrows and Simpson, 1962). Total carbohydrates were analysed by the method of Smith and Dubois, (1956). Percentage of fruit set and shedding was calculated. Samples were taken from various replicates when control fruits attained maturity (Abd El-Aziz et al, 1985).

Table 1. Soil and irrigation water analysis.

Samples	PH	soluble salts p.p.m	NO3 (%)	P (%)	K (%)	Na (%)	Cl- (%)	SO4- (%)	Hco- (%)
Before Application	7.98	1664	1.29	0.08	1.62	12.2	12.3	14.33	3.93
After Application									
Control 0.0 kg/tree	7.72	1671	1.28	0.09	1.60	12.0	12.9	14.9	3.59
0.75 Kg/tree	7.70	1680	1.26	0.08	1.71	12.1	13.03	14.2	3.41
1.5 Kg/tree	7.63	1690	1.29	0.08	1.73	11.92	12.9	14.9	3.39
2.0 Kg/tree	7.69	1681	1.29	0.09	1.72	12.21	13.1	14.3	3.51
Irrigation Water	7.60	1088	0.93	0.07	1.67	11.4	8.47	6.2	3.28

Individual fruit weight and diameter, firmness (Lb/cm²) using penetrometer (fruit tester) trade mark effegi with plunger of 5/16 inch and T.S.S. was determined using a hand refractometer (ATAGO No1 Brix 0 32%) made in Japan.

The strach content was estimated by the method of Lane and Eynon, 1960 and Whelan, 1955. Total number of fruits and yield/tree were estimated. Data were subjected to analysis of variance using the complete randomized design and averages were compared by L.S.D. (05) (Snedecor and Cochran, 1967).

RESULTS AND DISCUSSION

Vegetative Growth :

Mineral content of mature leaves on dry weight bases are presented in Table (2). It is evident that neither nitrogen nor phosphorous, in the leaves, were affected by potassium application in both seasons. However, potassium percentage was increased by the 1.5 and 2.0 Kgs soil application. Concerning Fe, Zn and Mn, they were increased by potassium application especially by any of the conducted potassium treatments. Leaves total soluble and non soluble carbohydrates were increased by higher concentrations of potassium especially at 2.0 Kgs. The effect of potassium

Table 2. Effect of Potassium Application on Anna Apple Leaves Nutrients and Carbohydrates.

K2O/tree (kgs)	Leaf elements concentrations						Dry weigh (%)	Carbohydrate		
	N (%)	P (%)	K (%)	Fe (%)	Zn (%)	Mn (%)		Soluble (%)	Non Soluble (%)	Total (%)
Season 1994										
Control	1.90	0.11	0.49	69	35	82	41.31	3.20	5.75	8.95
0.75	1.93	0.12	0.79	72	52	120	41.91	6.95	1.70	8.65
1.50	1.92	0.12	1.16	80	80	179	41.82	6.75	1.95	8.70
2.00	1.99	0.14	1.26	83	90	191	41.73	6.20	3.70	9.90
L.S.D 0.5	N.S.	N.S.	0.35	4.5.	6.0	5.0	N.S.	1.2	2.0	0.5
Season 1995										
Control	1.82	0.12	0.60	75	38	89	41.29	4.20	4.40	8.60
0.75	1.99	0.13	0.89	96	62	190	41.70	5.40	3.50	8.90
1.50	1.92	0.14	1.09	99	93	183	41.62	5.55	3.50	9.05
2.00	1.99	0.13	1.13	98	100	200	41.50	6.55	2.40	8.95
L.S.D 0.5	N.S	N.S	0.30	5.0	7.1	5.2	N.S	0.8	1.8	0.22

fertilization on leaf mineral content has been previously discussed. in the absence of K, leaf K content declined to 0.9% in bearing apple trees (Klein, 1992). Leaf content of "Cox's orange pippin" potassium was increased by soil application of potassium (Hansen, 1975). Nevertheless, the effect of potassium application on leaf mineral content has been extensively studied in many citrus cultivars. Many reports empha-

sized a significant increase (Koo and Reese, 1973, Reese and Koo, 1976 and Fahmy and Hassaballa, 1977. Badra and Yousif, 1978, Bakr et al, 1980 and Sarooshi et al, 1991). Concerning nitrogen content, some reports pointed out that it was reduced by K application (Takidze, 1971), increased (Badra and Yousif, 1978) or had no effect (Jones et al, 1973a). No effect had been attributed to K fertilization on P leaf content (Takidze, 1971 and Bazeleth, 1980). Ca became lower in leaves by K application (Takidze, 1971 Fahmy and Hassaballa, 1977). Less Mg content in leaves has been reported by Reese and Koo, 1976 and Badra and Yousif, 1978. However, Cu and Zn have been increased (Badra and Yousif, 1978) while Fe and Mn decreased (Kazak and Khalidy, 1974) by potassium application.

Fruit Set and Drop :

It is apparent that K application resulted in considerable increase in fruit set (Table 3).

Table 3. Effect of Potassium Application on Physical Apple fruit Characteristics and yield.

Treatments K ₂ O/tree (kgs)	Fruit Set (%)	Fruit drop (%)	Fruit No./tree (g) (g)	Yield tree (Kgs)
Season 1994				
Control	9.01	26.2	189	15.69
0.75	23.7	14.2	290	29.96
1.50	34.1	10.2	289	34.42
2.00	36.2	8.3	291	34.92
L.S.D 0.5	2.1	4.5	9.1	1.8
Season 1995				
Control	8.01	24.3	182	16.63
0.75	18.0	13.2	299	33.79
1.50	36.1	11.4	295	36.29
2.00	38.2	8.0	290	35.38
L.S.D 0.5	3.4	5.9	8.3	1.5

In the two seasons of the investigation. Both 1.5 and 2 kgs of K₂O were superior in this respect Fruit drop was also decreased by Potassium application. Differ-

ences were not constant between the 3 used concentrations. Potassium application enhanced fruit set (Wang et al, 1994). However non-significant variation in reproductive development of apple trees could be related to K fertilization (Schumacher and Stadler, 1991).

Mature fruits quality :

Fruit characters were compared when reached maturity (Abd El-Azia et al, 1985). Both fruit weight and volume were increased by Potassium application (Table 4). The increase was more evident with the high concentrations of Potassium.

Table 4. Effect of Potassium Application on Anna Apple Fruit Characteristics.

Treatments K ₂ O/tree (kgs)	Fruit Weight (g)	Fruit Volume (cm) ³	Fruit dimension		Firmness (16/cm ₂)
			Length cm	Width cm	
Season 1994					
Control	83.0	99	6.3	5.7	12.3
0.75	103.3	121	7.4	6.8	12.5
1.50	119.1	183	7.9	6.8	12.4
2.00	120.0	200	8.1	7.2	13.1
L.S.D 0.5	7.1	5.1	1.2	1.1	0.2
Season 1995					
Control	93	101	6.7	5.9	11.5
0.75	113	162	6.9	6.8	12.0
1.50	123	220	7.5	7.1	12.2
2.00	122	210	7.9	7.0	12.8
L.S.D 0.5	6.8	6.2	0.5	1.2	0.4

Fruit dimensions were enlarged except those of the equatorial diameter in the second season. Potassium application resulted in firmer fruits. This was more evident with the higher concentrations of Potassium in the first year of study. T.S.S. in the Juice was increased by Potassium application by 1.5 and 2.0 kgs KNO₃ in the first season and only kgs in the second one.

Acidity was not changed significantly by any of the conducted treatments in the two seasons of the investigation (Table 5).

Table 5. Effect of Potassium Application on some Anna Fruit Chemical.

K ₂ O/tree (kgs)	T.S.S. (%)	Acidity (%)	T.S.S. acid ratio	Ascorbic acid (mg/100g)	Starch (%)	Total sugars (mg/g D.W.)
Season 1994						
Control	11.3	0.90	12.56	13.0	3.5	175
0.75	12.0	0.80	15.00	13.7	3.1	200
1.50	12.9	0.80	16.13	13.9	3.1	210
2.00	14.2	0.79	17.97	14.5	3.0	225
L.S.D. 0.5	1.20	N.S	1.50	1.5	0.2	4.5
Season 1995						
Control	11.9	0.80	14.88	12.0	3.4	180
0.75	12.5	0.81	15.43	13.2	3.3	205
1.50	13.8	0.73	18.90	13.9	3.2	221
2.00	15.0	0.62	24.16	14.2	3.1	232
L.S.D. 0.5	2.0	N.S	1.8	2.0	0.2	4.9

T.S.S./ acid ratio was increased by potassium application especially with 1.5 and 2.0 kgs from K₂SO₄ that achieved, higher values as compared with control. Such increment has enhanced fruit maturity dramatically. Nevertheless, the rate of 750 kgs did not result in any apparent effect in the second season.

Ascorbic acid and starch contents were increased by potassium fertilization especially at the rate of 2.0 kgs. However, this rate was more effective than the other rates in the second season. Soluble total sugar content were increased by potassium application.

This was more apparent at the two kgs rate of application in both seasons and in the 1.5 kgs in the second only.

Previous studies confirmed that significant increase of fruit weight size and

firmness were induced by 450 or 900 gm. K_2SO_4 , (Kilany and Kilany, 1991). Fruit quality, however, was improved by K application (Wang et al, 1994).

Yield :

Number of mature fruits per tree was not changed by potassium application in the first season, however, it was increased significantly in the second (Table 3). Total yield was increased by potassium in the two years of the experiment. Differences attributed to different rates of applications were not apparent. Previous studies confirmed that significant increase of yield was induced by 450 or 900 K_2SO_4 /tree (Kilany and Kilany, 1991) .

Other investigations reported increase in the yield of apple fruits by K application (Zhigarevich and Treglazova, 1991 Klein, 1992, Solv'Eva et al, 1992, and Treglazova and Fataliev, 1994).

Conclusions :

Conclusively, the present results indicate that adding K_2SO_4 at 1.5 or 2.0 Kgs to young Anna apple trees under drip irrigation has its positive impact on yield and fruit quality allowing apple growers having similar conditions to adjust their fertilizer rates and frequencies more properly.

REFERENCES

1. Abd El-Aziz, E., I. Saied and G.R. Stino. 1985. Studies on performance of "Anna" apple trees on the two Malling Merton Stocks in Egypt. Bull, Fac. of Agric., Univ. of Cairo 36 (2) : 1337-47 .
2. Badra, T and M.G. Yousif. 1978. Comparative effects of potassium levels on growth and mineral composition of intact and nematized cowpea and sour orange seedlings . Nematologia Mediterranea, 7 (11 21-27). Soils and Fertilizers No. 43 : 7772.
3. Bakr, E.L., R.M. Barakat, A.M. Shahan and I.S. Salem. 1980. Effect of Potassium Fertilization on leaf nutrient content of valencia orange trees grown in sandy soils Arid Lands Development Abstracts, No. 3:2020.
4. Barrows, H.L. and E.C. Simpson. 1962. An EDTA method for the direct routine determinations of calcium and magnesium in soil and plant tissue. Soil Sci: Soc Amer, proc., 26:443-445 .
5. Bazeleth M.1980. Potassium Fertilization experiment in a shamout, orange grove. Israel-Agricultural Research Organization, Volcani Center, pamphlet (No. 220) 28 PD. Arid Lands Development Abstracts, No. 3:2025 .
6. Brown, I.O. and O.Lilleland. 1946. Rapid determination of Potassium and Sodium in plant material and soil extracts by Flamephotometry - Proc. Amer. Soc. Hort. Soc. Hort. Sci, 48 : 341-346 .
7. Fahmy, I. and A.I. Hassaballa. 1977. The effect of the different rootstocks and potassium sulfate fertilizer on leaf potassium, calcium and sodium in young clementine mandarin trees. Libyan Journal of Agriculture 6 (1) 1-5 Soil and Fertilizers No. 41 : 6734 .
8. Hansen, P. 1975. Effect of potassium and magnesium on "cox's orange pippin" on a sandy loam. Tidsskrift for plantearb 79 (2) 259-265. Soils and Fertilizers No. 38:5262.
9. Jones W.W., T.W. Embleton, H. J. foot and G.R. Platt. 1973. Response of young lemon trees to potassium and zinc application, yield and fruit quality. Soils and Fertilizers No, 37: 367 .
10. Kazak, M. and R. Khalidy. 1974. The effect to seven combinations of nitrogen, phosphorus and potassium fertilizer on leaf composition of Shammouti and Valencia oranges in South and North Lebanon. Idem 61-66. Hort. Abst. Vol. 46:1590.

11. Kilany, A.E., O.A. Kilany. 1991. Effect of potassium and boron nutrients on growth, yield and fruit quality of Anna apple trees. Bulletin of Faculty of Agriculture, Univesity of Cairo 42 (2) 415-428 .
12. Klein 1. 1992. The effect of potassium applied by drip irrigation on growth, fruit quality and storage of apples on a soil with marginal Mg content. Potash Review subject 9, 5th suite, No. 1,7 pp.
13. Koch and Mc Mee-Kin, 1924. Determination of total nitrogen by Nislar solution. J. Amer. Chem-soc. vol. 46 : 2066, 13.
14. Koo, R.C.J. and L.R. Reese. 1973. A comparison of Potash Sources and rates for citures. Potash Review 8/24. 7pp. Soil and Fertilizers. Hort. Abst. No. 37:1373 .
15. Lane, H.M. and S.I. Eynon. 1960. Analysis of fruit and vegetable products. Reducing and total sugar determinations published by British Crop production, 4th Edition (1960) 9-13.
16. Reese, R.L. and J.C.R. Koo. 1975. N and K fertilization effects on leaf analysis tree size and yield of three major florida orange cultivars. J. Amer. Soc. Hort. Sci., 100 (2), 195-198.
17. Reese, R.L. and J.C.R. Koo. 1976. Influence of fertility and irrigation on yield and leaf and soil analyses of (Temple) Orange proceedings of the Florida. Hort. Abst. Vol. 48-859.
18. Sarooshi, R., G.R. Weir and G.B. Coote. 1991. Effect of nitrogen, phosphorus and potassium fertilizer rates on fruit yield, leaf mineral concentration and growth of young orange trees in the sunraysia district. Australian journal of Experimental Agriculture 31 (2) 263-272. (Hort. Abst. Vol. 62:8270).
19. Schumacher, R., W. Stadler. 1991. Nitrogen and Potassium trial with Gravensteiner Stickstoff-Und Kaliumversuch bei Gravensteiner. Schweizerisch Zeitschrift fur obst-und weinbau 127 (12) 307-310.
20. Smith, F. and M. Dubois. 1956. Coloremtric method for determination of sugar and related substances. Anal, chem. 29 : 350.
21. Snedecor, C. W. and C.W. Cochran. 1967. Statistical Methods. Iowa state University press 6 th ed. Amer. Iowa USA.
22. Solv'EVA, M.A., B.I. Bershtein, N.F. Movchan. 1992. Effect of potassium fertilizers on the productivity and frost resistance of fruit trees.

- Sel'skokhozyaistvennaya Biologiya No. 3, 115-127.
23. Swietlik, D. and Miklos Foust. 1984. Foliar nutrition of fruit crops. Hort. Rev. 6: 287-355. Av., Publ. Co., Westport. Ct.
 24. Takidze, R. 1971. The diagnosis of Mandrian potassium nutrition subtropicheske. Kul Juvy, (6), 52-56. (Hort. Abst. Vol. 42:8349).
 25. Treglazova N.V., A.T. Fetaliev. 1994. Effect of combined mineral top dressing on apple storability. Sadovodstvo i vinogradarstvo No. 4, II {Ru}. Kuba, 37:3171 Azerbaijan.
 26. Troug, E. and A.H. Meyer. 1939. Improvement in deiness colorimetric method for phosphorus and arsenic. Ind. Eng. Chem. Anal. Ed. 136-139.
 27. Wang., J.K., Z.Zhang., G.Y. Wang, F.S. Guan., J.S., Feng. 1994. A study on the effects of different types of mineral fertilizer on apple trees. Journal of Fruit Science (1994) II (4) 240-241.
 28. Westwood, M.N. 1978. Temperate-Zone pomology. W.H. Freeman Comany San Fransisco. USA. pp. 428.
 29. Whelan, W.G. 1955. Starchighy cogen fructosans and similar polysaccharides spimger veriage P. 145-196.
 30. Zhigarevich, I. A., N.V. Treglazova. 1991. Combined mineral nutrition of apple trees Sadovodstv i vinogradastvo No. 10, 14-16 (Ru) Aznpo sisk, Kuba, Azerbaijan.

تأثير التسميد البوتاسى على اشجار التفاح صنف أنا النامية فى الأراضى الرملية فى مصر

ايمان صبحى عطا الله

معهد بحوث البساتين مركز البحوث الزراعية - الجيزة .

تمت دراسة تأثير إضافة معدلات مختلفة من كبريتا البوتاسيوم من صفر الى ٢ كجم/شجرة مقسمة على أربع دفعات على اشجار التفاح صنف الأنا المطعومة على أصل م.م. ١٠٦ والنامية فى الاراضى الرملية الحديثة الاستصلاح والتي تروى بنظام الرى بالتنقيط لم تكن هناك تأثيرات معنوية على محتوى الأوراق من عنصرى النيتروجين والفوسفور نتيجة إضافة السماد فقد زاد المحتوى من عناصر البوتاسيوم والحديد والزنك والمغنسيوم عند إضافة ١,٥ أو ٢ كجم من السماد للشجرة. وبالمثل كانت هناك زيادات واضحة فى الكربوهيدرات الكلية الذائبة والغير ذائبة كما كانت هناك زيادة معنوية فى عقد الثمار ونقص فى تساقطها، أدت إضافة ١,٥ كجم أو ٢ كجم من السماد الى حدوث زيادة فى وزن وحجم الثمار وصلابة اللحم ومحتواه من المواد الصلبة الذائبة الكلية مما أدى إلى زيادة نسبة المواد الصلبة الذائبة الى الحموضة مما سبب تبكير نضج الثمار لكن المحتوى من الحموضة لم يتغير .

أدت اضافة المعدلات العالية من كبريتات البوتاسيوم إلى حدوث زيادة معنوية فى المحصول.